

Simple Method for Standardization and Quantification of Linoleic Acid in *Solanum nigrum* Berries by HPTLC

Arpan Chakraborty¹, Arka Bhattacharjee², Pallab Dasgupta³, Debasmitta Manna⁴, Won Chun OH⁵ and Goutam Mukhopadhyay^{3*}

¹Department of Pharmaceutical Technology, Jadavpur University, Kolkata, West Bengal, India

²Department of Pharmaceutical Sciences and Technology, Birla Institute of Technology, Mesra, Ranchi, India

³BCDA College of Pharmacy and Technology, Kolkata, West Bengal, India

⁴NSHM College of Pharmaceutical Sciences, Kolkata, West Bengal, India

⁵Department of Material Science and Engineering, Hanseo University, South Korea

Abstract

Linoleic acid is a polyunsaturated omega-6 fatty acid found as phytoconstituent in many plant species. It is reported to have activity such as 5-alpha-reductase-inhibitor, antianaphylactic, antiarthritic, antiatherosclerotic, anticancer, anticoronary, antieczemic, antifibrinolytic and many others. In this research work we have developed a method by HPTLC to identify and standardize linoleic acid in methanolic extract of *Solanum nigrum* berries. Calibration curve of Linoleic acid was plotted and was found to have r^2 value of 0.99451. The percentage content of Linoleic acid in *Solanum nigrum* berries methanolic extract was found to be 9.32% w/w.

Keywords: Linoleic acid; *Solanum nigrum*; Finger print; HPTLC

Abbreviations: HPTLC: High Performance Thin Layer Chromatography; μg : Microgram; mL: Mililitre; R_f : Retention Factor; nm: Nanometer; μL : Microlitre; RSD: Relative Standard Deviation; TLC: Thin Layer Chromatography; °C: Degree Centigrade.

Introduction

Solanum nigrum (black nightshade) is a medicinal plant member belonging to the family Solanaceae. *S. nigrum* has been extensively used traditionally to treat various ailments. The juice of the berries used as an antiarrhoea, ophthalmopathy and hydrophobia. It is also used in anasarca and heart disease. Berries are reported to possess tonic, diuretic and cathartic properties. Seeds are useful in dysuria and giddiness [1].

Linoleic acid is a polyunsaturated omega-6 fatty acid. It is reported to have a number of useful physiological activities including 5-alpha-reductase-inhibitor, antianaphylactic, antiarthritic, antiatherosclerotic, anticoronary, anticancer, antieczemic, antifibrinolytic, antigranular, antihistaminic, anti-inflammatory, antimenorrhagic, antiprosthetic, hepatoprotective, hypocholesterolemic, immunomodulatory, insectifuge, metastatic, nematicide [2]. *Solanum nigrum* oil is reported to have linoleic acid as the most abundant unsaturated fatty acid found in it [3,4].

For the establishment of a consistent biological activity of any herbal item, standardization is an important step. It provides a consistent chemical profile, or simply a quality assurance program for production and manufacturing of an herbal drug. Standardization thus, is a tool in the quality control process. HPTLC is a modern adaptation of Thin Layer Chromatography with better and advanced separation efficiency and detection limits. HPTLC provides a number of advantages including easy separation process for with colored compounds. HPTLC can be used for different modes of evaluation, allowing identification of compounds having different light-absorption [5]. HPTLC is a method to standardize and identify the chemical ingredients which is expected to be present in a medicinal plant. This is done from regulatory perspective to ensure the efficacy, quality as well safety of the herbal drugs present in a plant. Thus it provides a very reliable way of determining the purity and percentage content of the active biomarker in the plant extracts.

Standardization of herbal products is a current issue of interest.

For quality control of these herbal materials or their extracts one needs to proceed by selecting one of the different phytoconstituents of the product, preferably the one showing maximum desired bioactivity and subsequent method of quantification of that specific constituent is required to be developed. The method so accepted should be simple and cost effective. Previously mentioned standardization methods of *Solanum nigrum* berries states use of gas chromatography, high performance liquid chromatography which by quantifying linoleic acid or other phytoconstituents [6,7]. But these methods are expensive. Standardization and quantification of linoleic acid in *Solanum nigrum* berries by HPTLC for quality control of this herbal material is an effective and simple method and may be used with necessary modifications for estimation of the a fore said phytoconstituent in other plant materials containing the same.

Materials and Methods

Instrumentation and reagents

The CAMAG HPTLC system used comprised of WINCATS software, LINOMAT V automatic sample applicator, and automatic development chamber, scanning densitometer CAMAG scanner 3 and photo documentation apparatus CAMAG reprostar-3. Stationary phase was used as aluminum based silica gel plate 60 F₂₅₄ (Merck, Mumbai) with 10 cm × 10 cm in a particle size of 5-10 μm . All the solvents were used of analytical grade. 100 μL syringe (Hamilton, Switzerland) was used for sample application on HPTLC plates. Linoleic acid was purchased from Sisco Research Laboratories (SRL). Methanol, n-hexane and ethyl acetate (analytical grade) were procured from Merck (Mumbai, India).

***Corresponding author:** Goutam Mukhopadhyay, BCDA College of Pharmacy and Technology, Kolkata, West Bengal-700 127, India, Tel: +919836333563; E-mail: gmkhopadhyay8@gmail.com

Received November 25, 2016; Accepted December 01, 2016; Published December 07, 2016

Citation: Chakraborty A, Bhattacharjee A, Dasgupta P, Manna D, Chun WOH, et al. (2016) Simple Method for Standardization and Quantification of Linoleic Acid in *Solanum nigrum* Berries by HPTLC. J Chromatogr Sep Tech 7: 342. doi: 10.4172/2157-7064.1000342

Copyright: © 2016 Chakraborty A, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

All the samples were filtered through Whatman's syringe filter (NYL 0.45 μm). The plant material was collected from local area of Kolkata, West Bengal, India and the identity was confirmed by taxonomist. The shade dried powdered berries of *Solanum nigrum* were extracted with methanol by cold maceration. The extract solution was dried under reduced pressure with the help of rotary evaporator.

Methods

Preparation of standard solution: About 1 mg of linoleic acid standard was taken in 1 mL eppendorf tube 1.0 mL of methanol was added to it and mixed in vortex mixture till the material got completely dissolved. It was then filtered through 0.45 μm syringe filter and kept for further study.

Preparation of calibration curve of linoleic acid: HPTLC analysis was performed using isocratic technique. The mobile phase was optimized with n-hexane and ethyl acetate in a ratio of 5:4 v/v. The temperature was kept at 25°C and mobile phase was developed in a twin trough glass chamber. The standard solution was applied 2, 4, 6, 8, 10 μL in a band wise fashion. After development the plate was dried. Then the dry plate was treated by spraying sulphuric acid-anisaldehyde reagent. The plate was kept at 110°C for 5 minute in hot air oven and evaluation was carried out at 366 and 540 nm. Calibration curve of linoleic acid was obtained by plotting peak areas versus concentrations of linoleic acid applied.

Identification and quantification of linoleic acid in *Solanum nigrum* berries

Preparation of sample solution: About 5 mg of *Solanum nigrum* berries methanolic extract was dissolved in 1 mL methanol in an eppendorf tubes. It was then mixed in vortex mixture and subjected to ultrasonication bath till the material completely dissolved. Then it was filtered through 0.45 μm syringe filter. Sample solution was applied consequently in the range of 4, 8 and 12 μL and was subjected to a fore mentioned chromatography.

Evaluation of method by some parameters

Instrumental precision: Instrumental precision was checked by repeated scanning (n=5) of the same spot of linoleic acid (2 $\mu\text{g/spot}$) and expressed as relative standard deviation (%RSD).

Repeatability: The repeatability of the method was affirmed by analyzing 2 $\mu\text{g/spot}$ of linoleic acid TLC plate (n=5) and expressed as %RSD.

Recovery: The accuracy of the method was assessed by performing recovery study at three different levels (50%, 100% and 125% addition of linoleic acid). The percent recoveries and the average percent recoveries were calculated.

Results

Calibration curve of linoleic acid

The calibration curve was found to be linear with the equation of $Y=3141.508 \times X+1366.840$ (correlation coefficient=0.9954), where X represents amount of linoleic acid and Y represents area under the curve. R_f value of standard linoleic acid was found to be 0.65.

Quantification of linoleic acid in *Solanum nigrum* berries

The percentage content of linoleic acid in *Solanum nigrum* berries methanolic extract was found to be 9.32% w/w. R_f value of standard linoleic acid was found to be 0.65. Specificity was confirmed by



comparing the R_f of standard and sample. Figures 1 and 2 represents HPTLC Chromatogram of standard Linoleic acid and that of extract of *Solanum nigrum* berries. Photodocumentation of *Solanum nigrum* berries methanolic extract at 366 nm and 540 nm are shown in Figures 3 and 4. Figure 5 represents all tracks at 540 nm having 5 standard and three samples respectively from left hand side.

Evaluation of method by some parameters

Instrumental precision was within range of 1.97-2.08 with% RSD of 1.65% whereas repeatability was found to be in range of 1.98-2.08 with % RSD of 1.95% percentage recovery data is reported in Table 1. with standard deviation of the amount of marker found.

Discussion

Chromatographic fingerprint analysis has proven to be a rational and feasible approach for the assessment of quality and authentication of species of traditional medicine [8-10]. It effectively uses chromatographic techniques to form specific patterns of recognition for phytochemicals. The developed fingerprint pattern of components can thereafter be utilized to check the presence of markers of interest as well as the ratio of all detectable analytes. Though there are some shortcomings of high performance thin layer chromatography, such as






Figure 3: HPTLC Fingerprint analysis of *Solanum nigrum* berries methanolic extract in UV 366 nm. Track 3a, 3b, 3c, 3d and 3e: Linoleic acid standard solution at 2, 4, 6, 8 and 10 μ L concentration respectively. Track 3sa, 3sb and 3sc: Sample solution at 4, 8 and 12 μ L concentration respectively.




Figure 4: HPTLC Fingerprint analysis of *Solanum nigrum* berries methanolic extract in UV 540 nm. Track 4a, 4b, 4c, 4d and 4e: Linoleic acid standard solution at 2, 4, 6, 8 and 10 μ L concentrations respectively. Track 4sa, 4sb and 4sc: Sample solution at 4, 8 and 12 μ L concentration respectively.

the limited developing distance and lower plate efficiency as compared to high performance liquid chromatography and gas chromatography, still it is an effective tool for evaluation of herbal drugs due to its simplicity. Moreover, the formerly mentioned limitations can be curbed by separately developing fractions of different polarity on two or several thin layer plates. The unique feature of the picture like image of HPTLC




Figure 5: HPTLC 3D Chromatogram of *Solanum nigrum* berries methanolic extract in UV 540 nm. Track 5a, 5b, 5c, 5d and 5e: Linoleic acid standard solution at 2, 4, 6, 8 and 10 μ L concentrations respectively. Track 5sa, 5sb and 5sc: Sample solution at 4, 8 and 12 μ L concentration respectively.

Marker	Amount of marker present (μ g)	Amount of marker added (μ g)	Amount of marker found (μ g)	Recovery (%)	Average recovery (%)
Linoleic acid	5	2.5	7.56 \pm 1.11	100.13	100.63
	5	5	10.08 \pm 1.05	100.86	
	5	6.25	11.35 \pm 1.04	100.91	

Table 1: Recovery data of linoleic acid.

coupled with the digital scanning profile is an attractive and useful tool for construction of herbal chromatographic fingerprint. The linoleic acid in sample of extract was identified and its presence was confirmed by comparing the R_f value of standard linoleic acid to that of the extract. This method is reproducible and has shown satisfactory results on precision, accuracy and recovery study data. There is no report of detection and quantification of linoleic acid in *Solanum nigrum* berries by HPTLC. Hence, we developed a simple and precise method for quantification of this marker.

Conclusion

The developed method provides a simple, precise and accurate analytical method for the identification and quantification of linoleic acid in berries of *Solanum nigrum*. Quick and easier process for sample preparation, high method sensitivity and reproducible results by the mobile phase are the important features of the developed method. It can be expected that this HPTLC technique may be applied successfully for evaluation of linoleic acid as marker in different herbal extracts.

References

1. Khattak JZK, Anwar Z, Aftab S, Afzal M, Islam M, et al. (2012) Solenium nigrum as potent therapy: A Review. British Journal of Pharmacology and Toxicology 3: 185-189.
2. Duke JA, Bogenschutz-Godwin MJ, Du-Cellier J, Duke PK (2000) CRC handbook of medicinal spices. CRC Press LLC, NW.
3. Sarma H, Sarma A (2011) Solanum nigrum L., a nutraceutical enriched herb or invasive weed?. International Conference on Environment and BioScience IPCBEE 21: 105-109.
4. Saleem TMS, Chetty CM, Ramkanth S, Alagusundaram M, Gnanaprakash K, et al. (2009) Solanum nigrum Linn.- A Review. Pharmacognosy review 3: 342-345.
5. Shivatare RS, Nagore DH, Nipanika SU (2013) HPTLC' an important tool in standardization of herbal medical product: A Review. Journal of Scientific and Innovative Research 2: 1086-1096.

6. Bai Y, Lin CH, Luo WM, Wang LY, Sun C, et al. (2012) Simultaneous quantification of four saponins, three alkaloids and three fatty acids in *Solanum nigrum* Linn. by HPLC-ELSD. Journal of Medicinal Plants Research 6: 3632-3639.
7. Xie P, Chen S, Liang Y, Wang X, Tian R, et al. (2006) Chromatographic fingerprint analysis-a rational approach for quality assessment of traditional Chinese herbal medicine. Journal of Chromatography A 1112: 171-180.
8. Qiao C, Han Q, Song J, Mo S, Kong L, et al. (2007) Chemical fingerprint and quantitative analysis of Fructus Psoraleae by high-performance liquid chromatography. Journal of Separation Science 30: 813-818.
9. Lu H, Liang Y, Chen S (2006) Identification and quality assessment of *Houttuynia cordata* injection using GC-MS fingerprint: a standardization approach. Journal of Ethnopharmacology 105: 436-440.
10. Li K, Wang S (2005) Fingerprint chromatogram analysis of extracts from the leaves of *Tripterygium wilfordii* Hook. F. by high performance liquid chromatography. Journal of Separation Science 28: 653-657.

Citation: Chakraborty A, Bhattacharjee A, Dasgupta P, Manna D, Chun WOH, et al. (2016) Simple Method for Standardization and Quantification of Linoleic Acid in *Solanum nigrum* Berries by HPTLC. J Chromatogr Sep Tech 7: 342. doi: [10.4172/2157-7064.1000342](https://doi.org/10.4172/2157-7064.1000342)

OMICS International: Open Access Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission>