

Smoking *Cannabis* is Especially Dangerous for Youth Diagnosed with Attention Deficit/Hyperactivity Disorder (ADHD)

Roger W Apple^{1*}, Donald E Greydanus¹ and Joav Merrick²⁻⁶

¹Department of Pediatric and Adolescent Medicine, Western Michigan University Homer Stryker MD School of Medicine, Kalamazoo, Michigan, USA

²National Institute of Child Health and Human Development, Jerusalem

³Office of the Medical Director, Health Services, Division for Intellectual and Developmental Disabilities, Ministry of Social Affairs and Social Services, Jerusalem

⁴Division of Pediatrics, Hadassah Hebrew University Medical Center, Mt Scopus Campus, Jerusalem, Israel

⁵Kentucky Children's Hospital, University of Kentucky School of Medicine, Lexington, Kentucky, USA

⁶Center for Healthy Development, School of Public Health, Georgia State University, Atlanta, USA

*Corresponding author: Apple RW, PhD, Assistant Professor, Department of Pediatric and Adolescent Medicine, Western Michigan University Homer Stryker MD School of Medicine, 1000 Oakland Drive, Kalamazoo, MI 49008-1284, United States. E-mail: Roger.Apple@med.wmich.edu

Received date: June 28, 2016; Accepted date: July 09, 2016; Published date: July 13, 2016

Copyright: © 2016 Apple RW, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Cannabis and Neurological Functioning

Lifetime prevalence of *cannabis* use is associated with multiple health and psychosocial risk factors and is greater than for all other illicit drugs [1]. For adolescents, this fact, combined with the symptoms associated with ADHD, makes *cannabis* use especially dangerous. Even before considering the impact of ADHD it is important to consider neurological changes that have been identified in adolescents who use *cannabis* compared to those who have not [2-17]. Neurological changes associated with chronic use of *cannabis* by adolescents include: excessive stimulation of cannabinoid receptors that has been shown to interfere with normal pruning of synapses during adolescence [7], asymmetrical increase in the size of hippocampi and amygdalae, enlargement of cerebellum, and thinning of the frontal cortex [9,10]. Regular *cannabis* use at early ages has also been shown to affect normal grey and white matter development, impaired axonal connectivity, and changes in cerebral blood flow [2,3,8,11,18,19].

Structural and functional brain differences have been shown in adolescents who use *cannabis* on a regular basis including inattention, impulsivity, impaired executive function, memory loss, decreased coordination, distorted visuospatial perception, altered awareness of passage of time, decline in intelligence quotient (IQ), and impaired novel learning [3-6]. Other studies suggest that the prevalence of *cannabis* abuse is increased in those with attention deficit hyperactivity disorder (ADHD) versus that found in the general population though the complexities of this relationship require more research [17,20-23]. This opinion is not trying to resolve the issue of whether or not those with ADHD are self-medicating or if the impulsive symptoms of ADHD contributed to *cannabis* use, but rather, that *cannabis* smoking combined with youth diagnosed with ADHD is dangerous.

Heritability of *cannabis* use among adolescents has also been discussed as problematic. One study reported that genetic factors were responsible for 40% of individual differences in *cannabis* initiation for adolescents but also acknowledged that environmental influences do increase with age [24]. This supports that a genetic predisposition exists for early initiation which could subsequently lead to the exacerbation of psychosocial and health risk factors [1]. Due to the vulnerability of these risk factors during adolescence, efforts at prevention during and before the adolescence developmental period, is paramount.

Cannabis and ADHD

Linking *cannabis* and ADHD is widely described in the literature [22,23,25,26]. One study found that 34% of adolescents receiving inpatient treatment for problems related to drug dependence were diagnosed with ADHD [22] and another found that males with ADHD and conduct disorder are at an increased risk for early initiation of substance use (including *cannabis*) [23]. With *cannabis* use and ADHD often co-occurring it is not surprising that many diagnosed with ADHD may turn to *cannabis* for self-medication according to some research [25]. *Cannabis* use is also associated with poorer levels of adolescent decision making, conduct disorder, and risky sexual behavior [26] all of which could negatively impact symptoms associated with ADHD. There is widespread belief that *cannabis* use improves mental health symptoms such as anxiety. While there may be empirical support for such claims it is important to note that *cannabis* is not a drug that will improve all aspects of mental health functioning, in fact, it can exacerbate problems such as cocaine-induced paranoia and other psychotic symptoms [27,28]. All of these factors clearly illustrate the hazards for youth diagnosed with ADHD should they decide to smoke *cannabis*.

Family influences cannot be ignored when discussing the dangers of *cannabis* use for youth diagnosed with ADHD [29-31]. Family factors found to contribute to *cannabis* use among youth include: permissiveness toward drug use, perception of parental control, social ties to parents, coercive family interaction, family integration, family stress, family income, adolescent disclosure, family activities, parental solicitation, parental knowledge of drug use, and parental monitoring [32-36]. Another study found additional family and peer influences in predicting *cannabis* use among adolescents including drinking to intoxication, peers who have experience with drugs, frequency of father's alcohol intoxication, and aggressive behavior with initiation of smoking by age of 12 as the most powerful predictor of *cannabis* use [37]. It should go without question that combining these factors along with problems with attention, concentration, impulsivity, and hyperactivity are a recipe for disaster.

Providers should understand that considerable research has shown that the developing brain, with its high neuronal plasticity, is vulnerable to exposure to exogenous cannabinoids, particularly during the adolescence developmental period [2-17]. Research also suggests that early initiation of *cannabis* use can increase risks for cognition

dysfunction, CNS changes, neuropsychiatric disorders, *cannabis* dependence, and consumption of additional illicit drugs [15,26]. All of these factors combined with the symptoms of ADHD should be of great concern to treating providers and awareness of this dangerous combination of factors could lead to improved identification, earlier treatment, and better outcomes.

Care Managers

Treatment and intervention services for youth diagnosed with ADHD who are also using *cannabis*, and may also be experiencing a myriad of psychosocial stressors, often proves exceedingly difficult; particularly if the physician is practicing independently with little or no ancillary staff. In an excellent study conducted by Ciccone, et al. [38] the introduction of care managers was evaluated in regard to health care management for patients with heart disease and diabetes. While this is a very different population than the one being discussed, the concept of care managers could potentially provide significant help in treatment and intervention. The results of their study found that utilizing care managers helped patients take a more active role in their treatment, in fact, making the patient the center of treatment. They also found care managers helped create partnerships among physicians and patients and worked to empower patients. These very concepts could be a significant help when applied to adolescents struggling with concentration and drug use issues. What care manager models represent are changes to the medical team, delivery of service, and the hope for improved outcomes which is exactly what could be beneficial for adolescents struggling with ADHD and *cannabis* use.

Conclusion

Smoking *cannabis* is dangerous for youth diagnosed with ADHD. When taking into account the documented neurological impact of smoking *cannabis*, combined with the vulnerability of early initiation, and the symptoms associated with ADHD, it is hard to argue that *cannabis* smoking is anything but high risk for youth diagnosed with ADHD. It is important to note the complexities revolving around such a topic as the dangers of *cannabis* smoking cannot be thought of as dichotomies or even in linear terms or the complexities would be lost. As mentioned above, there is research illustrating the health benefits of *cannabis* use; however, the opinion presented has research illustrating the dangers of smoking *cannabis* for youth diagnosed with ADHD. Though more research is needed to understand the effects of *cannabis* on the developing brain, adolescence is an especially dangerous time to smoke *cannabis* [39,40].

References

1. Hyshka E (2013) Applying a social determinants of health perspective to early adolescent cannabis use. An overview. *PsychINFO - ProQuest Drugs: Education, Prevention and Policy* 2013, pp: 110-119.
2. California Society of Addiction Medicine (2009) Impact of marijuana on children and adolescents.
3. Hadland SE, Harris SK (2014) Youth marijuana use: state of the science for the practicing clinician. *Curr Opin Pediatr* 26: 420-427.
4. Medina KL, Hanson KL, Schweinsburg AD, Cohen-Zion M, Nagel BJ, et al. Neuropsychological functioning in adolescent marijuana users: subtle deficits detectable after a month of abstinence. *J Int Neuropsychol Soc* 87: 114-119.
5. Meier MH, Caspi A, Ambler A, Harrington H, Houts R, et al. (2012) Persistent cannabis users show neuropsychological decline from childhood to midlife. *Proc Natl Acad Sci* 109: E2657-2664.
6. Crean RD, Crane NA, Mason BJ (2011) An evidence based review of acute and long term effects of cannabis use on executive cognitive functions. *J Addict Med* 5: 1-8.
7. Schacht JP, Hutchison KE, Filbey FM (2012) Associations between cannabinoid receptor-1 (CNR1) variation and hippocampus and amygdala volumes in heavy cannabis users. *Neuropsychopharmacology* 37: 2368-2376.
8. Zalesky A, Solowij N, Yucei M, Lubman DI, Takagi M, et al. (2012) Effect of long-term cannabis use on axonal fibre connectivity. *Brain* 135: 2245-2255.
9. Becker B, Wagner D, Gouzoulis-Mayfrank E, Spuentrup E, Daumann J (2010) Altered parahippocampal functioning in cannabis users is related to the frequency of use. *Psychopharmacology* 209: 361-374.
10. Churchwell JC, Lopez-Larson M, Yurgelun-Todd DA (2010) Altered frontal cortical volume and decision making in adolescent cannabis users. *Front Psychol* 1: 225.
11. Fernandez-Ruiz J, Berrendero F, Hernandez ML, Ramos JA (200) The endogenous cannabinoid system and brain development. *Trends Neurosci* 23: 14-20.
12. Kuepper R, van OJ, Lieb R, Wittchen HU, Höfler M, et al. (2011) Continued cannabis use and risk of incidence and persistence of psychotic symptoms: 10 year follow-up cohort study *BMJ* 1: 342.
13. Yücel M, Solowij N, Respondek C, Whittle S, Fornito A, et al. (2008) Regional brain abnormalities associated with long-term heavy cannabis use. *Arch Gen Psychiatry* 65: 694-701.
14. Sundram S (2006) Cannabis and neurodevelopment: implications for psychiatric disorders. *Hum Psychopharmacol* 2006;21: 245-254.
15. Schneider M (2008) Puberty as a highly vulnerable developmental period for the consequences of cannabis exposure. *Addict Biol* 13: 253-263.
16. Dinieri JA, Hurd YL (2012) Rat models of prenatal and adolescent cannabis exposure. *Methods Mol Biol* 829: 231-242.
17. Degenhardt L, Coffey C, Romaniuk H, Swift W, Carlin JB, et al. (2013) The persistence of the association between adolescent cannabis use and common mental disorders into young adulthood. *Addiction* 108: 124-133.
18. Jacobus J, Goldenberg D, Wierenga CE, Tolentino NJ, Liu TT, et al. (2012) Altered cerebral blood flow and neurocognitive correlates in adolescent cannabis users. *Psychopharmacology (Berl)* 222: 675-684.
19. Jacobus J, Squeglia LM, Meruelo AD, Castro N, Brumbach T, et al. (2015) Cortical thickness in adolescent marijuana and alcohol users: A three-year prospective study from adolescence to young adulthood. *Dev Cogn Neurosci* 16: 101-109.
20. Tamm L, Epstein JN, Lisdahl KM, Molina B, Tapert S, et al. (2013) Impact of ADHD and cannabis use on executive functioning in young adults. *Drug Alcohol Depend* 133: 607-614.
21. Elkins IJ, McGue M, Iacono WG (2007) Prospective effects of attention-deficit/hyperactivity disorder, conduct disorder, and sex on adolescent substance use and abuse. *Arch Gen Psychiatry* 64: 1145-1152.
22. Bidwell LC, Henry EA, Willcutt EG, Kinnear MK, Ito TA (2014) Childhood and current ADHD symptom dimensions are associated with more severe cannabis outcomes in college students. *Drug Alcohol Depend* 135: 88-94.
23. Ameringer KJ, Leventhal AM (2013) Associations between attention deficit hyperactivity disorder symptom domains and DSM-IV lifetime substance dependence. *Am J Addict* 22: 23-32.
24. Distel MA, Vink JM, Bartels M, Beijsterveldt CEM, Neale MC, et al. (2011) Age moderates non-genetic influences on the initiation of cannabis use: a twin-sibling study in Dutch adolescents and young adults. *Addiction* 106: 1658-1666.
25. Silva N, Szobot CM, Shih MC, Hoexter MQ, Anselmi CE, et al. (2014) Searching for a neurobiological basis for self-medication theory in ADHD comorbid with substance use disorders: an in vivo study of dopamine transporters using (99m)Tc-TRODAT-1 SPECT. *Clin Nucl Med* 39: e129-134.

26. Ross JM, Cox S, Schuster RM, Rojas A, Gonzalez R (2015) The moderating effects of cannabis use and decision making on the relationship between conduct disorder and risky sexual behavior. *J Clin Exp Neuropsychol* 37: 303–315.
27. Kalayasiri, R, Gelernter J, Farrer L, Weiss R, Brady K, et al. (2010) Adolescent cannabis use increases risk for cocaine-induced paranoia. *Drug and Alcohol Dependence* 107: 196–201.
28. Smeerdijk M, Linszen D, Kuipers T, Keet R (2002) Family motivational intervention in early psychosis and cannabis misuse. In: Lobban, Fiona; Barrowclough C, editor. *A casebook of family interventions for psychosis*, pp: 645–654.
29. Urban NB, Slifstein M, Thompson JL, Xu X, Grigis RR, et al. (2012) Dopamine release in chronic cannabis users: A [(11) C] raclopride positron emission tomography study. *Biol Psychiatry* 71: 677–683.
30. Lubman DI, Cheetham A, Yücel M (2014) Cannabis and adolescent brain development. *Pharmacol Ther* pp: S0163-7258.
31. Rubino T, Parolaro D (2014) Cannabis abuse in adolescence and the risk of psychosis: a brief review of the preclinical evidence. *Prog Neuropsychopharmacol Biol Psychiatry* 52: 41–44.
32. Becona E, Martinez U, Calafat A, Fernandez-Hermida JR, Juan M, et al. (2013) Parental permissiveness, control, and affect and drug use among adolescents. *Psicothema* 25: 292–298.
33. Bernburg JG, Thorlindsson T, Sigfusdottir ID (2009) The neighborhood effects of disrupted family processes on adolescent substance use. *Soc Sci Med* 69: 129–137.
34. Butters JE (2002) Family stressors and adolescent cannabis use: a pathway to problem use. *J Adolesc* 25: 645–654.
35. Kohn L, Kittel F, Piette D (2004) Peer, family integration and other determinants of cannabis use among teenagers. *Int J Adolesc Med Health* 16: 359–370.
36. Pinchevsky GM, Arria AM, Caldeira KM, Garnier-Dykstra LM, Vincent KB, et al. (2012) Marijuana exposure opportunity and initiation during college: parent and peer influences. *Prev Sci* 13: 43–54.
37. Korhonen T, Huizink A, Dick D, Pulkkinen L, Rose R, et al. (2008) Role of individual, peer and family factors in the use of cannabis and other illicit drugs: A longitudinal analysis among Finnish adolescent twins. *Drug Alcohol Depend* 97: 33–43.
38. Ciccone MM, Aquilino A, Cortese F, Scicchitano P, Sassara M, et al. (2010) Feasibility and effectiveness of a disease and care management model in the primary health care system for patients with heart failure and diabetes (Project Leonardo). *Vasc Health Risk Manag* 6: 297–305.
39. Greydanus DE, Hawver EK, Greydanus MM, Merrick J (2013) Marijuana: current concepts. *Front Public Health* 1: 42.
40. Greydanus DE, Kaplan G, Baxter LE (2015) Cannabis: the never-ending, nefarious nepenthe of the 21st century. *Dis Mon* 61: 118–75.