

Spontaneous Emergence of Information in Simple Biological Systems

Ricard J*

Jacques Monod Institute CNRS University, Paris, France

Short Communication

Information, no doubt plays an important role in the processes leading to the appearance of life on Earth. Hence, in this perspective, it is important to understand how a system, or a simple biological system, can generate such information. Let us consider, for instance, a simple process where a primitive catalyst, E, randomly binds two substrates, A and B, before releasing the products. The aim of the present paper is to raise the question to know whether such system can generate information.

Simple Enzyme System and its Information

Let us consider the simple process involving an enzyme E that randomly binds two substrates A and B before releasing the corresponding product(s). In such a system one can define the probabilities of occurrence of events A and B as

$$p(A) = \frac{K_1[A](1 + K_2[B]) + u_{EA}}{1 + K_1[A] + K_3[B] + K_1K_2[A][B] + u_{EA} + u_{EB} + u_E} \quad (1)$$

And

$$p(B) = \frac{K_3[B](1 + K_4[A]) + u_{EB}}{1 + K_1[A] + K_3[B] + K_1K_2[A][B] + u_{EA} + u_{EB} + u_E} \quad (2)$$

In these expressions, K_1, K_2, \dots are the apparent equilibrium constants of A and B to the various states of the protein. However, as the real system is not in equilibrium it has to be "corrected" by the so-called u functions defined as [1]

$$\begin{aligned} u_E &= \frac{k(k_{-1} + k_2[B])(k_{-3} + k_4[A])}{k_{-3}k_{-4}(k_{-1} + k_2[B]) + k_{-1}k_{-2}(k_{-3} + k_4[A])} \\ u_{EA} &= \frac{k_1[A]}{k_{-1} + k_2[B]} u_E \\ u_{EB} &= \frac{k_3[B]}{k_{-3} + k_4[A]} u_E \\ u_{EB}^* &= u_{EB} / k_3[B] \end{aligned} \quad (3)$$

Thus the biochemical network possesses nodes, i.e. the free enzyme E and the two enzyme-substrate complexes EA and EB. The u 's thus represent local reaction circuits that maintain an approximate steady state of the system. u_E is a reaction flow leading to the E node and both u_{EA} and u_{EB} are flows leading to nodes EA and EB. Hence the u 's are intrinsic flows that confer some dynamics to the system.

The information consumed, or generated, in such a system is defined as

$$I(A : B) = \log \frac{p(A|B)}{p(A)} \quad (4)$$

or as

$$I(A : B) = I(A) - I(A|B) \quad (5)$$

which implies that

$$I(A) = -\log p(A) \quad (6)$$

And

$$I(A|B) = -\log p(A|B) \quad (7)$$

Information is generated by the system if [2,3]

$$p(A|B) > p(A) \quad (8)$$

Looking back at expressions (1) and (2) allows to write

$$p(A|B) = \frac{[EAB]}{[EB] + [EAB]} = \frac{K_4[A]}{1 + K_4[A] + (u_{EB} / K_3[B])} \quad (9)$$

and it follows that

$$\frac{p(A|B)}{p(A)} = \frac{K_4[A]}{1 + K_4[A] + (u_{EB} / K_3[B])} \frac{1 + K_1[A] + K_3[B] + K_1K_2[A][B] + u_{EA} + u_{EB} + u_E}{K_1[A](1 + K_3[B]) + u_{EA}} \quad (10)$$

The numerator, N, and the denominator, D, of this expression can be rewritten as

$$N = K_4[A] + K_1K_4[A]^2 + K_3K_4[A][B] + K_1K_2K_4[A]^2[B] + K_4[A]u_E + K_4[A]u_{EA} + K_4[A]u_{EB} \quad (11)$$

$$D = K_1[A] + K_1K_2[A][B] + u_{EA} + K_1K_4[A]^2 + K_1K_2K_4[A]^2[B] + K_4[A]u_{EA} + K_1[A]u_{EB}^* + K_1K_2[A][B]u_{EB}^* + u_{EA}$$

and the difference $D-N$ can be expressed as

$$D-N = K_1[A] - K_4[A] + K_1K_2[A][B]u_{EB}^* - K_4[A]u_E + K_1[A]u_{EB}^* - K_4[A]u_{EB} + u_{EB}^*u_{EA} + u_{EA} \quad (12)$$

Moreover one has

$$K_1K_2[A][B]u_{EB}^* - K_4[A]u_E = \frac{K_1K_2[A][B] - K_4[A](k_{-3} + k_4[A])}{k_{-3} + k_4[A]} u_E \quad (13)$$

Similarly one has

$$K_1[A]u_{EB}^* - K_4[A]u_{EB} = \frac{(K_1[A] - K_4[A])k_3[B]}{k_{-3} + k_4[A]} u_E \quad (14)$$

Hence it appears that the differences above can be positive or negative whereas $u_{EB}^*u_{EA}$ and u_{EA} are of necessity positive, namely

$$K_1[A] - K_4[A]; K_1K_2[A][B]u_{EB}^* - K_4[A]u_E; K_1[A]u_{EB}^* - K_4[A]u_{EB}; u_{EB}^*u_{EA}; u_{EA} \quad (15)$$

It follows from these simple mathematical developments that the system considered can generate, or alternatively consume, information.

*Corresponding author: Jacques Ricard, Jacques Monod Institute CNRS University, Paris, France, Tel: 33 (0)1 57 27; E-mail: Jkricard@aol.com

Received January 08, 2015; Accepted January 10, 2015; Published January 13, 2015

Citation: Ricard J (2015) Spontaneous Emergence of Information in Simple Biological Systems. Curr Synthetic Sys Biol 3: 117. doi:10.4172/23320737.1000117

Copyright: © 2015 Ricard J. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Expressions (13) and (14) show that if $K_4 > K_1$ then the system generates information.

General Considerations and Conclusions

It can be noticed that, within a simple biochemical system such as an enzyme catalysed reaction, the interaction between several elements of this system can produce, or dissipate, information. We have seen previously that two situations have particularly interesting implications, namely:

Situation 1: $D(u)-N(u)>0$

Situation 2: $D(u)-N(u)<0$

The first situation (a) implies that

$$k_3[B] > k_3 + k_4[A] \quad (a)$$

$$K_2[B] < 1$$

And the second situation (b) means that

$$k_{-3} + k_4[A] > k_3[B] \quad (b)$$

$$K_2[B] > 1$$

Condition (a) means that EA and EB accumulate in the system and condition (b) means that E and EAB accumulate. The latter condition is indeed in favor of a chemical transformation of A and B into the corresponding products P and Q . In a way, one can consider that the system “choose” between these two possibilities. This choice is a consequence of the topology of the system. It also appears that the local reaction circuits u play a major role in the emergence, or the lack of emergence, of information in the system.

References

1. Ricard J (2010) Systems biology and the origins of life? Part II. Are biochemical networks possible ancestors of living systems? Networks of catalysed chemical reactions: Non-equilibrium, self-organization and evolution. Comptes Rendus Biologies 333: 769-778.
2. Gallager RG (1964) Information theory In: The Mathematics of Physics and Chemistry, Van Nostrand, Reinhold Company 2: 190-249.
3. Ricard J (2006) Emergent Collective Properties. Networks and Information in Biology. Elsevier, Amsterdam, Boston.

Citation: Ricard J (2015) Spontaneous Emergence of Information in Simple Biological Systems. Curr Synthetic Sys Biol 3: 117. doi:10.4172/23320737.1000117

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 400 Open Access Journals
- 30,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.editorialmanager.com/omicsgroup/>