

Strain Improvement of *Brevibacillus borostelensis* R1 for Optimization of α -Amylase Production by Mutagens

K Suribabu^{1*}, T Lalitha Govardhan¹ and KPJ Hemalatha²¹PG Department of Microbiology and Research Centre, Dr. Lankapalli Bullayya Post-graduate College, AP, India²Department of Microbiology, Andhra University, AP, India

Abstract

Physical and chemical mutagens are promising and are used for screening of high yielding strains. The exponential increase in the application of amylases in various fields has placed stress and demand in both qualitative improvement and quantitative enhancement through strain improvement. Ultraviolet light exerts its mutagenic effect by exciting electrons in molecules. The potent UV mutants which showed more than 20 mm zone of starch hydrolysis were screened and selected at 42% of survival time at 80minutes of exposure. The wild strain with fixed parameters yielded (3000 U/ml). The major findings of the strain improvement were out of ten mutants isolated, two (UV-3 and UV-10) showed 3000-4000 U/ml of amylase activity. The % of survival of *Brevibacillus borstelensis* R1 in Pikovskaya's medium was 25.75% at 120 minutes of exposure. Ten mutants (HNO₂-10, HNO₂-30, EMS-4, EtBr-40, EtBr-50, Acr-1, Acr-20, Acr-30, Acr-4 and 5'-FU-50) out of fifty mutants isolated showed 3000-4300U/ml of amylase activity, which was higher than the wild strain. The potent *Bacillus* species screened from marine water was *Brevibacillus borstelensis* R1. The α -amylase was found to be useful in bakery, food, fodder for poultry, automation dishwashing and laundry industries.

Keywords: *Brevibacillus borostelensis* R1; Ultra violet (UV) rays; Nitrous acid (HNO₂); Ethyl methane sulfonate (EMS); Ethidium bromide (EtBr); Acrylamide (Acr); 5'-fluoro uracil (5'-FU)

Introduction

Strain improvement

Physical and chemical mutagens are promising and are used for screening of high yielding strains [1]. Microorganisms usually produce commercially important metabolites in very low concentrations by their inherent control system. Although the yield may be increased by optimizing the cultural conditions, ultimately the productivity is controlled by the organism's genome [2]. In the last few decades, the exponential increase in the application of amylases in various fields has placed stress and demand in both qualitative improvement and quantitative enhancement through strain improvement. Such improved strains reduce the cost of the process. The challenge is to isolate those strains which are true mutants that carry beneficial mutations [3]. *Bacillus* spp. such as *B. subtilis* and *B. amyloliquefaciens* are the most commonly used organisms of choice for amylase production [4,5]. Highly active α -amylase is preferred for the conversion of starch into oligosaccharides. So, it is worthwhile to select a potent microbial strain for α -amylase production.

Physical mutagen

Ultraviolet (UV) light exerts its mutagenic effect by exciting electrons in molecules. The excitation of electrons in DNA molecules often results in the formation of extra bonds between adjacent pyrimidines in DNA. The UV light is the best studied mutagenic agent in prokaryotic organisms [6]. The UV mutant strains of *Bacillus* reported to show better ability to produce alpha amylase, which can be derived by mutagenesis and extensive screening [7-10].

Chemical mutagens are stronger mutagenic agents and cause permanent changes in DNA sequence. They bring about transitions from G: C \rightarrow A: T [11] and have preferential effect on DNA replication [12-16].

Nitrous acid (HNO₂, Hydroxidooxidonitrogen, O=NO) causes interstrand cross-linking of DNA. Nitrous acid acts as a mutagen by deamination of the NH₂ group of adenine and cytosine to an ether group, thus altering their base pairing. It causes oxidative deamination of particular bases. Conversion of the amino groups to keto groups changes the hydrogen bonding potential of the bases. Nitrous acid was reported to be a suitable mutagen for the improvement of α -amylase production in *Bacillus* spp. [17-20].

Ethyl Methane Sulfonate (EMS) is an alkylating agent which carries one, two or more alkyl groups in reactive form. The transfer of methyl or ethyl groups to the bases such that their base-pairing potentials are altered and transitions result. Thus it induces all types of mutations (transitions, transversions, frame shifts and even chromosome aberrations) with various relative frequencies. Ethyl methane sulfonate was reported as potent mutagen in amylase production in *Bacillus* spp. [20-22].

Acrylamide (acrylic amide) decomposes non-thermally to form ammonia and thermal decomposition produces carbon monoxide, carbon dioxide and oxides of nitrogen. It is effective in causing bacterial mutation. It shows direct and indirect modes of interaction with DNA results in alkylating DNA adducts [23].

Base analog mutagen (5'-fluoro uracil) is a thymine analog. It

***Corresponding author:** Dr. K Suribabu, PG Department of Microbiology and Research Centre, Dr. Lankapalli Bullayya Post-graduate College, Visakhapatnam-530 013, A.P, India, Tel: 9440788795; E-mail: ksuribabu_sda@yahoo.com

Received January 21, 2014; **Accepted** March 06, 2014; **Published** March 11, 2014

Citation: Suribabu K, Lalitha Govardhan T, Hemalatha KPJ (2014) Strain Improvement of *Brevibacillus borostelensis* R1 for Optimization of α -Amylase Production by Mutagens. J Microb Biochem Technol 6: 123-127. doi:[10.4172/1948-5948.1000132](https://doi.org/10.4172/1948-5948.1000132)

Copyright: © 2014 Suribabu K, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

has a higher likelihood for tautomerization to the enol form (FU*). Methyl group in 5th position of thymine and fluorine at 5'-position in 5'-FU have similar effects. After tautomeric shift to its enol form, 5'-fluorouracil pairs with guanine and thus causes A. T to G. C transitions. Base analogs cause transitions in the bidirectional mode.

Ethidium bromide (EtBr, Intercalating agent) acts as a mutagen because it intercalates double stranded DNA and deforms the DNA [24]. Ethidium bromide is a large, flat basic molecule that resembles a DNA base pair. This insertion causes "stretching" of the DNA duplex and the DNA polymerase is "fooled" into inserting an extra base opposite an intercalated molecule. This results in frame shift mutations. This affects DNA biological processes, like DNA replication and transcription.

Materials and Methods

Primary screening of α -amylase producing bacteria

The collected marine water samples were diluted by serial dilution technique. The diluted samples of 10^{-4} to 10^{-6} (0.1 ml) were spread with L-shaped glass rod by spread plate technique on the starch agar plates. After incubation at 37°C for 24 hours, the plates were flooded with Lugol solution (1% iodine in 2% potassium iodide w/v) [25].

Estimation of amylase by DNS method

Maltose produced by the hydrolytic activity of α -amylase on α -1, 4 linkages present in polysaccharides, reduce 3, 5 dinitro salicylate to an orange red colored 5-nitro 3-amino salicylate which can be measured at 520 nm. The starch substrate [0.5 ml of 0.5% in 0.1 M phosphate buffer (pH 6.8)] was mixed with 1% (0.2 ml) NaCl in a test tube and pre incubated at 37°C for 10 minutes. The supernatant collected from the centrifugation of the production media was used as enzyme source, 0.5 ml of this was added to the reaction mixture. The reaction was terminated by the addition of 1.0 ml of 3, 5-dinitrosalicylic acid reagent [1.0 gm DNS in 0.8% NaOH, 60% Na K tartrate] after incubation at 37°C for 15 minutes. The contents were mixed well and kept in boiling water bath for 10 minutes. Then they were cooled and diluted with 10 ml of distilled H₂O. The color developed was read at 520 nm. One unit of enzyme activity was defined as the amount of enzyme that releases 1.0 mmol of reducing sugar (maltose) per minute under the assay conditions [26].

UV survival curve of *Brevibacillus borstelensis* R1

Inoculate the *Brevibacillus borstelensis* R1 in nutrient broth and incubate in rotary shaking incubator at 120 rpm at 37°C for 24 hrs. Hundred ml of the culture was taken in a petridish and exposed to UV irradiation by placing at a distance of 30 cm from the UV radiation source (emitting the energy of 1.6×10^2 J/m²/s). After every 10 minutes of exposure 5.0 ml of the sample was drawn. One ml of the sample was serially diluted, 10^{-4} - 10^{-7} dilutions were used for the viable count method. The control was the culture sample without UV treatment. A graph was plotted with time in minutes on the X-axis and log cfu per ml on Y-axis in order to obtain survival curve.

Ultraviolet treatment to isolate mutants

Twenty four hours old bacterial culture prepared in nutrient broth was aseptically centrifuged at 5000 rpm for 15 minutes. The bacterial sediment was made suspension with sterile saline water. Ten ml of diluted suspension was transferred to sterilized petriplates placed under the UV lamp (emitting the energy of 1.6×10^2 J/m²/s) for 80

minutes (42% survival time) to 120 minutes. The bacterial suspension (0.5 ml) was collected after every 10 minutes interval and transferred to solidified starch agar petriplates (180×20 mm BOROSIL) by spread plate technique. The plates were invertedly incubated in an incubator at 37°C for 24 hrs.

Chemical treatment

Five ml of chemical mutagens [Nitrous acid (National), Ethyl methane sulfonate (HIMEIDA), Ethidium bromide (MERCK), Acrylamide (HIMEDIA) and 5'-fluoro uracil (ONCO FLUOR)] of 50-250 μ g/ml were transferred to sterilized centrifuge tube separately containing 5.0 ml of bacterial suspension. The tubes were placed at room temperature for 120 minutes. Then the tubes were centrifuged at 5000 rpm for 15 minutes. The supernatant was discarded to remove the chemical mutagen from the bacterial cells. Ten ml of sterile saline water was added to each centrifuge tube. The tubes were re-centrifuged thrice for the removal of traces of chemical mutagen from bacterial cell sediment. After washing the cells, 10 ml of sterilized saline water was added to each tube to form bacterial suspension. The suspension (0.5 ml) was transferred to the petriplates (180×20 mm) containing solidified starch agar by spread plate method and incubated at 37°C for 24 hrs.

From both physical and chemical mutagen treatment the mutant colonies showing more than 20 mm zone of starch hydrolysis in starch agar medium were isolated in pure culture (streak plate technique) on nutrient agar plates. The mutants were single line streaked over starch agar plates to confirm the size of the zone of starch hydrolysis. The confirmed mutants were subjected to submerged fermentation (SmF) at 120 rpm in rotary shaking incubator in Pikovskaya's fermentation medium (100 ml) in Erlenmeyer flasks with predetermined physical and chemical parameters. The α -amylase activity was assayed by DNS method.

Results and Discussion

Isolation of mutant strains by using physical and chemical mutagens

The optimum production of α -amylase by *Brevibacillus borstelensis* R1 was observed in Pikovskaya's medium at temperature-37°C, pH-7.0, salinity-1%, with 5% jaggary, 2% beef extract and 0.8% calcium chloride [27]. The activity of α -amylase produced by wild strain was found to be 3000 U/ml with zone of starch hydrolysis (Figure 1). The above optimized conditions were used for all physical and chemical mutagen experiments for strain improvement studies in our laboratory experiments.

Physical mutagen

The potent physical mutants which showed more than 20 mm zone of starch hydrolysis were screened and selected at different minutes of exposure of UV irradiation. Control is the culture unexposed to UV radiations. The potent UV mutants which showed more than 20 mm zone of starch hydrolysis were screened and selected at 42% of survival time at 80 minutes of exposure (Figure 2). The wild strain with fixed parameters yielded (3000 U/ml). Out of ten mutants isolated, two UV-3 and UV-10 showed 1.32 and 1.29 fold increase in of amylase activity, respectively (Table 1). The % of survival of *Brevibacillus borstelensis* R1 in Pikovskaya's medium was 25.75% after 120 minutes of exposure.

Jin et al. [28] have developed a hyper producing α -amylase mutant




Figure 1: Zone of starch hydrolysis *Brevibacillus borstelensis* R1.




Figure 2: Ultraviolet survival curve of *Brevibacillus borstelensis* R1.

S.no.	Time of exposure in minutes	Isolate code	Zone of starch hydrolysis (mm)	Alpha-amylase activity (U/ml) in Pikovskaya's medium
1	80	UV-1	21 ± 1	903 ± 6
2		UV-2	24 ± 4	2633 ± 58
3	90	UV-3	43 ± 6	3967 ± 58
4		UV-4	21 ± 4	1830 ± 52
5	100	UV-5	27 ± 4	2433 ± 58
6		UV-6	22 ± 2	1417 ± 29
7	110	UV-7	20 ± 2	1633 ± 58
8		UV-8	27 ± 4	2533 ± 58
9	120	UV-9	18 ± 4	823 ± 4
10		UV-10	33 ± 5	3867 ± 58

Values presented in the table are means of triplicates ± SD

Table 1: Ultra violet treated mutant isolates showing α -amylase potency in starch media and Pikovskaya's Medium.

of *Bacillus licheniformis* which yielded 50 times higher enzyme activity than the parental strain. The α -amylase production was decreased in 48 mutant strains out of 60 mutant strains isolated when compared to parent strain/wild strain. Similarly decrease in production of α -amylase was reported by Haq et al. [29] in mutant strain of *Bacillus amyloliquefaciens* UNG-16 when compared to the wild parental strain. The effect of UV irradiation on *Bacillus* spp. was most effective in the production of α -amylase [30-33].

Chemical mutagens

The wild strain with fixed physical parameters and chemical parameters yielded 2100 U/ml and 3000 U/ml enzyme activity

S.no.	Conc. of chemical mutagen (μ g/ml)	Isolate code	Zone of starch hydrolysis (mm)	Alpha-amylase activity (U/ml) in Pikovskaya's medium
1	50	HNO ₂ -1	21 ± 2	910 ± 17
2		HNO ₂ -10	35 ± 4	3967 ± 58
3	100	HNO ₂ -2	29 ± 3	2433 ± 58
4		HNO ₂ -20	25 ± 4	2833 ± 58
5	150	HNO ₂ -3	22 ± 4	1213 ± 23
6		HNO ₂ -30	33 ± 6	3833 ± 116
7	200	HNO ₂ -4	33 ± 5	2617 ± 29
8		HNO ₂ -40	24 ± 4	2033 ± 58
9	250	HNO ₂ -5	21 ± 3	2233 ± 58
10		HNO ₂ -50	25 ± 4	2017 ± 29
11	50	EMS-1	21 ± 2	2027 ± 46
12		EMS-10	33 ± 5	2367 ± 58
13	100	EMS-2	25 ± 4	2230 ± 52
14		EMS-20	21 ± 2	1567 ± 58
15	150	EMS-3	23 ± 5	1310 ± 17
16		EMS-30	28 ± 4	2825 ± 43
17	200	EMS-4	36 ± 2	4167 ± 58
18		EMS-40	24 ± 3	1313 ± 23
19	250	EMS-5	21 ± 5	1827 ± 46
20		EMS-50	27 ± 3	2167 ± 58
21	50	EtBr -1	26 ± 3	2233 ± 58
22		EtBr -10	21 ± 4	1023 ± 25
23	100	EtBr -2	28 ± 2	2367 ± 58
24		EtBr -20	27 ± 4	2433 ± 58
25	150	EtBr -3	29 ± 2	2767 ± 58
26		EtBr -30	21 ± 5	813 ± 23
27	200	EtBr -4	23 ± 2	1133 ± 58
28		EtBr -40	37 ± 3	4166 ± 58
29	250	EtBr -5	21 ± 5	1833 ± 58
30		EtBr -50	33 ± 5	4017 ± 76
31	50	Acr-1	33 ± 5	3950 ± 87
32		Acr-10	26 ± 2	2433 ± 58
33	100	Acr-2	25 ± 5	2833 ± 58
34		Acr-20	33 ± 5	4042 ± 72
35	150	Acr-3	22 ± 5	1333 ± 58
36		Acr-30	35 ± 5	3883 ± 29
37	200	Acr-4	33 ± 6	4237 ± 64
38		Acr-40	29 ± 2	2833 ± 58
39	250	Acr-5	39 ± 6	2833 ± 58
40		Acr-50	22 ± 4	1140 ± 53
41	50	5'-FU-1	30 ± 4	2783 ± 29
42		5'-FU-10	20 ± 5	1830 ± 52
43	100	5'-FU-2	21 ± 5	813 ± 23
44		5'-FU-20	27 ± 4	2577 ± 40
45	150	5'-FU-3	29 ± 2	2827 ± 46
46		5'-FU-30	30 ± 5	2810 ± 17
47	200	5'-FU-4	21 ± 5	717 ± 29
48		5'-FU-40	32 ± 5	2817 ± 58
49	250	5'-FU-5	20 ± 1	1227 ± 46
50		5'-FU-50	33 ± 5	3650 ± 50

Values presented in the table are means of triplicates ± SD

Table 2: Chemical mutant strains [Nitrous acid (HNO₂), Ethyl methane sulfonate (EMS), Ethidium bromide (EtBr), Acrylamide (Acr) and 5'-fluoro uracil (5'-FU)] showing α -amylase potency in starch media and Pikovskaya's medium.

respectively. Out of fifty mutants isolated ten mutants HNO₂-10, HNO₂-30, EMS-4, EtBr-40, EtBr-50, Acr-1, Acr-20, Acr-30, Acr-4 and 5'-FU-50 showed 1.32, 1.27, 1.38, 1.38, 1.34, 1.32, 1.35, 1.29, 1.41 and 1.22 fold increase in amylase activity, respectively which was higher than the wild strain (Table 2).

Haq et al. [12] reported that nitrous acid was found to be the suitable mutagen for improvement of α -amylase production in *Bacillus* spp. Similarly, maximum production of α -amylase (285 U/ml) was reported in *Aspergillus oryzae* NA17 and also in *Aspergillus oryzae* by Szafranec et al. [19] and Haq et al. [30].

Haq et al. [29] screened mutant strains of *Bacillus amyloliquefaciens* EMS-6 which gave 102.78 \pm 2.22 U/ml/min α -amylase activities. The activity was around 1.4 fold higher than the parental strain *Bacillus amyloliquefaciens* UNG-16. Similarly, Szafranec et al. [19] and Haq et al. [30] worked with *Aspergillus oryzae* (EMS-18) isolate which gave maximum production of α -amylase 347 U/ml.

Haq et al. [12] reported the mutagenesis for improvement of α -amylase production of bacterial strains with EMS. Similarly Sarikaya and Gürgün [21] worked on EMS mutants of *Bacillus subtilis*. Ikram-ul-haq et al. [22] reported on mutants of *Bacillus licheniformis* GCB-30UCM strain. Sarikaya and Gürgün [21] reported about the mutants of *Bacillus subtilis* obtained by ethidium bromide which increased amylase production.

Conclusions

Strain improvement of *B. borestelensis* R1 was carried out by using physical and chemical mutagens to develop a better yielding strain to improve the production of α -amylase. Various mutagenic agents such as ultraviolet rays (UV), nitrous acid (HNO₂), ethyl methyl sulfonate (EMS), ethidium bromide (EtBr), acrylamide (Acr), 5'-fluoro uracil (5'-FU) were used for strain improvement studies. Out of ten mutant strains isolated, two strains (UV-3 and UV-10) showed 3000-4000 U/ml of α -amylase activity. Out of fifty chemical mutant strains isolated, ten mutant strains (HNO₂-10, HNO₂-30, EMS-4, EtBr-40, EtBr-50, Acr-1, Acr-20, Acr-30, Acr-4 and 5'-FU-50) showed 3000-4300 U/ml of α -amylase activity which was higher than the wild parent strain. Chemical mutagens are more effective in strain improvement than physical mutagen.

Acknowledgements

We thank Management of Dr. Lankapalli Bullayya College, Visakhapatnam for the financial support and facilities provided to make this work possible.

References

1. Sidney PC, Nathan OK (1975) Compositions and Uses for an Alpha-Amylase Polypeptide of Bacillus Spp. Meth. InEnz. 3: 26.
2. Stanbury A, Whitaker, Hall SJ (1995) Fermentation Technology' Principles of Fermentation Technology', Pergamon Press, Oxford, 1-24.
3. Parekh S, Vinci VA, Strobe RJ (2000) Improvement of microbial strains and fermentation processes. App Microbiol Biotechnol 54: 287-301.
4. Haq I, Shamim N, Ashraf H, Ali S, Qadeer MA(2005) Effect of surfactants on the biosynthesis of alpha-amylase by Bacillus subtilis GCBM-25. Pak J Bot 37: 373-379.
5. Sivaramakrishnan S, Gangadharan D, Nampoothiri KM, Socol CR, Pandey A (2006) α -Amylases from Microbial sources-An Overview on Recent Developments. Food Techn Biotechnol 44: 173-184.
6. Meenu M, Santhosh D, Kamia C, Randhir S (2000) Strain improvement of Aspergillusflavus for enhanced production Ind J Microbiol 40: 25-32.
7. Dinu DL, Jurcoane S, Vamanu E, Burcea M, Campeanu G (2001) Studies for enhancement of alkaline amylase production to the strain Bacillus sp. ICCF 276 Rom. Biotechnology Lett 6:2
8. Rivera MH, López-Munguía A, Soberón X, Saab-Rincón G (2003) Alpha-amylase from Bacillus licheniformis mutants near to the catalytic site: effects on hydrolytic and transglycosylation activity. Protein Eng 16: 505-514.
9. Shiao RJ, Hung HC, Jeang CL (2003) Improving the thermostability of raw-starch-digesting amylase from a Cytophaga sp. by site-directed mutagenesis. Appl Environ Microbiol 69: 2383-2385.
10. Surendra S, Vinni S, Manohar LS (2011) Biotechnological applications of industrially important amylase enzyme. Int J Pharm Bio Sci 2: 45-51.
11. Miller JH (1972) In: Experiments in molecular genetics, 11th edition (Cold Spring Harbor Laboratory, Cold Spring Harbor, New York), 113.
12. Haq I, Ashraf H, Rani S, Qadeer MA (2002) Biosynthesis of alpha amylase by chemically treated mutant of Bacillus subtilis GCBU-20. Pak J Biol Sci 2: 73-75.
13. Joh JH, Kim BGT, Kong WS, Yoo YB, Chu KS, et al. (2004) Isolation and characterization of dikaryoticmutans from Pleurotusostreatus by UV irradiation. Mycobiology 32: 88-96.
14. Paul ND, Jacobson RJ, Taylor A, Wargent JJ, Moore JP (2005) The use of wavelength-selective plastic cladding materials in horticulture: understanding of crop and fungal responses through the assessment of biological spectral weighting functions. Photochem Photobiol 81: 1052-1060.
15. Wu CH, Apweiler R, Bairoch A, Natale DA, Barker WC, et al. (2006) The Universal Protein Resource (UniProt): an expanding universe of protein information. Nucleic Acids Res 34: D187-191.
16. Besoain XA, Perez LM, Araya A, Lefever L, Sanguinetti M, et al. (2007) New strains obtained after UV treatment and protoplast fusion of native Trichodermaharzianum: their biocontrol activity onPyrenochaetalycopersici. Elec J Biotechnol 10: 16-28.
17. Azin M, Noroozi E (2001) Random mutagenesis and use of 2-deoxy-D-glucose as an antimetabolite for selection of α -amylase overproducing mutants of Aspergillusoryzae. W J Microbiol Biotechnol 17: 747-750.
18. Rubinder K, Chadha BS, Singh N, Saini HS, Singh S (2002) Amylase hyperproduction by deregulated mutants of the thermophilic fungus Thermomyceslanuginosus. J Ind Microbiol Biotechnol 29: 70-74.
19. Szafranec K, Wloch DM, Sliwa P, Borts RH, Korona R (2003) Small fitness effects and weak genetic interactions between deleterious mutations in heterozygous loci of the yeast Saccharomyces cerevisiae. Genet Res 82: 19-31.
20. Karanam SK, Medicherla NR (2008) Enhanced lipase production by mutation induced Aspergillusjaponicus. Afri J Biotechnol 7: 2064-2067.
21. Sarikaya E, Gürgün V (2000) Increase of the α -amylase yield by some Bacillus strains. Turk J Biol 24: 299-308.
22. Ikram-ul-haq S, Ali A, Saleem, Javed MM (2009) Mutagenesis Of Bacillus licheniformis through Ethyl Methanesulfonate for alpha Amylase Production. Pak J Bot 41: 1489-1498.
23. Besaratinia A, Pfeifer GP (2005) DNA adduction and mutagenic properties of acrylamide. Mutat Res 580: 31-40.
24. McCann J, Ames B N (1975) Detection of carcinogens as mutagens in the Salmonella/microsome test: assay of 300 chemicals. Proc Natl Acad Sci U S A 72: 5135-5139.
25. Amoozegar MA, Malekzadeh F, Malik KA (2003) Production of amylase by newly isolated moderate halophile, Halobacillus sp. strain MA-2. J Microbiol Methods 52: 353-359.
26. Miller GL (1959) Use of Dinitro salicylic acid reagent for determination of reducing sugar. Anal Chem 31: 426 - 429.
27. Suribabu K, Lalitha GT, Hemalatha KPJ (2014) Optimization of physical parameters of α -amylase Producing Brevibacillusborostelensis R1 in submerged fermentation. Int Jr Rech Eng Tech 3: 517-525.
28. Jin B, VanLeeuwen HJ, Patel B, Yu Q (1998) Utilization of starch Process in waste water for production of microbial biomass protein and fungal α -amylase by Aspergillusoryzae. Bioresour Technol 66: 201-206.
29. Haq I, Ali S, Javed MM, Hameed Um Saleem A, et al. (2010) Production of alpha amylase from a randomly induced mutant strain of Bacillus amyloliquefaciens and its application as a desizer in textile industry. Pak J Bot 42: 473-484.

-
30. Haq I, Ashraf H, Abdullah R, Shah AH (2002b) Isolation and screening of Fungi for the biosynthesis of alpha- amylase. Biotechnol 2:66.
31. Pandey A, Nigam P, Soccol CR, Soccol VT, Singh D, et al. (2000) Advances in microbial amylases. Biotechnol Appl Biochem 31: 135-152.
32. Varalakshmi KN, Kumudini BS, Nandini BN, Solomon JD, Mahesh B, et al. (2008) Characterization of Alpha Amylase from Bacillus sp.1/isolated from paddy seeds. J Appl Bio Sci 1: 46-53.
33. Roheena Abdullah (2005) Studies on the production of alpha amylase by *Aspergillusoryzae* using submerged fermentation. Institute of Industrial Biotechnology, GC University Lahore. Thesis 20-91.

Citation: Suribabu K, Lalitha Govardhan T, Hemalatha KPJ (2014) Strain Improvement of *Brevibacillus borostelensis* R1 for Optimization of α -Amylase Production by Mutagens. J Microb Biochem Technol 6: 123-127. doi:[10.4172/1948-5948.1000132](https://doi.org/10.4172/1948-5948.1000132)

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 300 Open Access Journals
- 25,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.editorialmanager.com/jmbt>

