

Sulfamic Acid: An Efficient and Recyclable Solid Acid Catalyst for the Synthesis of Quinoline-4-Carboxylic Acid Derivatives in Water

Yahya S, Beheshtiha SH, Majid M* and Dehghani M

Department of Chemistry, School of Science, Alzahra University, Vanak, Tehran, Iran

Abstract

A simple, cost-effective, environmentally, and convenient procedure for the synthesis of quinoline-4-carboxylic acid derivatives is described through a one-pot MCR condensation of pyruvic acid, various aniline derivatives, and differently substituted aryl aldehydes in the presence of sulfamic acid ($\text{NH}_2\text{SO}_3\text{H}$) (SA) as an efficient and recyclable catalytic system under mild reaction conditions in water.

Keywords: Quinoline-4-carboxylic acid derivatives; One-pot MCR; Sulfamic acid; Recyclable catalyst; Green chemistry

Introduction

Quinoline [1] 1-aza-naphthalene and benzo[b]pyridine are nitrogen containing heterocyclic aromatic compounds. Quinolines are very important compounds due to their wide spectrum of biological activities behaving as anti-malarial, anti-bacterial, anti-fungal, anti-asthmatic, anti-hypertensive, anti-inflammatory, anti-platelet activity [2-9]. In addition quinolines have been employed in the study of bioorganic and bio organo-metallic processes [10,11]. Considering the significant applications in the fields of medicinal, bioorganic, industrial, and synthetic organic chemistry, there has been tremendous interest in developing efficient methods for the synthesis of quinolines. Some derivatives of such as quinoline-4-carboxylic acid elicited profound changes in the morphology of typical tips of *Botrytis cinerea* [12] Quinoline-4-carboxylic acids are one of the most important series of quinoline derivatives because they exhibit a wide variety of medicinal effects and are applied as active components in industrial antioxidants [13,14]. Meanwhile, quinoline-4-carboxylic acids are the key precursors for the synthesis of other useful quinoline derivatives [15]. Despite remarkable efforts in the last decade [16-19] the development of effective methods for the synthesis of quinoxaline ring is still an important challenge and much in demands. In recent years, many methods for the synthesis of these quinoline acids have successively been reported. The conventional synthesis involves the Pfitzinger reaction [20,21]. Doebner reaction [22,23], Friedlander and Combes methods [24-31]. However, these synthetic methods require expensive and hazardous solvents and chemicals, as well as harsh reaction condition. Therefore, the development of simple, convenient, and environmentally benign approaches for the synthesis of quinolines is still desirable. In recent years, the use of solid acids as heterogeneous catalysts has received tremendous interest in various areas of organic synthesis [32-34]. Heterogeneous solid acids are advantageous over conventional homogeneous acid catalysts as they can be easily recovered from the reaction mixture by simple filtration and can be reused after activation or without activation and more importantly without appreciable loss in activities. SA has emerged as a substitute for conventional Bronsted- and Lewis acid catalysts. This catalyst is an amino acid containing sulfur element with mild acidity. It is fast coming up as a stable, white odorless crystalline, commercially available, also not volatile and corrosive, thus can be considered as green catalyst in organic synthesis [35-37]. Interestingly, sulfamic acid exists not only in its amino sulfonic acid form, but also as $\text{H}_3\text{N}^+\text{SO}_3^-$ zwitterionic units [38] immiscible with commonly employed non-polar organic solvents [35]. Also, it is a white crystalline solid [39]. In recent years, SA has been extensively used as an efficient heterogeneous catalyst for acid catalyzed reactions, such as, acetalization, [40] esterification, [41-43] functional group protections and deprotections [44] the Michael addition [45]

amino Diels-Alder reactions [46], Biginelli condensations [47] and Beckmann rearrangement [48].

As the aforesaid conditions are not compatible with heat- or acid-sensitive substrates, there is a need to develop an effective synthesis of quinoxalines employing more ecofriendly conditions and catalysts [49-60]. With considering green chemistry principles, especially using water greenest and most abundant solvent [61-70]. We have recently reviewed the application of Doebner reaction in synthesis of heterocycles including quinolone-4-carboxylic acids [71]. We also frequently used Sulfamic acid as an efficient catalyst in different organic transformations [72-80] and published a review highlighting the application of Sulfamic acid in organic synthesis [81,82]. Armed with these experiences, in response to the need for the facile, efficient and green synthesis of quinolone-4-carboxylic acid derivatives, herein we wish to report the high yielding, one pot MCRs synthesis of this heterocyclic *via* condensation of pyruvic acid, various amines and aryl benzaldehyde in the presence of SA as a relatively inexpensive and available catalyst in water under mild reaction conditions. These eco-friendly protocols offer several advantages such as green, convenient and cost-effective procedures with high yield, shorter reaction time, simpler work-up, recovery, and reusability of solid acid heterogeneous catalyst (SA) in subsequent reactions.

Results and Discussion


To optimize the reaction conditions, a mixture of pyruvic acid (1.2 mmol), aniline (1.1 mmol), and benzaldehyde (1.0 mmol) was selected as a model reaction (Scheme 1) and was examined under different conditions. Initially, we examined various catalysts such as $\text{NH}_2\text{SO}_3\text{H}$, $\text{H}_6\text{P}_2\text{Mo}_{18}\text{O}_{62}$, HNO_3 , DABCO, Nano- Fe_3O_4 and also tested the un-catalyzed reaction (Table 1). Because of type of reaction that needs acidic or basic catalyst, or in another word needs a stimulus to begin a reaction and also base on previous work [82] we choose these catalysts to examine in this reaction (Scheme 1). As shown in Table 1, among the catalysts tested, $\text{NH}_2\text{SO}_3\text{H}$ (SA) gave the highest yield for the corresponding quinoline derivative (Table 1) [83-85].

*Corresponding author: Majid M, Department of Chemistry, School of Science, Alzahra University, Vanak, Tehran, Iran, Tel: +989121329147; E-mail: mmh1331@yahoo.com

Received October 19, 2016; Accepted November 26, 2016; Published November 30, 2016

Citation: Yahya S, Beheshtiha SH, Majid M, Dehghani M (2016) Sulfamic Acid: An Efficient and Recyclable Solid Acid Catalyst for the Synthesis of Quinoline-4-Carboxylic Acid Derivatives in Water. Mod Chem appl 4: 195. doi: 10.4172/2329-6798.1000195

Copyright: © 2016 Yahya S, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



Entry	Catalyst (mol %)	Time (hr)	Yield ^b (%)
1	NH ₂ SO ₃ H	3	90
2	H ₆ P ₂ Mo ₁₈ O ₆₂	3	85
3	HNO ₃	3	81
4	DABCO	4.5	59
5	Nano-Fe ₃ O ₄	6	68
6	-	18	Trace

Table 1: Synthesis of 4a in presence of different catalysts in reflux condition. Synthesis of 4a in presence of different catalysts in reflux condition. a 1.2 mmol (1), 1.1 mmol (2a) and 1.0 mmol (3a) in presence of 3 mol% of catalyst and 5 mL of H₂O. ^bRefers to the isolated yield.

The other catalysts examined such as HNO₃ and H₆P₂Mo₁₈O₆₂ were also effective but generally not as much (Table 1). To find a suitable solvent, the model reaction was carried out in different solvents (Table 2). The best result was achieved by conducting the reaction in the presence of 3 mol% of SA under reflux conditions in water (Table 2) [82,86]. It was also found out that the best result is achieved in the presence of 3 mol% of catalyst (SA) under reflux condition (Table 3). To establish, the substrate scope of this methodology, we reacted divergent benzaldehydes, and amines with pyruvic acid under the optimized reaction conditions to obtain the corresponding quinoline-4-carboxylic acids in excellent yields (Scheme 2; 4a-i). All compounds were known and their structures were confirmed by comparison of physical and spectral data with those of already reported (Table 4). The results were listed in Table 4 [82]. A proposed mechanism for the SA-catalyzed formation of quinoline-4-carboxylic acid derivatives is proposed and shown in Scheme 3. The mechanism of this reaction using other catalysts was suggested and explained in recent years [90-93]. The proposed Sulfamic acid-catalyzed reaction involves the 1,2- addition of the aniline to aromatic aldehyde to form adducts 5, which followed by elimination of H₂O to achieve intermediate 6. The latter is subjected to Mannich reaction with pyruvic acid which followed by intermolecular cyclization, to afford the intermediate 8, which in turn dehydrated and then dehydrated and oxidized to give the corresponding quinoline 4 [82] (Scheme 3). From the green chemistry points of view, experiments concerning the recycling and reuse of the catalyst (SA) was carried out. The insolubility of the catalyst SA in water made the catalyzed reaction, heterogeneous, thus the catalyst is separated by a simple filtration. Upon the completion of the reaction, monitored and observed with TLC, the catalyst SA was separated by filtration and reused. As shown in Figure 1, the catalyst could efficiently have recovered even after 4 cycles without suffering any significant drop in its catalytic activity or the yield of reaction (Figure 1) [82,83]. In summary, an extremely facile, high-yielding method for the preparation of quinoline-4-carboxylic acid derivatives under mild reaction conditions, *via* a one-pot MCR, in the presence of an efficient, highly effective, economically, and reusable solid acid heterogeneous catalyst in was developed. This catalyst could easily be separated from the reaction mixture with high yields and purity and can be directly reused after simple extraction. In addition, using water as green solvent and operational simplicity are the merits of this method. The procedure has many advantages such as short reaction time, the requirement of small amount of catalyst, mild acidity conditions, giving virtually quantitative yields and being performed under environmentally friendly conditions.

Experimental Section

Materials and equipment's

Chemicals and solvents were purchased from Merck-Aldrich and used directly with high-grade quality, without any purification. TLC

Entry	Solvent	Time (hr)	Temperature (°C)	Yield ^b (%)
1	H ₂ O	3	Reflux	90
2	CH ₃ CH ₂ OH	6.5	Reflux	80
3	CH ₃ CN	18	Reflux	Trace
4	-	26	100	69


Table 2: Synthesis of 4a in presence of different solvents. Synthesis of 4a in presence of different solvents. a 1.2 mmol (1), 1.1 mmol (2a) and 1.0 mmol (3a) in presence of 3 mol% of catalyst (Sulfamic acid) and 5 mL of solvent. ^bRefers to the isolated yield.


Entry (3212)	Catalyst mol %	Time (hr)	Temperature (°C)	Yield ^b (%)
1	2	4.5	Reflux	78
2	3	3	Reflux	90
3	4	3	Reflux	90
4	5	3	Reflux	90
5	2	5	100	62
6	3	5.5	100	50
7	2	8.5	120	43
8	4	9	130	75
9	3	4	70-80	71
10	3	5.5	90-100	77
11 ^c	3	8	110-120	80

Table 3: Synthesis of 4a in presence of different amount of catalyst and temperature. Synthesis of 4a in presence of different amount of catalyst (Sulfamic acid) and temperature. a 1.2 mmol (1), 1.1 mmol (2a) and 1.0 mmol (3a). ^bRefers to the isolated yield. ^cIn solvent free condition.


Entry	R ¹	R ²	Product	Time (hr)	Yield ^b (%)	Mp (°C)/Lit. Mp [Ref]
1	H	H	4a	3	90	211-212/212-214 [82]
2	4-Me	H	4b	4	88	213-214/218-219 [83]
3	4-NO ₂	H	4c	11	43	233-235/238-239 [84]
4	3-Cl	H	4d	4	80	256-259/256-260 [85]
5	H	4-OMe	4e	7	83	213-215/216-217 [86]
6	H	4-OH	4f	3	89	268-270/271-273 [87]
7	H	4-NO ₂	4g	3	90	319-322/324 [88]
8	4-Me	4-OMe	4h	3.5	79	231-233/235-237 [89]
9	4-Me	4-NO ₂	4i	5.5	91	263/265-266 [82]

Table 4: Synthesis of quinoline-4-carboxylic acid derivatives (4a-i) under optimized conditions Synthesis of quinoline-4-carboxylic acid derivatives (4a-i) under optimized conditions. a 1.2 mmol (1), 1.1 mmol (2) and 1.0 mmol (3) in presence of 3 mol% of catalyst (Sulfamic acid) and 5 mL H₂O at reflux condition. ^bRefers to the isolated yield.





Scheme 2: Synthesis of quinoline-4-carboxylic acid derivatives.



Scheme 3: Proposed mechanism of the synthesis of 4.

analyses were done using percolated TLC silica gel 60 F₂₅₄ (Merck) plates. The ¹H NMR spectra were recorded by Bruker Ultrashield 500 MHz respectively advance instrument, with CDCl₃ used as solvent. Proton resonances are designated as singlet (s), doublet (d), triplet (t) and multiplet (m). FT-IR spectra were recorded using KBr disks on FTIR Bruker Tensor 27 instrument in the 500-4000 cm⁻¹ region. The vibration transition frequencies are reported in wave numbers (cm⁻¹). Band intensities are assigned as weak (*w*), medium (*m*), and strong (*s*). Melting points were measured using a capillary tube method with a Barnstead Electrothermal 9200 apparatus. All products were known and identified by comparison of their physical and spectroscopic data with those of authentic compounds and found being identical. All yields refer to isolated products.

The synthesis of quinoline-4-carboxylic acid derivatives (4a-i)

General procedure: A mixture of pyruvic acid (1.2 mmol), amine (1.1 mmol), and benzaldehyde (1.0 mmol) in 5 ml of water was well stirred with NH₂SO₃H (3 mol%). This mixture was magnetically stirred under reflux for appropriate time according to Table 4. The progress of reaction was monitored by TLC (7:3, *n*-hexane/ethyl acetate). Upon completion of the reaction, the mixture was cooled to room temperature. After filtration, the catalyst which reminded in the aqueous phase can be recovered simply by removing the water through heating and the aqueous phase, and stored for use in subsequent run under the same conditions without any treatment.

The remaining solution is concentrated by increasing slightly CH₂Cl₂, and was magnetically stirred for 5 min. The precipitate was filtered to give quinoline-4-carboxylic acid as a crude. This crude was recrystallized with acetic acid to get pure product (Table 4; 4a-i) All the triazoles are known compounds, and their physical data were found to be identical with those of authentic samples [82-89].

Compound 4a: m.p.: 211-212°C [82]. IR (KBr): ν =3429, 3335, 3042, 1679, 1590, 1487, 1378 cm⁻¹; ¹H NMR (500 MHz, CDCl₃): δ =8.76 (d, 1H, J=8.5 Hz), 8.59 (s, 1H), 8.22 (d, 1H, J=8.5 Hz), 8.36 (d, 2H, J=7.4 Hz), 7.30 (t, 1H, J=7.5 Hz), 7.53 (t, 1H, J=7.5 Hz), 7.95 (m, 2H), 6.50 (m, 1H).

Compound 4b: m.p.: 213-214°C [87]. ¹H NMR (500 MHz, CDCl₃): δ =7.50 (s, 1H), 8.55 (s, 1H), 8.20 (m, 2H), 8.10 (d, 1H, J=8.4 Hz), 7.36 (d, 1H, J=8.4 Hz), 7.70 (t, 2H, J=7.4 Hz), 7.22 (m, 1H), 2.62 (s, 3H).

Compound 4c: m.p.: 233-235°C [88]. ¹H NMR (500 MHz, CDCl₃): δ =8.86 (d, 1H, J=8.9 Hz), 7.45 (m, 2H), 7.00-7.40 (m, 5H), 8.15 (m, 1H).

Compound 4d: m.p.: 256-259°C [89]. ¹H NMR (500 MHz, CDCl₃): δ =8.52 (s, 1H), 8.20-8.39 (m, 2H), 7.49 (m, 5H), 7.89 (d, 1H, J=9.1 Hz).

Compound 4e: m.p.: 213-215°C [90]. ¹H NMR (500 MHz, CDCl₃): δ =8.20 (s, 1H), 7.23 (d, 1H, J=8.9 Hz), 7.10 (d, 2H, J=8.0 Hz), 7.42 (t, 1H, J=7.0 Hz), 7.30 (d, 2H, J=8.9 Hz), 8.47 (d, 1H, J=8.0 Hz), 6.76 (t, 1H, J=7.1 Hz), 3.72 (s, 3H).

Compound 4f: m.p.: 268-270°C [91]. ¹H NMR (500 MHz, CDCl₃): δ =8.23 (d, 1H, J=8.7 Hz), 7.51 (s, 1H), 8.22 (m, 3H), 7.50 (t, 1H, J=7.3 Hz), 8.38 (d, 2H, J=8.7 Hz), 6.62 (t, 1H, J=7.3 Hz).

Compound 4g: m.p.: 319-322°C [92]. ¹H NMR (500 MHz, CDCl₃): δ =8.25 (d, 1H, J=8.0 Hz), 7.35 (s, 1H), 8.83 (m, 2H), 7.13 (d, 1H, J=8.1 Hz), 7.78 (t, 1H, J=7.6 Hz), 7.56 (t, 1H, J=7.5 Hz), 7.00 (m, 2H).

Compound 4h: m.p.: 231-233°C [93]. ¹H NMR (500 MHz, CDCl₃): δ =8.22 (s, 1H), 8.41 (s, 1H), 8.29 (d, 2H, J=8.3 Hz), 7.16 (d, 1H, J=8.2 Hz), 7.85 (d, 1H, J=8.3 Hz), 7.25 (d, 2H, J=8.4 Hz), 2.90 (s, 3H), 2.62 (s, 3H).

Compound 4i: m.p.: 263°C [82]. ¹H NMR (500 MHz, CDCl₃): δ =8.95 (s, 1H), 7.52 (s, 1H), 8.47 (m, 4H), 8.49 (d, 1H, J=8.7 Hz), 7.23 (d, 1H, J=9.4 Hz), 2.85 (s, 3H).

Acknowledgements

The authors are thankful to Alzahra University Research Council for partial financial assistance.

References

1. Abadi AH, Hegazy GH, Zaher AAE (2005) Synthesis of novel 4-substituted-7-trifluoromethylquinoline derivatives with nitric oxide releasing properties and their evaluation as analgesic and anti-inflammatory agents. *Bio org Med Chem* 13: 5759-5765.
2. Larsen RD, Corley EG, King AO, Carrol JD, Davis P, et al. (1996) Practical Route to a New Class of LTD4 Receptor Antagonists. *Org Chem* 61: 3398-3405.
3. Chen YL, Fang KC, Sheu JY, Hsu SL, Tzeng CC, et al. (2001) Synthesis and antibacterial evaluation of certain quinolone derivatives. *J Med Chem* 44: 2374-2377
4. Roma G, Braccio MD, Grossi G, Mattioli F, Ghia M, et al. (2000) Synthesis and antioxidant properties of novel quinoline-chalcogenium compounds. *Eur J Med Chem* 35: 1021
5. Kalluraya B, Sreenivasa S (1998) Synthesis and pharmacological properties of some quinoline derivatives. *Farmaco* 53: 399-404.
6. Doube D, Blouin M, Brideau C, Chan C, Desmarais S (1998) *Bioorg Med Chem Lett* 8: 1255.
7. Ko TC, Hour MJ, Lien JC, Teng CM, Lee KH (2001) Synthesis of 4-alkoxy-2-phenylquinoline derivatives as potent antiplatelet agents. *Bioorg Med Chem Lett* 11: 279-282.
8. Ferrarini PL, Mori C, Badweh M, Manera C, Martinelli A, et al. (1997) *Heterocycl Chem* 34: 1501.
9. Maguire MP, Sheets KR, McVety K, Spada AP, Zilberstein A, et al. (1994) A new series of PDGF receptor tyrosine kinase inhibitors: 3-substituted quinoline derivatives. *J Med Chem* 37: 2129-2137.
10. Saito I, Sando S, Nakatani K (2001) Improved selectivity for the binding of naphthyridine dimer to guanine-guanine mismatch. *Bioorg Med Chem* 9: 2381-2385.
11. He C, Lippard SJ, *Am J* (2001) *Chem Soc* 40: 1414.
12. Strigacova J, Hudcová D, Varecká L, Lasiková A (2000) Some biological properties of new quinoline-4-carboxylic acid and quinoline-4-carboxamide derivatives. *Folia Microbiol (Praha)* 45: 305-309.
13. Michael JP (2005) Quinoline, quinazoline and acridone alkaloids. *Nat Prod Rep* 22: 627-646.
14. Michael JP (2004) Quinoline, quinazoline and acridone alkaloids. *Nat Prod Rep* 21: 650-668.
15. Yu XY, Hill JM, Yu G, Yang Y (2001) A series of quinoline analogues as potent inhibitors of *C. albicans* prolyl tRNA synthetase. *Biol Med Chem Lett* 11: 541-544.
16. Gobec S, Urleb U (2004) *Sci. Synth.* 16: 845.
17. Kim SY, Park KH, Chung YK (2005) Manganese(IV) dioxide-catalyzed synthesis of quinoxalines under microwave irradiation. *Chem Commun* 10: 1321-1323.
18. Raw SA, Wilfred CD, Taylor RJK (2003) Preparation of quinoxalines, dihydropyrazines, pyrazines and piperazines using tandem oxidation processes. *Chem Commun* 18: 2286-2287.
19. Antoniotti S, Dunach E (2002) Direct and catalytic synthesis of quinoxaline derivatives from epoxides and ene-1, 2-diamines. *Tetrahedron Lett* 43: 3971-3973.
20. Hoi NPB, Xuong ND, Lavit D (1953) Fluorine-containing analogs of 4-hydroxypropiophenone. *J Org Chem* 18: 910-915.
21. Atwell GJ, Baguley BC, Denny WA (1989) Potential antitumor agents. 57. 2-Phenylquinoline-8-carboxamides as "minimal" DNA-intercalating antitumor agents with in vivo solid tumor activity. *J Med Chem* 32: 396-401.
22. Dobner VO, Gieseke M, Liebig J (1887) *Ann Chem* 242: 290.
23. Allen CFH, Spangler FW, Webster ER (1951) 2-aminopyridine and the doebner reaction. *J Org Chem* 16: 17-20.

24. Jones G, Katritzky AR, Rees CW, Scriven EFV (1996) Pergamon, New York, USA 5: 167.
25. Cho CS, Oh BH, Kim TJ, Shim SC (2000) Synthesis of quinolines via ruthenium-catalyzed amine exchange reaction between anilines and trialkyl amines. *Chem Commun*, pp: 1885-1886.
26. Jiang B, Si YC (2002) Zn(II)-Mediated Alkynylation-Cyclization of *o*-Trifluoroacetyl Anilines: One-Pot Synthesis of 4-Trifluoromethyl-Substituted Quinoline Derivatives. *J Org Chem* 67: 9449-9451.
27. Skraup H (1880) *Chem Ber* 13: 2086.
28. Friedlander P (1882) *Chem Ber* 15: 2572.
29. Mansake RHF, Kulka M (1953) Skraup Synthesis of Quinolines. *Org React* 7: 59.
30. Linderman RJ, Kirillos SK (1990) *Tetrahedron Lett* 31: 2689-2798.
31. Theoclitou ME, Robinson LA (2002) Novel facile synthesis of 2, 2, 4 substituted 1, 2-dihydroquinolines via a modified Skraup reaction. *Tetrahedron Lett* 43: 3907-3910.
32. Clark JH (2002) Solid acids for green chemistry. *Acc Chem Res* 35: 791-797.
33. Helwani Z, Othman MR, Aziz N, Kim J, Fernando WJN (2009) Solid heterogeneous catalysts for transesterification of triglycerides with methanol: a review. *Appl Catal A: General* 363: 1-10.
34. Climent MJ, Corma A, Iborra S (2010) Heterogeneous catalysts for the one-pot synthesis of chemicals and fine chemicals. *Chem Rev* 111: 1072-1133.
35. Wang B (2005) Sulfamic acid: a very useful catalyst. *Synlett* 2005: 1342-1343.
36. Bo W, Ming YL, Shuan SJ (2003) Ionic liquid-regulated sulfamic acid: chemoselective catalyst for the transesterification of β -ketoesters. *Tetrahedron Lett* 44: 5037-5039.
37. Wang B, Gu Y, Luo C, Yang T, Yang L, et al. (2004) Sulfamic acid as a cost-effective and recyclable catalyst for liquid Beckmann rearrangement, a green process to produce amides from ketoximes without waste. *Tetrahedron Lett* 45: 3369-3372.
38. Harbison GS, Kye YS, Penner GH, Grandin M, Monette M (2002) ^{14}N Quadrupolar, ^{14}N and ^{15}N Chemical Shift, and ^{14}N - ^1H Dipolar Tensors of Sulfamic Acid. *J Phys Chem B* 106: 10285-10291.
39. Nonose N, Kubota M (1998) Determination of metal impurities in sulfamic acid by isotope dilution electrothermal vaporization inductively coupled plasma mass spectrometry. *J Anal At Spectrom* 13: 151-156.
40. Jin TS, Sun G, Li YW, Li TS (2002) An efficient and convenient procedure for the preparation of 1, 1-diacetates from aldehydes catalyzed by $\text{H}_2\text{NSO}_3\text{H}$. *Green Chem* 4: 255-256.
41. Yiming L, Huaxe ML (1988) 39: 407.
42. Jin L, Ru-Qi Z (2000) *Hecheng Huaxue* 8: 364.
43. Wang B, He J, Sun RC (2010) Carbamate synthesis from amines and dialkyl carbonate over inexpensive and clean acidic catalyst-Sulfamic acid. *Chinese Chem Lett* 21: 794-797.
44. An LT, Zou JP, Zhang LL, Zhang Y (2007) Sulfamic acid-catalyzed Michael addition of indoles and pyrrole to electron-deficient nitroolefins under solvent-free condition. *Tetrahedron Lett* 48: 4297-4300.
45. Nagarajan R, Magesh CJ, Perumal PT (2004) Inter- and intramolecular imino diels-alder reactions catalyzed by sulfamic acid: a mild and efficient catalyst for a one-pot synthesis of tetrahydroquinolines. *Synthesis* 2004: 69-74.
46. Li JT, Han JF, Yang JH, Li TS (2003) An efficient synthesis of 3, 4-dihydropyrimidin-2-ones catalyzed by $\text{NH}_2\text{SO}_3\text{H}$ under ultrasound irradiation. *Ultrason Sonochem* 10: 119-122.
47. Heravi MM, Alishiri T (2014) Dimethyl acetylenedicarboxylate. *Adv Heterocycl Chem* 113: 1.
48. Heravi MM, Talaei B (2014) Ketenes as privileged synthons in the synthesis of heterocyclic compounds, part 1: three- and four-membered heterocycles. *Adv Heterocycl Chem* 113: 143-244.
49. Heravi MM, Khaghaninejad S, Mostofi M (2015) Pechmann reaction in the synthesis of coumarin derivatives. *Adv Heterocycl Chem* 112: 1.
50. Heravi MM, Khaghaninejad S, Nazari N (2015) Bischler-Napieralski reaction in the synthesis of isoquinolines. *Adv Heterocycl Chem* 112: 183-226.
51. Heravi MM, Talaei B (2015) Chapter Three-Ketenes as Privileged Synthons in the Syntheses of Heterocyclic Compounds Part 2: Five-Membered Heterocycles. *Adv Heterocycl Chem* 114: 147-225.
52. Heravi MM, Vavsari VF (2015) Chapter Two-Recent Advances in Application of Amino Acids: Key Building Blocks in Design and Syntheses of Heterocyclic Compounds. *Adv Heterocycl Chem* 114: 77-145.
53. Khaghaninejad S, Heravi MM (2014) Paal-Knorr Reaction in the Synthesis of Heterocyclic Compounds. *Chem Inform*, p: 45.
54. Heravi MM, Zadsirjan V (2015) Chapter Five. Recent Advances in the Synthesis of Benzo [b] furans. *Adv Heterocycl Chem* 117: 261-376.
55. Heravi MM, Talaei B (2016) Chapter Five. Ketenes as Privileged Synthons in the Synthesis of Heterocyclic Compounds. Part 3: Six-Membered Heterocycles. *Adv Heterocycl Chem* 118: 195-291.
56. Heravi MM, Asadi S, Lashkariani BM (2013) Recent progress in asymmetric Biginelli reaction. *Molecular diversity* 17: 389-407.
57. Heravi MM, Hashemi E, Beheshtiha YS, Kamjou K, Toolabi M, et al. (2014) Solvent-free multicomponent reactions using the novel N-sulfonic acid modified poly (styrene-maleic anhydride) as a solid acid catalyst. *J Mol Catal A Chem* 392: 173-180.
58. Heravi MM, Mousavizadeh F, Ghobadi N, Tajbakhsh M (2014) A green and convenient protocol for the synthesis of novel pyrazolopyranopyrimidines via a one-pot, four-component reaction in water. *Tetrahedron Lett* 55: 1226-1228.
59. Nemati F, Heravi MM, Elhampour A (2015) Magnetic nano- $\text{Fe}_3\text{O}_4@ \text{TiO}_2/\text{Cu}_2\text{O}$ core-shell composite: an efficient novel catalyst for the regioselective synthesis of 1, 2, 3-triazoles using a click reaction. *RSC Adv* 5: 45775-45784.
60. Heravi MM, Sadjadi S, Oskooie HA, Shoar RH, Bamoharram FF (2008) Heteropolyacids as heterogeneous and recyclable catalysts for the synthesis of benzimidazoles. *Catal Commun* 9: 504-507.
61. Heravi MM, Zadsirjan V, Bakhtiari K, Oskooie HA, Bamoharram FF (2007) Green and reusable heteropolyacid catalyzed oxidation of benzylic, allylic and aliphatic alcohols to carbonyl compounds. *Catal Commun* 8: 315-318.
62. Heravi MM, Sadjadi S, Oskooie HA, Shoar RH, Bamoharram FF (2008) The synthesis of coumarin-3-carboxylic acids and 3-acetyl-coumarin derivatives using heteropolyacids as heterogeneous and recyclable catalysts. *Catal Commun* 9: 470-474.
63. Heravi MM, Hashemi E, Beheshtiha YS, Ahmadi S, Hosseinejad T (2014) PdCl₂ on modified poly (styrene-co-maleic anhydride): A highly active and recyclable catalyst for the Suzuki-Miyaura and Sonogashira reactions. *J Mol Catal A: Chem* 394: 74-82.
64. Heravi MM, Tajbakhsh M, Ahmadi AN, Mohajerani B (2006) Zeolites. Efficient and eco-friendly catalysts for the synthesis of benzimidazoles. *Monatsh Chem* 137: 175-179.
65. Heravi MM, Vazin Fard M, Faghihi Z (2013) Heteropoly acids-catalyzed organic reactions in water: doubly green reactions. *Green Chem Lett Rev* 6: 282-300.
66. Mirsafaei R, Heravi MM, Ahmadi S, Moslemi MH, Hosseinejad T (2015) *In situ* prepared copper nanoparticles on modified KIT-5 as an efficient recyclable catalyst and its applications in click reactions in water. *J Mol Catal A: Chem* 402: 100-108.
67. Heravi MM, Khorasani M, Derikvand F, Oskooie HA, Bamoharram FF (2007) Highly efficient synthesis of coumarin derivatives in the presence of H₁₄ [NaP₅W₃₀O₁₁₀] as a green and reusable catalyst. *Catal Commun* 8: 1886-1890.
68. Heravi MM, Bakhtiari K, Zadsirjan V, Bamoharram FF, Heravi OM (2007) Aqua mediated synthesis of substituted 2-amino-4H-chromenes catalyzed by green and reusable Preysslter heteropolyacid. *Bioorg Med Chem Lett* 17: 4262-4265.
69. Heravi MM, Asadi SH, Azarakhshi F (2014) *Curr Org Chem* 11: 701.
70. Heravi MM, Alinejad H, Derikvand F, Oskooie HA, Baghernejad B, et al. (2012) $\text{NH}_2\text{SO}_3\text{H}$ and $\text{H}_6\text{P}_2\text{W}_{18}\text{O}_{62} \cdot 18\text{H}_2\text{O}$ -Catalyzed, Three-Component, One-Pot Synthesis of Benzo [c] acridine Derivatives. *Syn Commun* 42: 2033-2039.
71. Heravi MM, Saeedi M, Beheshtiha YS, Oskooie HA (2011) One-pot chemoselective synthesis of novel fused pyrimidine derivatives. *Chem Heterocycl Compd (NY)* 47: 737-744.
72. Saeedi M, Beheshtiha YS, Heravi MM (2011) Synthesis of Novel Tetrahydro [4,5] imidazo [2,1-b]-chromeno [4,3,2-de] quinazoline and Benzothiazol-2-ylaminoxanthrone Derivatives. *Heterocycles* 83: 1831-1841.

73. Heravi MM, Saeedi M, Beheshtiha YS, Oskooie HA (2011) One-pot synthesis of benzochromeno-pyrazole derivatives. *Molecular diversity* 15: 239-243.
74. Heravi MM, Alinejhad H, Bakhtiari K, Oskooie HA (2010) Sulfamic acid catalyzed solvent-free synthesis of 10-aryl-7, 7-dimethyl-6, 7, 8, 10-tetrahydro-9H-[1, 3]-dioxolo [4, 5-b] xanthen-9-ones and 12-aryl-9, 9-dimethyl-8, 9, 10, 12-tetrahydro-11H-benzo [a] xanthen-11-ones. *Molecular diversity* 14: 621-626.
75. Heravi MM, Bakhtiari K, Alinejhad H, Saeedi M, Malakooti R (2010) MCM-41 Catalyzed Efficient Regioselective Synthesis of β -Aminoalcohol under Solvent-free Conditions. *Chin J Chem* 28: 269-272.
76. Heravi MM, Derikvand F, Ranjbar L (2010) Sulfamic acid-catalyzed, three-component, one-pot synthesis of [1,2,4] triazolo/benzimidazolo quinazolinone derivatives. *Synth Commun* 40: 677-685.
77. Heravi MM, Ranjbar L, Derikvand F, Alimadadi B (2008) Three-component one-pot synthesis of 4, 6-diarylpyrimidin-2 (1H)-ones under solvent-free conditions in the presence of sulfamic acid as a green and reusable catalyst. *Molecular diversity* 12: 191-196.
78. Heravi MM, Ranjbar L, Derikvand F, Bamoharram FF (2007) Sulfamic acid as a cost-effective catalyst instead of metal-containing acids for the one-pot synthesis of β -acetamido ketones. *J Mol Catal A Chem* 276: 226-229.
79. Heravi MM, Baghernejad B, Oskooie HA (2009) Application of sulfamic acid in organic synthesis-A short review. *Curr Org Chem* 13: 1002-1014.
80. Wang LM, Hu L, Chen HJ, Sui YY, Shen W (2009) One-pot synthesis of quinoline-4-carboxylic acid derivatives in water: Ytterbium perfluorooctanoate catalyzed Doebner reaction. *J Fluor Chem* 130: 406-409.
81. Wang L, Han J, Sheng J, Tian H, Fan Z (2005) Rare earth perfluorooctanoate [RE (PFO)₃] catalyzed one-pot Mannich reaction: three component synthesis of β -amino carbonyl compounds. *Catal Commun* 6: 201-204.
82. Iijima H, Kato T, Söderman O (2000) Variation in degree of counterion binding to cesium perfluorooctanoate micelles with surfactant concentration studied by ¹³³Cs and ¹⁹F NMR. *Langmuir* 16: 318-323.
83. Furó I, Sitnikov R (1999) Order parameter profile of perfluorinated chains in a micelle. *Langmuir* 15: 2669-2673.
84. Dvinskikh SV, Furo I (2000) Order parameter profile of perfluorinated chains in a lamellar phase. *Langmuir* 16: 2962-2967.
85. Boa AN, Canavan SP, Hirst PR, Ramsey C, Stead AM, et al. (2005) Synthesis of brequinar analogue inhibitors of malaria parasite dihydroorotate dehydrogenase. *Bioorg Med Chem* 13: 1945-1967.
86. McCloskey CM (1952) 6-Nitrocinchophen and Related Substances. *J Am Chem Soc* 74: 5922-5924.
87. Rapport MM, Seneor AE, Mead JF, Koepfli JB (1946) The Synthesis of Potential Antimalarials. 2-Phenyl- α -(2-piperidyl)-4-quinolinemethanols. *J Am Chem Soc* 68: 2697-2703.
88. Lindwall HG, Bandes J, Weinberg I (1931) Preparation of certain brominated cinchophens. *J Am Chem Soc* 53: 317-319.
89. Belen'kaya RS, Boreko EI, Zemtsova MN, Kalinina MI, Timofeeva MM, et al. (1981) Synthesis and antiviral activity of 2-[aryl (hetaryl)] quinoline-4-carboxylic acids. *Pharm Chem J* 15: 171-176.
90. Schneider W, Pothmann A (1941) Über die Lösungsfarben von Phenolbetainen der Chinolinreihe. *Berichte der deutschen chemischen Gesellschaft (A and B Series)* 74: 471-493.
91. Boykin Jr DW, Patel AR, Lutz RE (1968) Antimalarials. IV. New synthesis of α -(2-pyridyl)-and α -(2-piperidyl)-2-aryl-4-quinolinemethanols. *J Med Chem* 11: 273-277.
92. Wu YC, Liu L, Li HJ, Wang D, Chen YJ (2006) Skraup-Doebner-von Miller quinoline synthesis revisited: Reversal of the regiochemistry for γ -aryl- β , γ -unsaturated α -ketoesters. *J Org Chem* 71: 6592-6595.
93. Denmark SE, Venkatraman S (2006) On the mechanism of the Skraup-Doebner-Von Miller quinoline synthesis. *J Org Chem* 71: 1668-1676.

Citation: Yahya S, Beheshtiha SH, Majid M, Dehghani M (2016) Sulfamic Acid: An Efficient and Recyclable Solid Acid Catalyst for the Synthesis of Quinoline-4-Carboxylic Acid Derivatives in Water. *Mod Chem appl* 4: 195. doi: [10.4172/2329-6798.1000195](https://doi.org/10.4172/2329-6798.1000195)

OMICS International: Open Access Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission>