

# SWI/SNF Chromatin Remodeling Complex in Regulating Mesenchymal Stem Cell Lineage Specification

De-Meng Chen<sup>1</sup>, Xin-Qi Zhong<sup>2</sup>, Kai Wang<sup>3</sup> and Yi-Zhou Jiang<sup>1\*</sup>

<sup>1</sup>Institute for Advanced Study, Shenzhen University, Shenzhen,

<sup>2</sup>The Third Affiliated Hospital of Guangzhou Medical University, Guangzhou, and

<sup>3</sup>Clinical and Translational Research Center, Shanghai First Maternity and Infant Hospital, Tongji University School of Medicine, Shanghai, China

## Abstract

Mesenchymal Stem Cells (MSC) can be obtained from various tissues and differentiate into many different lineages, including osteoblasts, adipocytes, chondrocytes, cardiomyocytes, hepatocytes and neural cells both *in vivo* and *in vitro*. However, the ability of MSC to differentiate into specific lineages seems to be restricted and requires a deeper understanding of the genetic and epigenetic mechanisms. Epigenetic mechanism refers to a process that regulates heritable alterations in gene expression without changing the DNA sequence. SWI/SNF (SWI/SNF/Sucrose Non-Fermentable), a chromatin-remodeling complex serves as an ideal intervention point for lineage manipulation of MSC. In this review, we discuss the importance of SWI/SNF chromatin remodeling complex in regulating the fate determination of MSC. We propose that selectively manipulation of subunits of SWI/SNF will enhance the lineage-specific differentiation of MSC and improve therapeutic application of MSC.

**Keywords:** SWI/SNF; MSC; Lineage commitment; Tissue engineering

## Introduction

Mesenchymal Stem Cells (MSC), also called multipotential stromal cells or mesenchymal stromal cells, are a heterogeneous subset of stem cells that can be isolated from various adult tissues, including bone marrow, adipose tissue and other sources [1-3]. Initial studies have showed that MSC are able to differentiate into the mesoderm cell lineages, including bone, cartilage, adipose tissue, skeletal muscle and cardiac tissue [1,4,5]. Recently, studies have also found that MSC can differentiate into endoderm and ectoderm lineages, such as hepatocytes, neurons, epidermal-like cells [6-8]. MSC are an outstanding choice for regenerative medicine, tissue engineering and clinical therapy. Differentiation of MSC is mediated by intrinsic and extrinsic regulatory mechanisms, including specific growth factors, signaling molecules and epigenetic modifications [9]. These regulatory factors together define a selective transcription of discrete combination of genes, which define a differentiation program and determine the specific lineage and phenotype. Better understanding of how specific lineage commitment is achieved and especially how epigenetic mechanisms take part in this process will facilitate the development of more effective protocols for MSC tissue engineering and cellular therapy. In this review, we summarize our current knowledge of how SWI/SNF chromatin-remodeling complex modulate multi-lineage differentiation of MSC.

## SWI/SNF chromatin-remodeling complexes

Gene transcription is dynamically regulated at different levels across a wide range of cellular processes, including stemness, differentiation and proliferation. The expression level of certain gene largely depends on the state of its associated chromatin. Histone modification enzymes or chromatin-associated multisubunit protein complexes can affect whether a gene is transcriptionally activated or repressed [10,11]. Based on the identity of their catalytic subunit, chromatin-remodeling complexes can be divided into four different classes, including SWI/SNF, ISWI, CHD and INO80 [12-14]. Switch/Sucrose Non Fermentable (SWI/SNF) enzymes are a family of evolutionarily conserved, multi-subunit complexes that depend on energy from Adenosine Triphosphate (ATP) to regulate the chromatin structure [15]. SWI/SNF complex consist either BRG1 or BRM, two alternative ATPases that

catalyze the chromatin complex remodeling, along with 8-12 BRG1/BRM-associated factors (BAFs) [15]. BRG1 and BRM share 75% amino acid sequence identity and have similar domains: the QLQ domain (protein-protein interaction), the HSA domain (DNA binding), the ATPase domain and the bromodomain (acetyl-lysine histone mark recognition) [16].

The complexes share several common members, such as BAF170, BAF155, BAF60a/b/c, BAF57, BAF53a/b, BAF47, BAF45a/b/c/d and  $\beta$ -actin [17]. BRG1-containing complexes can be further classified into Polybromo-Associated Factor (PBAF) complexes, which uniquely contain BAF200 and BAF180 or BAF complexes, which uniquely contain BAF250A or BAF250B [16]. Among all BAFs members, BAF170, BAF155 and BAF47 are the functional core subunits for chromatin remodeling *in vitro* [15]. While other BAFs members are essential for interactions with transcriptional activators and repressors *in vivo*. The composition of SWI/SNF can be reconfigured during cell fate transitions through cell type-specific and developmental stage-specific expression patterns of certain subunits, which will be discussed in details later.

To regulate transcription activity, SWI/SNF chromatin-remodeling complexes are recruited by transcription factors to the promoters of target genes, where they can disrupt histone-DNA contacts and allow transcription factors to access their sequence-specific DNA [16,18]. Generally, SWI/SNF-mediated chromatin remodeling leads to sliding of nucleosome, eviction of nearby nucleosome or selective removal of H2A/H2B dimers near the Transcription Start Site (TSS) [18]. These

**\*Corresponding author:** Yi-Zhou Jiang, Institute for Advanced Study, Shenzhen University, Nanhai Road 3688, Shenzhen, Guangdong, China 518060, Tel: (+86) 0755-26530611; E-mail: [jiangyz@szu.edu.cn](mailto:jiangyz@szu.edu.cn)

**Received** August 22, 2015; **Accepted** September 10, 2015; **Published** September 17, 2015

**Citation:** Chen DM, Zhong XQ, Wang K, Jiang YZ (2015) SWI/SNF Chromatin Remodeling Complex in Regulating Mesenchymal Stem Cell Lineage Specification. J Tissue Sci Eng 6: 154. doi:10.4172/2157-7552.1000154

**Copyright:** © 2015 Chen DM, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

activities modulate RNA Polymerase II occupancy and transcriptional initiation. Depending on whether a transcriptional activator or repressor recruits SWI/SNF, transcription can be up regulated or down regulated. The energy of ATP hydrolysis is harnessed to disrupt histone-DNA contacts and move nucleosomes away from the TSS or toward the TSS [16,18].

SWI/SNF is essential for self-renewal of stem cells and is critical for cell differentiation. Depletion of core subunits of SWI/SNF complexes, such as BRG1, BAF155 and BAF47, affects the survival of totipotent cells and leads to peri-implantation lethality in mouse [19-21]. Deficiency of BRG1, BAF155 or BAF250a/b impaired the ability of mESCs to differentiate into ectoderm, mesoderm and endoderm layers [22- 26]. BRG1 has been shown to co-occupy a large number of the genomic loci with OCT4 and SOX2 by ChIP-seq analysis, indicating the importance of SWI/SNF in regulating stem cell property [23]. The altered expression of BRG1 promoted the senescence of MSC with suppression of the NANOG transcription, which is part of the transcriptional circuitry governing stem cell functions. Overall, regulation of gene expression patterns during differentiation of any cell lineage requires cooperation with the SWI/SNF chromatin-remodeling complex [28].

### SWI/SNF and osteogenesis

Bone formation of MSC can be achieved by treatment of Bone Morphogenetic Proteins (BMPs) and Wnt proteins, which target a number of osteoblast-determining transcription factors, such as Runx2, Osterix (Osx), Id1, Dlx3 [29, 30]. Both Runx2 and Osx knockout mice fail to develop mature osteoblasts. Hence these two factors are widely accepted as master transcription factors for osteogenesis. Osx acts downstream of Runx2 during bone formation and regulates the expression of osteoblast markers, including collagen type I, Osteopontin (OP) and Osteocalcin (OC) [30]. Several studies have shown that SWI/SNF activity plays a critical role for induction of the osteoblast phenotype. For example, Brg1 is expressed in developing skeletal tissues of the mouse embryo and in *ex vivo* osteoblast cultures [31]. This expression of Brg1 depends on BMP2-induced Runx2 expression. Deficiency of Brg1 in calvarial cells and NIH3T3 cells impairs the expression of Alkaline Phosphatase (ALP), an early marker of osteoblast differentiation regulated by Runx2 [31]. BRG1 can be recruited to the promoter region OC via transcription factor C/EBP $\beta$  and activate the expression of OC mediated by RNA polymerase II [32]. During osteogenesis, Osx can recruit p300 and Brg1 to the promoter of its target genes to form transcriptionally active complex *in vivo*, which is enhanced by p38 [33]. Moreover, C/EBP $\beta$ -LAP\* can interact with both Brg1 and Baf47 and bind to the Ric-8B promoter in differentiating osteoblasts, which causes repressed Ric-8B expression and osteogenesis [34]. The Retino Blastoma (RB) protein family plays a major function in the osteoblast lineage differentiation of MSC through directly target several osteogenic genes, including ALP. Both pRB and p107 can recruit BRG1 and RNA polymerase II to occupy the promoter region ALP gene and activate its transcription. However, the binding activities between pRB and p107 are mainly not overlapped [35].

### SWI/SNF and adipogenesis

MSC can be induced to differentiate into adipocytes under appropriate conditions. The commitment of adipogenic lineage requires upregulation of a variety of genes, which are largely mediated by the three classes of transcription factors, including peroxisome proliferator-activated receptor- $\gamma$  (PPAR $\gamma$ ), CCAAT/enhancer binding proteins (C/EBPs) and basic helix-loop-helix family (ADD1/SREBP1c) [36]. Chromatin remodeling is an important step in promoter

activation during adipogenic lineage commitment and differentiation. For example, X/EBP1 $\alpha$ , X/EBP1 $\beta$  and PPAR $\gamma$ 2 require the modification of the chromatin structure by SWI/SNF ATPases BRG1 and hBRM to induce the adipogenic differentiation of uncommitted fibroblasts [38,39]. Moreover, depletion of BAF47, one of the core subunit of SWI/SNF, in murine 3T3-L1 preadipocytes and human MSC abrogates adipocyte differentiation through cooperation with C/EBP $\beta$  and PPAR $\gamma$ 2 [40]. Study has shown that binding of Brg1 to PPAR $\gamma$  promoter can be regulated by Prmt4 or Prmt5, members of protein arginine methyltransferases [40-43]. Furthermore, disruption of *Brd2* expression in mouse leads to elevated adiponectin expression in adipose tissue, suggesting Brd2 plays an important role in adipogenic regulation. Indeed, Brd2 can repress the activity of PPAR $\gamma$  and inhibit adipogenesis of 3T3-L1 pre-adipocytes [44]. Characterization of genes associated with adipocyte development using microarray analysis revealed that BAF47 is upregulated in this induction. Importantly, even though BRM is dispensable for normal development, this gene is critical for balance of lineage selection between osteoblasts and adipocytes of MSC. BRM depletion favors osteoblast lineage of MSC over adipocyte lineage, which is confirmed by BRM knockout mice that display resistance to age-related osteoporosis [45].

### SWI/SNF and chondrogenesis

During developmental stage, chondrocytes derive from mesenchymal progenitors and express cartilage-specific collagens II, IX and XI. The differentiation of chondrocytes is subject to complex regulation by interplay of the transforming growth factor  $\beta$ , Wnt, BMP and fibroblast growth factor (FGF) signaling pathways [46]. The current known master transcription factor in regulating chondrogenesis is Sox9, which is expressed predominantly in the mesenchymal condensations during the early development of embryonic skeletons [47]. In mouse chimaeras, Sox9<sup>-/-</sup> cells are excluded from all chondrogenic tissues but are present as juxtaposed mesenchymal cells that show no expression of the chondrocyte-specific markers [48]. Previous studies showed that SWI/SNF-mediated chromatin remodeling complexes are enriched in cartilage-expressed transcripts, which are identified by comprehensive mapping [49]. Through downstream mediator Sp1, BMP2 can induce expression of FGF receptor 3 (FGFR3), which is essential for cartilage development. Importantly, Brg1 is shown selectively remodel a Sp1-binding sites-containing chromatin region locates in the transcription start site of FGFR3 and induces FGFR3 expression [50].

### SWI/SNF and cardiomyocytes

Adult cardiomyocytes have little or no regenerative ability. It is challenging to efficiently differentiate multipotential stem cells into cardiomyocytes. In general, FGF2, Activin A, BMP4, vascular endothelial growth factor (VEGF) and the Wnt inhibitor DKK-1 can promote the differentiation of cardiomyocytes [52]. Multiple transcription factors, including GATA4, MEF-2, NKX2.5, HAND1, HAND2 and TEAD, are predominantly expressed in myocardium and regulate the expression of cardiac genes. Study has found that BRG1 can interact with LIM-only protein CRP2 and induce expression of smooth muscle genes in adult cardiomyocytes [52]. Deficiency of BAF250a causes early development arrest and missing of mesoderm layer in mouse embryos [22,53]. BAF250a is required for mouse ESC to differentiate into mesoderm-derived functional cardiomyocytes and adipocytes, while dispensable for ectoderm-derived neuron formation [22]. Deletion of Baf250a in the sinoartial node, the region responsible for heart beating, leads to de-repression of Nkx2.5 and causes sick sinus diseases, indicating a critical role of Baf250a in normal heart function

[54]. Mechanistically, BAF250a directly interacts with subunits of Nucleosome Remodeling and Histone Deacetylase (NURD) and occupied the regulatory regions of cardiomyocyte associated genes [53]. In addition, Baf60c can collaborate with cardiomyocyte-specific transcription factors Gata4 and Tbx5 to promote the cardiomyocyte differentiation when Nodal/BMP signaling pathways are inhibited [55].

### SWI/SNF and hepatocytes

Hepatocyte-like cells generated from MSCs serve as a promising cell source for liver regeneration or tissue engineering. However, how to improve the efficacy and consistency of hepatic differentiation from MSCs remains challenging. Previous studies have revealed that The SWI/SNF chromatin-remodeling complex is essential for hepatocyte differentiation. For example, study has showed a differentiation-specific up-regulation of BRM expression in hepatocytes and C2C12 cells, which is mediated by transcription factors C/EBP $\beta$ , GATA2 and GATA3 [56]. During the early stage of liver development, BRG1 or BRM can repress the expression of tryptophan oxygenase, a late stage-specific gene [57]. As liver cells differentiate, the expression of BRM is gradually up-regulated while BRG1 is down-regulated. Deficiency of BRG1 and BRM leads to compromised expression of albumin gene in hepatocytes. BRM and BRG1 directly bind to the promoter region of the albumin gene, along with RB family proteins and C/EBP $\alpha$  [58]. The sumoylation of C/EBP $\alpha$  can block the interaction between C/EBP $\alpha$  and BRG1, which inhibits the proliferation of hepatocytes [59]. BRG1 and BRM are also involved in transcription of ATP-binding cassette transporter A1 (ABCA1) through associated with liver X receptor (LXR) and retinoid X receptor (RXR) in Hep3B cells, a hepatocyte cell line [60]. Deletion of BAF47 in mouse hepatocytes leads to neonatal lethality due to severe hypoglycemia, glycogen storage failure and impaired energetic metabolism. Loss of BAF47 is accompanied by downregulation of a majority of liver development genes [61]. BAF60a, together with PGC-1 $\alpha$ , can induce the expression of PPAR $\alpha$  target genes and stimulate fatty  $\beta$ -oxidation in hepatocyte [62].

### SWI/SNF and neurogenesis

MSC have a capacity to differentiate into neurons and astrocytes when treated with retinoic acid and brain-derived neurotrophic factors [63]. Previous reports have demonstrated that mutations in Brg1 and its associated proteins lead to development of neuronal disorders [64, 65]. During neuronal differentiation, BAF53A is repressed and replaced by BAF53B, suggesting SWI/SNF activity is tightly regulated for proper development of neuron [66]. Indeed, conditional depletion of BAF subunits in mice found that the Brg1 and BAF complex play an essential role at various steps during neurogenesis [66]. These mice display a neonatal lethal phenotype and have thin cortices. Specific depletion of BAF53b showed dramatic defects in dendritic development and memory [67]. Brg1 and other BAF subunits mutant mice showed decrease of proliferation in neural progenitors and growth retardation in cerebellum [68, 69]. These genetic studies provide useful insights to BAF functions in neurogenesis and are helpful for designing more efficient strategy for differentiation of MSC to neural lineage.

### Small-molecules targeting SWI/SNF complex

Overall, our understanding of epigenetic regulation on MSC differentiation could be beneficial for developing better strategy of regenerative medicine. Targeting BRM in MSC display a dramatic effect on promoting osteoblast lineage rather than adipogenic lineage, demonstrating great potential of SWI/SNF chromatin-remodeling complex as therapeutic targets for stem cell-based regenerative medicine

and the treatment of human metabolic disease such as osteoporosis and obesity [45]. Indeed, the screening of small cell-permeable inhibitors of SWI/SNF chromatin-remodeling complex has been intensively pursued over the last decade. For instance, inhibitor I-BET151 can specifically bind to the bromodomains of BRD3/4 and inhibit the transcription of target genes [70]. Importantly, many subunits of SWI/SNF chromatin-remodeling complex contain this druggable bromodomains, including BRG1, BAF190B, BRD9 and BAF180. Hence, the strategy of using small molecule inhibitors in combination with lineage-specific induced medium will help the cell-based therapeutic approach generating promising results.

### Conclusion

As a solution for the low regenerative capacity of adult tissues, MSC has attracted many researchers' attention for its self-renewable and pliable traits. By using MSC, donor site morbidity and immune reactions can be overcome. A critical caveat in the field is the relatively poor understanding of the molecular mechanisms that governing MSC self-renewal and lineage specification. Here, we propose that differentiation efficiency of MSC into cell types of interest will be dramatically improved by altering the composition of SWI/SNF components. As discussed above, ectopic expression of subunits of SWI/SNF complex can lead to activation of tissue-specific transcriptional program and drive cell fate selection or convert one cell type into another. In addition, more and more data suggest that the combinatorial composition of mutually exclusive BAF members in different SWI/SNF complex determine their binding ability for lineage-specific transcription factors. Pharmacological targeting of the SWI/SNF complex of MSC and its progeny represents a novel strategy for enhancing tissue-engineering and cell-based therapy.

### References

1. Pittenger MF, Mackay AM, Beck SC, Jaiswal RK, Douglas R, et al. (1999) Multilineage potential of adult human mesenchymal stem cells. *Science* 284: 143-147.
2. Jiang Y, Jahagirdar BN, Reinhardt RL, Schwartz RE, Keene CD, et al. (2002) Pluripotency of mesenchymal stem cells derived from adult marrow. *Nature* 418: 41-49.
3. Mohammadian M, Shamsasenjan K, Lotfi Nezhad P, Talebi M, Jahedi M, et al. (2013) Mesenchymal stem cells: new aspect in cell-based regenerative therapy. *Adv Pharm Bull* 3: 433-437.
4. Orlic D, Kajstura J, Chimenti S, Jakoniuk I, Anderson SM, et al. (2001) Bone marrow cells regenerate infarcted myocardium. *Nature* 410: 701-705.
5. Caplan AI (2005) Review: mesenchymal stem cells: cell-based reconstructive therapy in orthopedics. *Tissue Eng* 11: 1198-1211.
6. Theise ND, Nimmakayalu M, Gardner R, Illei PB, Morgan G, et al. (2000) Liver from bone marrow in humans. *Hepatology* 32: 11-16.
7. Brazelton TR, Rossi FM, Keshet GI, Blau HM (2000) From marrow to brain: expression of neuronal phenotypes in adult mice. *Science* 290: 1775-1779.
8. Lin Y, Weisdorf DJ, Solovey A, Hebbel RP (2000) Origins of circulating endothelial cells and endothelial outgrowth from blood. *J Clin Invest* 105: 71-77.
9. Augello A, De Bari C (2010) The regulation of differentiation in mesenchymal stem cells. *Hum Gene Ther* 21: 1226-1238.
10. Conaway JW (2012) Introduction to theme "Chromatin, epigenetics, and transcription". *Annu Rev Biochem* 81: 61-64.
11. Bartholomew B (2014) Regulating the chromatin landscape: structural and mechanistic perspectives. *Annu Rev Biochem* 83: 671-696.
12. Becker PB, Hörz W (2002) ATP-dependent nucleosome remodeling. *Annu Rev Biochem* 71: 247-273.
13. Clapier CR, Cairns BR (2009) The biology of chromatin remodeling complexes. *Annu Rev Biochem* 78: 273-304.

14. Hargreaves DC, Crabtree GR (2011) ATP-dependent chromatin remodeling: genetics, genomics and mechanisms. *Cell Res* 21: 396-420.
15. Wilson BG, Roberts CW (2011) SWI/SNF nucleosome remodellers and cancer. *Nat Rev Cancer* 11: 481-492.
16. Tang L, Nogales E, Ciferri C (2010) Structure and function of SWI/SNF chromatin remodeling complexes and mechanistic implications for transcription. *Prog Biophys Mol Biol* 102: 122-128.
17. Euskirchen G, Auerbach RK, Snyder M (2012) SWI/SNF chromatin-remodeling factors: multiscale analyses and diverse functions. *J Biol Chem* 287: 30897-30905.
18. Tolstorukov MY, Sansam CG, Lu P, Koellhoffer EC, Helming KC, et al. (2013) Swi/Snf chromatin remodeling/tumor suppressor complex establishes nucleosome occupancy at target promoters. *Proc Natl Acad Sci U S A* 110: 10165-10170.
19. Bultman SJ, Gebuhr TC, Magnuson T. (2005) A Brg1 mutation that uncouples ATPase activity from chromatin remodeling reveals an essential role for SWI/SNF-related complexes in beta-globin expression and erythroid development. *Genes Dev* 19: 2849-2861.
20. Kim JK, Huh SO, Choi H, Lee KS, Shin D, et al. (2001) Srg3, a mouse homolog of yeast SWI3, is essential for early embryogenesis and involved in brain development. *Mol Cell Biol* 21: 7787-7795.
21. Klochendler-Yeivin A, Fiette L, Barra J, Muchardt C, Babinet C, et al. (2000) The murine SNF5/INI1 chromatin remodeling factor is essential for embryonic development and tumor suppression. *EMBO Rep* 1: 500-506.
22. Gao X, Tate P, Hu P, Tjian R, Skarnes WC, et al. (2008) ES cell pluripotency and germ-layer formation require the SWI/SNF chromatin remodeling component BAF250a. *Proc Natl Acad Sci U S A* 105: 6656-6661.
23. Ho L, Jothi R, Ronan JL, Cui K, Zhao K, et al. (2009) An embryonic stem cell chromatin remodeling complex, esBAF, is an essential component of the core pluripotency transcriptional network. *Proc Natl Acad Sci U S A* 106: 5187-5191.
24. Kaeser MD, Aslanian A, Dong MQ, Yates JR 3rd, Emerson BM (2008) BRD7, a novel PBAF-specific SWI/SNF subunit, is required for target gene activation and repression in embryonic stem cells. *J Biol Chem* 283: 32254-32263.
25. Ho L, Ronan JL, Wu J, Staahl BT, Chen L, et al. (2009) An embryonic stem cell chromatin remodeling complex, esBAF, is essential for embryonic stem cell self-renewal and pluripotency. *Proc Natl Acad Sci U S A* 106: 5181-5186.
26. Ho L, Miller EL, Ronan JL, Ho WQ, Jothi R, et al. (2011) esBAF facilitates pluripotency by conditioning the genome for LIF/STAT3 signalling and by regulating polycomb function. *Nat Cell Biol* 13: 903-913.
27. Squillaro T, Severino V, Alessio N, Farina A, Di Bernardo G, et al. (2015) De-regulated expression of the BRG1 chromatin remodeling factor in bone marrow mesenchymal stromal cells induces senescence associated with the silencing of NANOG and changes in the levels of chromatin proteins. *Cell Cycle* 14:1315-26.
28. Wu JI (2012) Diverse functions of ATP-dependent chromatin remodeling complexes in development and cancer. *Acta Biochim Biophys Sin (Shanghai)* 44: 54-69.
29. Minguell JJ, Erices A, Conget P (2001) Mesenchymal stem cells. *Exp Biol Med (Maywood)* 226: 507-520.
30. Fakhry M, Hamade E, Badran B, Buchet R, Magne D (2013) Molecular mechanisms of mesenchymal stem cell differentiation towards osteoblasts. *World J Stem Cells* 5: 136-148.
31. Young DW, Pratap J, Javed A, Weiner B, Ohkawa Y, et al. (2005) SWI/SNF chromatin remodeling complex is obligatory for BMP2-induced, Runx2-dependent skeletal gene expression that controls osteoblast differentiation. *J Cell Biochem* 94: 720-730.
32. Villagra A, Cruzat F, Carvallo L, Paredes R, Olate J, et al. (2006) Chromatin remodeling and transcriptional activity of the bone-specific osteocalcin gene require CCAAT/enhancer-binding protein  $\gamma$ -dependent recruitment of SWI/SNF activity. *J. Biol. Chem* 281: 22695-22706.
33. Ortuño MJ, Ruiz-Gaspà S, Rodríguez-Carballo E, Susperregui AR, Bartrons R, et al. (2010) p38 regulates expression of osteoblast-specific genes by phosphorylation of osterix. *J Biol Chem* 285: 31985-31994.
34. Aguilar R, Grandy R, Meza D, Sepulveda H, Pihan P, et al. (2014) A functional N-terminal domain in C/EBP $\beta$ -LAP\* is required for interacting with SWI/SNF and to repress Ric-8B gene transcription in osteoblasts. *J Cell Physiol* 229: 1521-1528.
35. Flowers S, Patel PJ, Gleicher S1, Amer K1, Himelman E1, et al. (2014) p107-Dependent recruitment of SWI/SNF to the alkaline phosphatase promoter during osteoblast differentiation. *Bone* 69: 47-54.
36. Scott MA, Nguyen VT, Levi B, James AW (2011) Current methods of adipogenic differentiation of mesenchymal stem cells. *Stem Cells Dev* 20: 1793-1804.
37. Pedersen TA, Kowenz-Leutz E, Leutz A, Nerlov C (2001) Cooperation between C/EBP $\alpha$  TBP/TFIIB and SWI/SNF recruiting domains is required for adipocyte differentiation. *Genes Dev* 15: 3208-3216.
38. Salma N, Xiao H, Mueller E, Imbalzano AN. (2004) Temporal recruitment of transcription factors and SWI/SNF chromatin-remodeling enzymes during adipogenic induction of the peroxisome proliferator-activated receptor gamma nuclear hormone receptor. *Mol Cell Biol* 24: 651-6663.
39. Caramel J, Medjkane S, Quignon F, Delattre O (2008) The requirement for SNF5/INI1 in adipocyte differentiation highlights new features of malignant rhabdoid tumors. *Oncogene* 27: 2035-2044.
40. LeBlanc SE, Konda S, Wu Q, Hu YJ, Osowski CM, et al. (2012) Protein arginine methyltransferase 5 (Prmt5) promotes gene expression of peroxisome proliferator-activated receptor  $\beta$ 2 (PPAR $\beta$ 2) and its target genes during adipogenesis. *Mol Endocrinol* 26: 583-597.
41. Yadav N, Cheng D, Richard S, Morel M, Iyer VR, et al. (2008) CARM1 promotes adipocyte differentiation by coactivating PPARgamma. *EMBO Rep* 9: 193-198.
42. Pal S, Yun R, Datta A, Lacomis L, Erdjument-Bromage H, et al. (2003) mSin3A/histone deacetylase 2- and PRMT5-containing Brg1 complex is involved in transcriptional repression of the Myc target gene cad. *Mol Cell Biol* 23: 7475-7487.
43. Xu W, Cho H, Kadam S, Banayo EM, Anderson S, et al. (2004) A methylation-mediator complex in hormone signaling. *Genes Dev* 18: 144-156.
44. Wang F, Liu H, Blanton WP, Belkina A, Lebrasseur NK, et al. (2009) Brd2 disruption in mice causes severe obesity without Type 2 diabetes. *Biochem J* 425: 71-83.
45. Nguyen KH, Xu F1, Flowers S1, Williams EA1, Fritton JC1, et al. (2015) SWI/SNF-Mediated Lineage Determination in Mesenchymal Stem Cells Confers Resistance to Osteoporosis. *Stem Cells* .
46. Pei M, Chen D, Li J, Wei L (2009) Histone deacetylase 4 promotes TGF-beta1-induced synovium-derived stem cell chondrogenesis but inhibits chondrogenically differentiated stem cell hypertrophy. *Differentiation* 78: 260-268.
47. Wright E, Hargrave MR, Christiansen J, Cooper L, Kun J, et al. (1995) The Sry-related gene Sox9 is expressed during chondrogenesis in mouse embryos. *Nat Genet* 9: 15-20.
48. Bi W, Deng JM, Zhang Z, Behringer RR, de Crombrughe B (1999) Sox9 is required for cartilage formation. *Nat Genet* 22: 85-89.
49. Yager TD, Dempsey AA, Tang H, Stamatou D, Chao S, et al. (2004) First comprehensive mapping of cartilage transcripts to the human genome. *Genomics* 84: 524-535.
50. Sun F, Chen Q, Yang S, Pan Q, Ma J, et al. (2009) Remodeling of chromatin structure within the promoter is important for bmp-2-induced fgfr3 expression. *Nucleic Acids Res* 37: 3897-3911.
51. Yang L, Soonpaa MH, Adler ED, Roepke TK, Kattman SJ, et al. (2008) Human cardiovascular progenitor cells develop from a KDR+ embryonic-stem-cell-derived population. *Nature* 453: 524-528.
52. Chang DF, Belaguli NS, Chang J, Schwartz RJ (2007) LIM-only protein, CRP2, switched on smooth muscle gene activity in adult cardiac myocytes. *Proc Natl Acad Sci U S A* 104: 157-162.
53. Singh AP, Archer TK. (2014) Analysis of the SWI/SNF chromatin-remodeling complex during early heart development and BAF250a repression cardiac gene transcription during P19 cell differentiation. *Nucleic Acids Res* 42: 2958-2975.
54. Wu M, Peng S1, Yang J2, Tu Z2, Cai X3, et al. (2014) Baf250a orchestrates an epigenetic pathway to repress the Nkx2.5-directed contractile cardiomyocyte program in the sinoatrial node. *Cell Res* 24: 1201-1213.
55. Cai W, Albini S, Wei K, Willems E, Guzzo RM, et al. (2013) Coordinate Nodal and BMP inhibition directs Baf60c-dependent cardiomyocyte commitment. *Genes Dev* 27: 2332-2344.

56. Itoh T, Miyake K, Iijima S (2008) Differentiation-specific expression of chromatin remodeling factor BRM. *Biochem Biophys Res Commun* 366: 827-833.
57. Inayoshi Y, Kaneoka H, Machida Y, Terajima M, Dohda T, et al. (2005) Repression of GR-mediated expression of the tryptophan oxygenase gene by the SWI/SNF complex during liver development. *J Biochem* 138: 457-465.
58. Inayoshi Y, Miyake K, Machida Y, Kaneoka H, Terajima M, et al. (2006) Mammalian chromatin remodeling complex SWI/SNF is essential for enhanced expression of the albumin gene during liver development. *J Biochem* 139: 177-188.
59. Sato Y, Miyake K, Kaneoka H, Iijima S (2006) Sumoylation of CCAAT/enhancer-binding protein alpha and its functional roles in hepatocyte differentiation. *J Biol Chem* 281: 21629-21639.
60. Huuskonen J1, Vishnu M, Fielding PE, Fielding CJ (2005) Activation of ATP-binding cassette transporter A1 transcription by chromatin remodeling complex. *Arterioscler Thromb Vasc Biol* 25: 1180-1185.
61. Gresh L, Bourachot B, Reimann A, Guigas B, Fiette L, et al. (2005) The SWI/SNF chromatin-remodeling complex subunit SNF5 is essential for hepatocyte differentiation. *EMBO J* 24: 3313-3324.
62. Li S, Liu C, Li N, Hao T, Han T, et al. (2008) Genome-wide coactivation analysis of PGC-1alpha identifies BAF60a as a regulator of hepatic lipid metabolism. *Cell Metab* 8: 105-117.
63. Sanchez-Ramos J, Song S, Cardozo-Pelaez F, Hazzi C, Stedeford T, et al. (2000) Adult bone marrow stromal cells differentiate into neural cells in vitro. *Exp Neurol* 164: 247-256.
64. Santen GW, Aten E, Sun Y, Almomani R, Gilissen C, et al. (2012) Mutations in SWI/SNF chromatin remodeling complex gene ARID1B cause Coffin-Siris syndrome. *Nat Genet* 44: 379-380.
65. Tsurusaki Y, Okamoto N, Ohashi H, Kosho T, Imai Y, et al. (2012) Mutations affecting components of the SWI/SNF complex cause Coffin-Siris syndrome. *Nat Genet* 44: 376-378.
66. Lessard J, Wu JI, Ranish JA, Wan M, Winslow MM, et al. (2007) An essential switch in subunit composition of a chromatin remodeling complex during neural development. *Neuron* 55: 201-215.
67. Vogel-Ciernia A, Matheos DP, Barrett RM, Kramár EA, Azzawi S, et al. (2013) The neuron-specific chromatin regulatory subunit BAF53b is necessary for synaptic plasticity and memory. *Nat Neurosci* 16: 552-561.
68. Moreno N, Schmidt C2, Ahlfeld J3, Pöschl J3, Dittmar S1, et al. (2014) Loss of Smarck proteins impairs cerebellar development. *J Neurosci* 34: 13486-13491.
69. Zhan X, Shi X, Zhang Z, Chen Y, Wu JI (2011) Dual role of Brg chromatin remodeling factor in Sonic hedgehog signaling during neural development. *Proc Natl Acad Sci U S A* 108: 12758-12763.
70. Dawson MA, Prinjha RK, Dittmann A, Giotopoulos G, Bantscheff M, et al. (2011) Inhibition of BET recruitment to chromatin as an effective treatment for MLL-fusion leukaemia. *Nature* 478: 529-533.