

# Synthesis and Antimicrobial Activity of New Synthesized Benzimidazole Derivatives and their Acyclic Nucleoside Analogues

Hamada H Amer<sup>1,2\*</sup>, SM El-Kousy<sup>3</sup>, WM Salama<sup>3</sup> and A H-H Sheleby<sup>2</sup>

<sup>1</sup>Chemistry Department, Faculty of Medical and Applied Science, Turabba Campus, Taif University, Saudi Arabia

<sup>2</sup>Faculty of Veterinary Medicine, Sadat City University, Sadat City, Menoufia, Egypt

<sup>3</sup>Chemistry Department, Faculty of Science, Menoufia University, Egypt

## Abstract

New benzimidazole derivatives and their N-substituted acyclic nucleoside analogues were prepared. The synthesized compounds were tested for their antimicrobial activity against *Escherichia coli*, *Staphylococcus aureus*, *Micrococcus*, *Salmonella typhi* and *Salmonella para typhi*. The synthesized compounds were tested also against fungi species such as *Aspergillus flavus*, *Aspergillus fumigates*, *Aspergillus ochraceus* and *Candida albicans*. Most of tested compounds exhibited moderate to high antimicrobial activity while few compounds were found to exhibit little or no activity against the tested microorganisms.

**Keywords:** Benzimidazole derivatives; Sugar hydrazones; Acyclic nucleosides; Antimicrobial activity

## Introduction

Imidazoles are common scaffolds in highly significant biomolecules, including biotin, the essential amino acid histidine, histamine, the pilocarpine alkaloids [1] and other alkaloids, which have been shown to exhibit interesting biological activities such as antimicrobial, anticryptococcal, inhibition of nitric oxide synthase and cytotoxic activities [2]. Imidazole derivatives have also been found to possess many pharmacological properties and are widely implicated in biochemical processes. Members of this class of diazoles are known to possess NO synthase inhibition [3], antibiotic [4], antifungal [5-8] and neuropeptide Y antagonistic activities [9]. In addition, these heterocycles include several inhibitors of p38 MAP kinases [10-13] which are thought to be involved in a variety of inflammatory and immunological disorders, and some derivatives such as mitronidazole, etomidate and ketoconazole which have found application in drug therapy [14]. Recently certain imidazole based compounds were reported to possess antimicrobial activities [15]. Further recent literature revealed that N1-substitution of imidazoles improves the antibacterial activity [16]. Among the five-membered nitrogen heterocycles, the 1,3,4-oxadiazoles are associated with broad spectrum of biological activities [17-19]. Their derivatives have been known to possess antibacterial [20], antimicrobial [21], insecticidal [22], herbicidal, fungicidal [23], anti-inflammatory [24], hypoglycemic [25] characteristics, antiviral [26], and anti-tumour activities [27]. On the other hand, the acyclic C-nucleoside analogues possess a wide range of biological properties, including antibiotic, antiviral, and anti-tumour activities [28-37]. Consequently, newly synthesized compounds containing imidazole and 1,3,4-oxadiazole moieties as well as their substituted sugar derivatives will be expected of enhanced biological activities. Owing to the above facts and our interest in the attachment of carbohydrate residues to newly synthesized heterocycles [38-40] searching for potent leads as antimicrobial agents, our aim is the synthesis and antimicrobial evaluation of new substituted 5-nitroimidazol and their acyclic nucleoside analogues.

## Experimental

Melting points were determined with a Kofler block apparatus and are uncorrected. The IR spectra were recorded on a Perkin-Elmer model 1720 FTIR spectrometer for KBr discs. NMR spectra were recorded on a Varian Gemini 200 NMR Spectrometer at 300 MHz for <sup>1</sup>H NMR with TMS as a standard. The progress of the reactions was monitored by TLC using aluminum silica gel plates 60 F 245. Elemental analyses

were performed at the Microanalytical data centre at Faculty of science, Cairo University, Egypt.

### 1-Ethoxycarbonylmethyl-2-methylbenzimidazole (172)

Ethylchloroacetate (1.4 g, 0.01 mol) was added to a solution of 2-methylbenzimidazol (171) (1.32 g, 0.01 mol) in dry acetone (30 ml) and anh. Potassium carbonate (1.38 g, 0.01 mol). The reaction mixture was heated under reflux for 5 h, poured on crushed-ice, filtered off, and recrystallized from ethanol to give 172 as a white needles (1.82 g, 83%); m.p.=115-117°C. IR spectrum (KBr),  $\nu$ , cm<sup>-1</sup>: 1730 (C=O), 1455 (CH<sub>2</sub>), 1375 (CH<sub>3</sub>). <sup>1</sup>H NMR spectrum (300 MHz, DMSO-d<sub>6</sub>),  $\delta$ , ppm (J, Hz): 1.32 (t, 3H, J=5.6 Hz, CH<sub>3</sub>CH<sub>2</sub>), 1.87(s,3H, CH<sub>3</sub>), 4.20 (q, 2H, J=5.6 Hz, CH<sub>3</sub>CH<sub>2</sub>), 5.08 (s, 2H, NCH<sub>2</sub>), 7.12- 8.14(m, 4H, Ar-H). Mass spectrum, m/z (I, %): 218[M<sup>+</sup>] (20).

### 1-(2-Methylbenzimidazol-1-yl) acetic acid hydrazide (173)

A solution of the respective ester 172 (2.18 g, 0.01 mol) in absolute ethanol (30 ml) and hydrazine hydrate (1.5 g, 0.03 mol) was refluxed for 3 h. The solvent was removed under reduced pressure and the remaining precipitate was collected, dried, and recrystallized from ethanol to afford 173 as a pale yellow powder (1.7 g, 83%); m.p.=88-90°C. IR spectrum (KBr),  $\nu$ , cm<sup>-1</sup>: 1637 (C=O), 3351 (NH), 3567 (NH<sub>2</sub>). Mass spectrum, m/z (I, %): 204 [M<sup>+</sup>] (100).

### General procedure for the synthesis of Sugar (2-methylbenzimidazol-1-yl)acetylhydrazones 178-181

Hydrazide 173 (0.65 g, 0.032 mol) in ethanol (10 ml) was added to a stirred solution of the respective monosaccharide (0.032 mol) in water (1 ml) and glacial acetic acid (1 ml). The mixture was heated under reflux, the excess of ethanol was removed under reduced pressure

\*Corresponding author: Hamada H Amer, Assistant Professor, Chemistry Department, Faculty of Medical and Applied Science, Taif University, Saudi Arabia, Tel: 00966537348857; E-mail: [dr.hamada1435@gmail.com](mailto:dr.hamada1435@gmail.com)

Received: December 16, 2015; Accepted: January 06, 2016; Published: January 12, 2016

Citation: Amer HH, El-Kousy SM, Salama WM, H-H Sheleby A (2016) Synthesis and Antimicrobial Activity of New Synthesized Benzimidazole Derivatives and their Acyclic Nucleoside Analogues. Organic Chem Curr Res 5: 157. doi:10.4172/2161-0401.1000157

Copyright: © 2016 Amer HH, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

and the residue was triturated with (10 ml) diethyl ether to produce compounds as brown viscous material to afford the corresponding sugar hydrazones **178-181**.

#### L(-)-Arabinose(2-methylbenzimidazol-1-yl)acetylhydrazone (178)

Pale brown gum (0.75 g, 70%); IR spectrum (KBr),  $\nu$ ,  $\text{cm}^{-1}$ : 3350 (OH), 3469 (NH).  $^1\text{H}$  NMR spectrum (300 MHz, DMSO- $d_6$ ),  $\delta$ , ppm ( $J$ , Hz): 1.00 (s, 1H,  $\text{CH}_3$ ), 3.25-3.74 (m, 5H, H-2', H-3', H-4', H-5', H-5''), 4.45 (brs, 2H, 2xOH), 4.60 (s, 2H,  $\text{CH}_2$ ), 4.96 (brs, 2H, 2xOH), 7.09 (d, 1H,  $J=2.5$  Hz, H-1'), 7.12-7.45 (m, 4H, Ar-H), 9.63 (brs, 1H, NH).

#### D-(+)-Xylose(2-methylbenzimidazol-1-yl)acetylhydrazone (179)

Pale brown gum (0.82 g, 76%); IR spectrum (KBr),  $\nu$ ,  $\text{cm}^{-1}$ : 3373 (OH), 3426 (NH).  $^1\text{H}$  NMR spectrum (300 MHz, DMSO- $d_6$ ),  $\delta$ , ppm ( $J$ , Hz): 0.96 (s, 1H,  $\text{CH}_3$ ), 3.21-3.70 (m, 5H, H-2', H-3', H-4', H-5', H-5''), 4.43 (bs, 2H, 2xOH), 4.65 (s, 2H,  $\text{CH}_2$ ), 5.22 (bs, 2H, 2xOH), 7.09 (d, 1H,  $J=2.5$  Hz, H-1'), 7.12-7.45 (m, 4H, Ar-H), 9.63 (brs, 1H, NH).

#### D-(+)-Galactose(2-methylbenzimidazol-1-yl)acetylhydrazone (180)

Pale brown gum (0.95 g, 81%); IR spectrum (KBr),  $\nu$ ,  $\text{cm}^{-1}$ : 3295(OH), 3402 (NH).  $^1\text{H}$  NMR spectrum (300 MHz, DMSO- $d_6$ ),  $\delta$ , ppm ( $J$ , Hz): 1.00 (s, 3H,  $\text{CH}_3$ ), 3.24-3.85 (m, 5H, H-3', H-4', H-5', H-6', H-6''), 4.19 (m, 1H, H-2'), 4.42 (bs, 1H, OH), 4.55 (bs, 2H, 2xOH), 4.62 (s, 2H,  $\text{CH}_2$ ), 5.00 (bs, 2H, 2xOH), 7.11 (d, 1H,  $J=2.5$  Hz, H-1'), 7.38-7.94 (m, 4H, Ar-H), 8.88 (brs, 1H, NH). Mass spectrum,  $m/z$  ( $I$ , %): 366 [ $\text{M}^+$ ] (100).

#### D-(+)-Mannose(2-methylbenzimidazol-1-yl)acetylhydrazone (181)

Pale brown gum (0.85 g, 72%); IR spectrum (KBr),  $\nu$ ,  $\text{cm}^{-1}$ : 3295(OH), 3402 (NH).  $^1\text{H}$  NMR spectrum (300 MHz, DMSO- $d_6$ ),  $\delta$ , ppm ( $J$ , Hz): 1.00 (s, 3H,  $\text{CH}_3$ ), 3.20-3.65 (m, 5H, H-3', H-4', H-5', H-6', H-6''), 4.19 (m, 1H, H-2'), 4.44 (bs, 1H, OH), 4.55 (bs, 2H, 2xOH), 4.65 (s, 2H,  $\text{CH}_2$ ), 5.00 (bs, 2H, 2xOH), 7.11 (d, 1H,  $J=2.5$  Hz, H-1'), 7.38-7.94 (m, 4H, Ar-H), 9.00 (brs, 1H, NH).

#### General procedure for the synthesis of sugar of tetra-O-acetyl- and penta-O-acetyl(2-methylbenzimidazol-1-yl)acetylhydrazones 183-187

Acetic anhydride (3.06 g, 0.03 mol) was added to a solution of sugar hydrazones **178-181** (0.001 mol) in pyridine (7 ml) with stirring at room temperature for overnight. The mixture was cooled and poured on crushed ice. Hydrochloric acid was added with stirring until the odor of pyridine was removed and extracted by chloroform. The product was separated in pure form as brown viscous material to afford **182-185**.

#### 2,3,4,5-Tetra-O-acetyl-L(-)-arabinose(2-methylbenzimidazol-1-yl)acetylhydrazone (182)

Pale yellow gum (0.54 g, 60%);  $^1\text{H}$  NMR spectrum (300 MHz, DMSO- $d_6$ ),  $\delta$ , ppm ( $J$ , Hz): 0.87 (s, 3H,  $\text{CH}_3$ ), 1.99, 2.01, 2.10, 2.49 (4s, 12H, 4x $\text{CH}_3\text{CO}$ ), 3.40, 3.41 (2m, 2H, H-5', H-5''), 4.23 (m, 1H, H-4'), 4.60 (s, 2H,  $\text{CH}_2$ ), 4.97 (m, 1H, H-3'), 5.73 (m, 1H, H-2'), 7.26 (d, 1H,  $J=2.5$  Hz, H-1'), 7.38-7.94 (m, 4H, Ar-H), 9.43 (brs, 1H, NH).

#### 2,3,4,5-Tetra-O-acetyl-D-(+)-xylose(2-methylbenzimidazol-

#### 1-yl)acetylhydrazone (183)

Pale yellow gum (0.50 g, 66%), m.p.  $\rightarrow 300^\circ\text{C}$ . NMR spectrum (300 MHz, DMSO- $d_6$ ),  $\delta$ , ppm ( $J$ , Hz): 0.87 (s, 3H,  $\text{CH}_3$ ), 1.99, 2.01, 2.10, 2.49 (4s, 12H, 4x $\text{CH}_3\text{CO}$ ), 3.40, 3.41 (2m, 2H, H-5', H-5''), 4.23 (m, 1H, H-4'), 4.60 (s, 2H,  $\text{CH}_2$ ), 4.97 (m, 1H, H-3'), 5.73 (m, 1H, H-2'), 7.26 (d, 1H,  $J=2.5$  Hz, H-1'), 7.38-7.94 (m, 4H, Ar-H), 9.43 (brs, 1H, NH).

#### 2,3,4,5,6-Penta-O-acetyl-D-(+)-galactose(2-methylbenzimidazol-1-yl)acetylhydrazone (184)

Pale brown gum (0.53 g, 67%);  $^1\text{H}$  NMR spectrum (300 MHz, DMSO- $d_6$ ),  $\delta$ , ppm ( $J$ , Hz): 0.89 (s, 3H,  $\text{CH}_3$ ), 1.98, 2.02, 2.04, 2.07, 2.48 (5s, 15H, 5x $\text{CH}_3\text{CO}$ ), 3.40, 3.42 (2m, 2H, H-6', H-6''), 4.12 (m, 1H, H-5'), 4.14 (s, 2H,  $\text{CH}_2$ ), 5.00 (m, 1H, H-4'), 5.30 (m, 1H, H-3'), 5.49 (m, 1H, H-2'), 7.27 (d, 1H,  $J=2.5$  Hz, H-1'), 7.68-7.69 (m, 4H, Ar-H), 9.43 (brs, 1H, NH).

#### 2,3,4,5,6-Penta-O-acetyl-D-(+)-mannose(2-methylbenzimidazol-1-yl)acetylhydrazone (185)

Pale brown gum (0.60 g, 77%);  $^1\text{H}$  NMR spectrum (300 MHz, DMSO- $d_6$ ),  $\delta$ , ppm ( $J$ , Hz): 0.91 (s, 3H,  $\text{CH}_3$ ), 1.99, 2.00, 2.04, 2.10, 2.48 (5s, 15H, 5x $\text{CH}_3\text{CO}$ ), 3.40, 3.42 (2m, 2H, H-6', H-6''), 4.10 (m, 1H, H-5'), 4.52 (s, 2H,  $\text{CH}_2$ ), 5.07 (m, 1H, H-4'), 5.30 (m, 1H, H-3'), 5.49 (m, 1H, H-2'), 7.32 (d, 1H,  $J=2.5$  Hz, H-1'), 7.55-7.66 (m, 4H, Ar-H), 9.33 (brs, 1H, NH).

#### General procedure for the synthesis of 4-acetyl-5-(tetra- and penta-O-acetylalditolyl)-2-(2-methylbenzimidazol-1-yl)-1, 3, 4-oxadiazolines 186- 189

A solution of sugar hydrazones **178-181** (0.01 mol) in acetic anhydride (5 ml) was heated at  $100^\circ\text{C}$  for 5 h. The resulting solution was poured onto crushed-ice and the product that separated out was filtered off, washed with a saturated solution of sodium bicarbonate followed by water, and then dried. The products were recrystallized from ethanol to give **186-189**.

#### 4-Acetyl-5-(1,2,3,4-tetra-O-acetyl-L-arabinotetritolyl)-2-(2-methylbenzimidazol-1-yl)-1,3,4-oxadiazoline (186)

Pale yellow gum (0.53 g, 65%); IR spectrum (KBr),  $\nu$ ,  $\text{cm}^{-1}$ : 1741 ( $\text{COCH}_3$ ), 1650 ( $\text{C}=\text{N}$ ).  $^1\text{H}$  NMR spectrum (300 MHz, DMSO- $d_6$ ),  $\delta$ , ppm ( $J$ , Hz): 0.89 (s, 3H,  $\text{CH}_3$ ), 1.83, 1.90, 2.01, 2.10, 2.48 (5s, 15H, 5x $\text{CH}_3\text{CO}$ ), 3.85 (s, 2H,  $\text{CH}_2$ ), 4.01, 4.09 (2m, 2H, H-4', H-4''), 4.97 (m, 1H, H-3'), 5.18 (m, 1H, H-2'), 5.40 (dd, 1H,  $J=3.2, 6.2$  Hz, H-1'), 5.89 (d, 1H,  $J=8.8$  Hz, oxadiazoline H-5), 7.68-7.69 (m, 4H, Ar-H).

#### 4-Acetyl-5-(1,2,3,4-tetra-O-acetyl-D-xylotetritolyl)-2-(2-methylbenzimidazol-1-yl)-1,3,4-oxadiazoline (187)

Pale yellow gum (0.45 g, 55%); IR spectrum (KBr),  $\nu$ ,  $\text{cm}^{-1}$ : 1738 ( $\text{COCH}_3$ ), 1650 ( $\text{C}=\text{N}$ ).  $^1\text{H}$  NMR spectrum (300 MHz, DMSO- $d_6$ ),  $\delta$ , ppm ( $J$ , Hz): 0.89 (s, 3H,  $\text{CH}_3$ ), 1.91, 1.95, 1.97, 1.98, 2.11 (5s, 15H, 5x $\text{CH}_3\text{CO}$ ), 3.89 (s, 2H,  $\text{CH}_2$ ), 4.00, 4.12 (2m, 2H, H-4', H-4''), 5.07 (m, 1H, H-3'), 5.22 (m, 1H, H-2'), 5.47 (dd, 1H,  $J=3.2, 6.2$  Hz, H-1'), 5.89 (d, 1H,  $J=8.8$  Hz, oxadiazoline H-5), 7.68-7.69 (m, 4H, Ar-H).

#### 4-Acetyl-5-(1,2,3,4,5-penta-O-acetyl-D-galactopentitolyl)-2-(2-methylbenzimidazol-1-yl)-1,3,4-oxadiazoline (188)

Pale yellow gum (0.62 g, 73%); IR spectrum (KBr),  $\nu$ ,  $\text{cm}^{-1}$ : 1738 ( $\text{COCH}_3$ ), 1674 ( $\text{C}=\text{N}$ ).  $^1\text{H}$  NMR spectrum (300 MHz, DMSO- $d_6$ ),  $\delta$ , ppm ( $J$ , Hz): 0.89 (s, 3H,  $\text{CH}_3$ ), 1.90, 1.91, 1.95, 1.97, 1.98, 2.11

(6s,18H,6xCH<sub>3</sub>CO), 3.89 (s, 2H, CH<sub>2</sub>), 4.05, 4.12 (2m,2H,H-5',H-5''), 4.98 (m,1H,H-4'), 5.23 (m, 1H, H-3'), 5.33 (m,1H,H-2'), 5.43 (dd,1H, J=3.2, 6.2 Hz,H-1'),5.89 (d, 1H, J=8.8 Hz,oxadiazoline H-6), 7.68-7.69 (m, 4H, Ar-H).

#### 4-Acetyl-5-(1,2,3,4,5-penta-O-acetyl-D-mannopentitolyl)-2-(2-methylbenzimidazol-1-yl)-1,3,4-oxadiazole (189)

Pale yellow gum (0.58 g, 69%); IR spectrum (KBr),  $\nu$ , cm<sup>-1</sup>: 1742(COCH<sub>3</sub>), 1679 (C=N). <sup>1</sup>H NMR spectrum (300 MHz, DMSO-d<sub>6</sub>),  $\delta$ , ppm (J, Hz): 0.89(s,3H, CH<sub>3</sub>), 1.90, 1.91, 1.95, 2.03, 2.10, 2.13 (6s, 18H, 6xCH<sub>3</sub>CO), 3.95 (s, 2H, CH<sub>2</sub>), 4.0.89(s,3H, CH<sub>3</sub>), 11, 4.17 (2m, 2H, H-5',H-5''), 4.92 (m, 1H, H-4'), 5.19 (m, 1H, H-3'), 5.31 (m, 1H, H-2'), 5.43 (dd, 1H, J=3.2, 6.2 Hz, H-1'),5.89 (d, 1H, J=8.8 Hz, oxadiazoline H-6), 7.68-7.69 (m, 4H, Ar-H).

#### 2-[(2-methylbenzimidazol-1-yl)methyl]-1,3,4-oxadiazole-2-thiol (194)

To a solution of (173) (4.8 g, 0.02 mole) in ethanol (50 mL) was added a solution of potassium hydroxide (1.12 g, 0.02 mole) in water (2 mL) and carbon disulphide (5 mL). The solution was heated under reflux for 15 h. The solvent was evaporated and the residue was dissolved in water, filtered, and acidified with dilute hydrochloric acid. The precipitate was filtered off, washed with water and recrystallized from ethanol. Yield 80%; mp=200-202°C; IR (KBr):  $\nu$  3415 (NH), 1622 cm<sup>-1</sup> (C=N). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz):  $\delta$ 1.00 (s,1H, CH<sub>3</sub>), 4.58 (s, 2H, CH<sub>2</sub>), 7.97-8.22 (m, 4H, Ar-H), 12.80 (s, 1H, SH). *m/z*=246 [M<sup>+</sup>].

#### 2-(Methylthio)-2-[(2-methylbenzimidazol-1-yl)methyl]-1,3,4-oxadiazole (195)

To a solution of (194) (2.46 g, 0.01 mole) and potassium hydroxide (0.56 g, 0.01 mole) in a mixture of water (25 mL) and ethanol (10 mL), was added the methyl iodide (0.01 mole). The solution was stirred at room temperature for 4 h. The resulting precipitate was filtered off and crystallized from ethanol to give 195. Yield 71%; mp=87-89°C; IR (KBr):  $\nu$  1622 cm<sup>-1</sup> (C=N). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz):  $\delta$  0.9(s, 3H, CH<sub>3</sub>), 2.65 (s, 1H, SCH<sub>3</sub>), 5.55 (s, 2H, CH<sub>2</sub>), 7.98-8.22(m, 4H, Ar-H).

#### 2-Hydrazinyl-2-[(2-methylbenzimidazol-1-yl)methyl]-1,3,4-oxadiazole (196)

A mixture of (195) (0.01 mole) EtOH (8 mL), and N<sub>2</sub>H<sub>4</sub>.H<sub>2</sub>O (1 g, 0.02 mole) was refluxed for 4 h and the solvent was removed under reduced pressure. The remaining precipitate was collected, dried, and recrystallized from EtOH to afford compound 196. Yield 74%; m.p.=174-176°C; IR (KBr):  $\nu$  1620 cm<sup>-1</sup> (C=N). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz):  $\delta$ 1.00(s,3H, CH<sub>3</sub>), 5.58 (s, 2H, CH<sub>2</sub>), 5.80 (s, 2H, NH<sub>2</sub>), 7.95-8.22(m, 4H, Ar-H),9.45(brs, 1H, NH).

#### General procedures for the reaction of (196) with aromatic aldehydes to afford Schiff's bases 200-202

A solution of 196 (2.44 g, 0.01 mol), an aromatic aldehyde (0.01 mol) in abs. ethanol (30 mL) and glacial acetic acid (1 ml) was refluxed for 4-6 h (TLC). The solvent was evaporated under reduced pressure and the residue was filtered off and recrystallized from ethanol to afford 200-202.

#### N'-(Benzylidene)-2-[(2-methylbenzimidazol-1-yl)methyl]-1,3,4-oxadiazole (200).

yellow powder (2.78 g, 83%); m.p.=270-272°C. <sup>1</sup>H NMR spectrum (300 MHz, DMSO-d<sub>6</sub>),  $\delta$ , ppm (J, Hz):0.94(s,3H, CH<sub>3</sub>), 4.51 (s, 2H,

CH<sub>2</sub>), 7.80 (s, 1H, CH), 7.86-8.01 (m, 8H, Ar-H), 9.93 (brs, 1H, NH).

#### N'-(4-Chlorobenzylidene)-2-[(2-methylbenzimidazol-1-yl)methyl]-1,3,4-oxadiazole (201)

White powder (3.24 g, 88%); m.p.=145-147°C. <sup>1</sup>H NMR spectrum (300 MHz, DMSO-d<sub>6</sub>),  $\delta$ , ppm (J, Hz):0.94(s, 3H, CH<sub>3</sub>), 4.55 (s, 2H, CH<sub>2</sub>), 8.00 (s, 1H, CH), 8.42-8.64 (m, 8H, Ar-H), 9.93 (brs, 1H, NH). Mass spectrum, *m/z* (I, %): 366 [M<sup>+</sup>] (60).

#### N'-[5-(Methylfuran-2-yl)methylene]-2-[(2-methylbenzimidazol-1-yl)methyl]-1,3,4-oxadiazole (201).

White powder (3.10 g, 92%); m.p.=299-300°C. <sup>1</sup>H NMR spectrum (300 MHz, DMSO-d<sub>6</sub>),  $\delta$ , ppm (J, Hz):0.95(s,3H, CH<sub>3</sub>), 2.32 (s, 3H, CH<sub>3</sub>), 4.65 (s, 2H, CH<sub>2</sub>), 6.24 (s, 1H, H-4-furan), 6.79 (s, 1H, H-3-furan), 7.22-7.55 (m, 4H, Ar-H), 8.22 (s, 2H, CH=N),9.96 (brs, 1H, NH). Mass spectrum, *m/z* (I, %): 336 [M<sup>+</sup>] (23).

#### General procedure for the synthesis of sugar hydrazinyl derivatives 205,206

To a well stirred solution of the respective monosaccharides (xylose and galactose) (0.01 mole) in water (2 mL), and glacial acetic acid (0.2 mL) was added to 196 (2.44 g, 0.01 mole) in ethanol (10 mL). The mixture was heated under reflux for 3 h, the resulting solution was concentrated and left to cool. The precipitate formed was filtered off, washed with water and ethanol, then dried and recrystallized from ethanol.

#### 2-D-Xylotetritolylidenehydrazinyl)-2-((2-methylbenzimidazol-1-yl)methyl)-1,3,4-oxadiazole (205)

Yield 80%; mp=194-196°C; IR (KBr):  $\nu$  3445 (OH), 3346 (NH) 1612 cm<sup>-1</sup> (C=N). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz):  $\delta$ 1.05(s,3H, CH<sub>3</sub>), 3.69 (m, 2H, H-5', H-5''), 4.18 (m, 1H, H-4'), 4.37 (dd, 1H, J=2.8 Hz, J=5.8 Hz, H-3'), 4.44 (t, 1H, J=5.8 Hz, H-2'),4.61 (m, 1H, OH), 4.75 (d, 1H, J=6.3 Hz, OH), 4.90 (m, 1H, OH), 4.98 (t, 1H, J=4.5 Hz, OH), 4.97 (s, 2H, CH<sub>2</sub>), 7.45-7.86(m,4H,Ar-H) 8.10 (d, 1H, H-1'), 11.18 (s, 1H, NH).

#### 2-[2-D-Galactopentitolylidenehydrazinyl)-2-((2-methylbenzimidazol-1-yl)methyl)-1,3,4-oxadiazole (206)

Yield 82%; mp=182-184°C; IR (KBr):  $\nu$  3448 (OH), 3382 (NH), 1612 cm<sup>-1</sup> (C=N). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz):  $\delta$  1.05(s,3H, CH<sub>3</sub>), 3.48 (m, 2H, H-6', H-6''), 3.52 (m, 1H, H-5'), 4.18 (m, 1H, H-4'), 4.36 (dd, 1H, J=2.8 Hz, J=5.8 Hz, H-3'), 4.42 (t, 1H, J=5.8 Hz, H-2'), 4.58 (s, 2H, CH<sub>2</sub>), 4.61 (m, 1H, OH), 4.74 (d, 1H, J=6.3 Hz, OH), 4.91 (m, 1H, OH), 4.97 (t, 1H, J=4.5 Hz, OH), 5.37 (t, 1H, J=4.5 Hz, OH), 7.45-7.86(m,4H,Ar-H), 8.44 (d, 1H, H-1'), 11.14 (s, 1H, NH).

#### General procedure for the synthesis of per-O-acetyl-sugar hydrazinyl derivatives 209,210

To a solution of compounds 205,206 (0.001 mole) in pyridine (7 mL) was added to acetic anhydride (1.02 g, 0.01 mol). The mixture was stirred at room temperature for 5 h. The resulting solution was poured onto crushed ice, and the product that separated out was filtered off, washed with a solution of sodium hydrogen carbonate followed by water and then dried. The products were recrystallized from ethanol.

#### 2-[2-(2,3,4,5-Tetra-O-acetyl-D-xylotetritolylidene)hydrazinyl]-2-[(2-methylbenzimidazol-1-yl)methyl]-1,3,4-oxadiazole (209)

Yield 72%; mp=174-176°C; IR (KBr):  $\nu$  1740 (C=O), 3282 (NH),

1614  $\text{cm}^{-1}$  (C=N).

## 2-[2-(2,3,4,5,6-Penta-O-acetyl-D-galactopentitolylidene)hydrazinyl]-2-[(2-methylbenzimidazol-1-yl)methyl]-1,3,4-oxadiazole (210)

Yield 78%; mp=188-190°C; IR (KBr):  $\nu$  1740 (C=O), 3282 (NH), 1614  $\text{cm}^{-1}$  (C=N).  $^1\text{H}$  NMR (DMSO- $d_6$ , 300 MHz):  $\delta$  1.02(s,1H,  $\text{CH}_3$ ), 1.93, 1.95, 2.03, 2.07, 2.10 (5s, 15H,  $5\times\text{CH}_3\text{CO}$ ), 3.98 (dd, 1H,  $J=11.2$  Hz,  $J=2.8$  Hz, H-6'), 4.07 (dd, 1H,  $J=11.2$  Hz,  $J=3.2$  Hz, H-6''), 4.18 (m, 1H, H-5'), 4.22 (t, 1H,  $J=7.5$  Hz, H-4'), 4.82 (s, 2H,  $\text{CH}_2$ ), 5.18 (dd, 1H,  $J=2.8$  Hz,  $J=6.5$  Hz, H-3'), 5.24 (dd, 1H,  $J=7.5$  Hz,  $J=9.5$  Hz, H-2'), 7.20 (d,1H,  $J=7.8$  Hz, H-1'), 7.45-7.86(m,4H,Ar-H), 11.20 (s, 1H, NH).

### Antimicrobial testing

**Antimicrobial screening:** The agar diffusion method reported by Cruickshank et al. [41] was used for the screening process. The bacteria and fungi were maintained on nutrient agar and Czapek's-Dox agar media, respectively. The assay medium flasks containing 50 ml of nutrient agar for bacteria and Czapek's-Dox agar medium for fungi respectively were allowed to reach 40-50°C to be inoculated with 0.5 ml of the test organism cell suspension. The flasks were mixed well and poured each into a Petri dish (15  $\times$  2 cm) and allowed to solidify. After solidification, holes (0.6 cm diameter) were made in the agar plate by the aid of a sterile cork pooper (diameter 6 mm). The synthesized target compounds were dissolved each in 2 ml DMSO. In these holes, 100  $\mu\text{l}$  of each compound was placed using an automatic micropipette. The Petri dishes were left at 5°C for 1 h to allow diffusion of the samples through the agar medium and retard the growth of the test organism. Plates were incubated at 30°C for 24 h for bacteria and 72 h of incubation at 28°C for fungi. DMSO showed no inhibition zones. The diameters of zone of inhibition were measured and compared with that of the standard and the solvent alone was used as negative control, the values were tabulated. Ciprofloxacin [42,43] (50  $\mu\text{g}/\text{ml}$ ) and Nystatin [44] (50  $\mu\text{g}/\text{ml}$ ) were used as standard for antibacterial and antifungal activity respectively. The observed zones of inhibition are presented in Tables 1 and 2.

The synthesized compounds were screened *in vitro* for their antimicrobial activities against *Staphylococcus aureus* (G+ve bacteria), *Micrococcus* (G+ve bacteria), *Salmonella typhi* (G-ve bacteria), *Salmonella para typhi* (G-ve bacteria) and *Escherichia coli* (G-ve bacteria) in addition to strains of fungi species *Aspergillus flavus*, *Aspergillus fumigates*, *Aspergillus ochraceus* and *Candida albicans* which were isolated from milk and milk products. The diameters of zone of inhibition were measured and compared with that of the standard and the solvent alone was used as negative control. The values of minimal inhibitory concentrations (MICs) of the tested compounds are presented in Tables 1 and 2. The MIC values of the results obtained revealed that compounds showed varying degrees of inhibition against the tested microorganisms. The results indicated generally that tested compounds did not show high activity against fungi under test except **194** that exhibited high antifungal activities against (*Aspergillus fumigates* and *Candida albicans*) and some antifungal activity against *Aspergillus flavus*, in addition to **196** that exhibited some antifungal activities against *Candida albicans*. Also the results indicated that among the compounds tested in antibacterial screening Compound **209** revealed high activities against *Micrococcus* while Compounds **182**, **183**, **184**, **185**, **186** and **187** exhibited good antibacterial activities in addition to Compounds **188**, **189**, **201**, **191** that exhibited appreciable activity against *Micrococcus*. Compounds **172**, **181**, **182**, **183**, **184**, **185**, **186**, **187** and **188** exhibited mild to moderate antibacterial

activity against *Salmonella para typhi*. Compounds **189**, **194**, **196**, **200**, **201**, **202**, **205** and **206** exhibited notable antibacterial activity against *Salmonella typhi*. Compounds **172**, **184**, **188**, **189**, **194**, **209** and **210** exhibited notable antibacterial activity against *Escherichia coli* followed by compounds **183**, **186**, **195** and **206**. Compounds **179**, **181**, **184** and **205** exhibited notable antibacterial activity against *Staphylococcus aureus* followed by compounds **182**, **189**, **201**, **202** and **205**.

### Results and Discussion

2-Methylbenzimidazole (**171**) was allowed to react with ethylchloroacetate in dry acetone and in the presence of anhydrous potassium carbonate to afford 1-ethoxycarbonylmethyl-2-methylbenzimidazole (**172**) in 83% yield. Treatment of **172** with hydrazine hydrate in absolute ethanol at reflux temperature gave the corresponding hydrazide **173** in 83% yield following the reported procedure<sup>146</sup>. The structure of **172** was confirmed by IR spectra which showed the presence of characteristic absorption bands corresponding to ( $\text{CH}_2$ ) groups at  $\nu$  1455  $\text{cm}^{-1}$  and (CO) group at  $\nu$  1730  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR showed a signal corresponding to  $\text{CH}_2\text{CH}_2$  group as a triplet at  $\delta$  1.32 ppm, signal corresponding to  $\text{CH}_2\text{CH}_2$  group as a quartet at  $\delta$  4.20 ppm, signal corresponding to  $\text{NCH}_2$  group as a singlet at  $\delta$  5.08 ppm, and signal for aromatic at  $\delta$  7.12 - 8.14 ppm. Mass spectrum,  $m/z$  ( $I$ , %): 218[ $\text{M}^+$ ] (20). The structure of **173** was confirmed by IR spectra which showed the presence of characteristic absorption bands corresponding to (CO) group at  $\nu$  1637  $\text{cm}^{-1}$ , (NH) group at  $\nu$  3351  $\text{cm}^{-1}$  and ( $\text{NH}_2$ ) group at  $\nu$  3567  $\text{cm}^{-1}$  in addition confirmed by mass spectra revealed the presence characteristic signals corresponding to the molecular ion peaks corresponding to their molecular formulas. 1-(2-Methylbenzimidazol-1-yl) acetic acid hydrazide (Scheme 1). **173** was allowed to react with L-(-)-arabinose, D-(+)-xylose, D-(+)-galactose and D-(+)-mannose in an aq. ethanolic solution with a catalytic amount of acetic acid, to afford the corresponding sugar *N*-acylhydrazones **178-181** in 70-81% yields. The structures of these compounds were confirmed by analytical and spectral data. The IR spectra of **178-181** showed the presence of characteristic absorption bands corresponding to (OH) group in the region  $\nu$  3295-3350  $\text{cm}^{-1}$ . The  $^1\text{H}$  NMR spectra showed signals of the sugar chain protons at  $\delta$  3.24-3.85 ppm, the C-1 methine proton as a doublet in the range  $\delta$  7.09-7.11 ppm, in addition to the aromatic protons in the region  $\delta$  7.12-7.94 ppm. Treatment of the sugar hydrazones **178-181** with acetic anhydride in pyridine at room temperature gave the corresponding per-*O*-acetyl derivatives **182-185** in 60-77% yields. The  $^1\text{H}$  NMR spectra of **182-185** showed the signals of the *O*-acetyl-methyl protons at  $\delta$  1.98-2.48 ppm. The rest of the sugar protons appeared in the range  $\delta$  3.40-5.73 ppm. The reaction of sugar arylhydrazones with boiling acetic anhydride is well known to give either the corresponding per-*O,N*-acetyl derivatives or the respective per-*O,N*-acetyl-1,3,4-oxadiazolin derivatives<sup>147-149</sup>. However, reaction of the sugar hydrazones **178-181** with acetic anhydride at 100°C gave the sugar-substituted 1,3,4-oxadiazoline derivatives **186-189** in 74-81% yields. The IR spectra of **188-192** showed the presence of characteristic absorption bands corresponding to (CO) group in the region  $\nu$  1738-1742  $\text{cm}^{-1}$  and bands corresponding to (C=N) group in the region  $\nu$  1650-1679  $\text{cm}^{-1}$ . The  $^1\text{H}$  NMR spectra of **186-189** showed signals of the *O*- and *N*-acetyl-methyl protons as singlets in the range  $\delta$  1.83-2.48 ppm. The rest of the sugar chain protons appeared in the range  $\delta$  4.05-5.34 ppm and the aromatic protons as multiplets in the region  $\delta$  7.68-7.69 ppm (Scheme 2). When the hydrazide **173** was reacted with carbon disulphide in ethanol in the presence of potassium hydroxide at reflux temperature, it afforded 2-[(2-methylbenzimidazol-1-yl)methyl]-1,3,4-oxadiazole-2-thiol

Compound	Gram-positive	Gram-positive	Gram-negative	Gram-negative	Gram-negative
	<i>Staphylococcus aureus</i>	<i>Micrococcus</i>	<i>Salmonella typhi</i>	<i>Salmonella para typhi</i>	<i>Escherichia coli</i>
171	0	0	0	0	0
172	0	0	0	0	220
178	0	250	130	125	0
179	150	225	0	0	0
180	0	0	0	0	0
181	150	0	125	125	0
182	80	0	100	0	0
183	0	130	weak	100	80
184	125	500	0	225	75
185	0	0	0	0	0
186	0	100	130	125	75
187	0	125	0	0	0
188	0	240	140	0	100
189	75	350	125	225	100
190	75	0	100	110	100
191	0	110	100	125	75
192	0	0	0	0	0
205	110	200	0	0	75
206	0	0	0	0	0
207	0	0	0	0	0
208	0	0	0	0	0
209	0	0	0	0	0
210	0	0	0	0	0
211	0	0	0	0	0
212	75	350	0	0	125
213	0	0	0	0	0
214	75	250	0	0	125
215	0	0	0	0	0
216	0	0	0	0	0
217	0	0	0	0	0
218	125	225	0	225	130
219	0	0	0	0	0
220	0	100	0	140	0
221	0	100	0	0	75
222	0	0	0	0	0
223	0	125	0	0	75
Ciprofloxacin	125	110	220	225	225
DMSO	0	0	0	0	0




Table 1: Minimum inhibitory concentrations (MIC in µg/ml) of the title Compounds against bacteria species. The negative control DMSO showed no activity.

Compound	Fungi	Fungi	Fungi	Fungi
	<i>Aspergillus flavus</i>	<i>Aspergillus fumigates</i>	<i>Aspergillus ochraceus</i>	<i>Candida albicans</i>
171	0	0	0	0
172	0	0	0	0
179	0	0	0	0
181	0	0	0	0
184	0	0	0	0
187	0	0	0	0
189	0	0	0	0
190	0	0	0	0
191	0	0	0	0
205	0	0	0	0
209	0	0	0	0
210	0	0	0	0
211	0	0	0	0
212	120	500	0	500
214	0	0	0	0
216	0	0	0	0

218	0	0	0	0
221	0	0	0	110
222	0	0	0	0
223	0	0	0	0
Nystatin DMSO	250	150	45	225
	0	0	0	0



**Table 2:** Minimum inhibitory concentrations (MIC in  $\mu\text{g/ml}$ ) of the title Compounds against fungi species. The negative control DMSO showed no activity.

(194) in 80% yield. The  $^1\text{H}$  NMR spectrum of the 1,3,4-oxadiazole-2-thione showed a signal corresponding to the  $\text{CH}_2$  group as a singlet at  $\delta$  4.58 ppm, Ar-H signals as multiplet at  $\delta$  7.97 - 8.48 ppm in addition to the SH signal at  $\delta$  12.80 ppm. Alkylation of the 1,3,4-oxadiazolethione **194** with methyl iodide in alkaline medium afforded the corresponding S-methyl derivative **195** in 71% yield. Hydrazinolysis of **195** gave 2-hydrazinyl-2-[(2-methylbenzimidazol-1-yl)methyl]-1,3,4-oxadiazole (**196**) in 74% yield. The  $^1\text{H}$  NMR spectra of **195** showed the signals of the methyl group as a singlet which disappeared in the  $^1\text{H}$  NMR spectrum of **196** in which the  $\text{NH}_2$  signal appeared at  $\delta$  5.80 ppm (Scheme 3). 2-Hydrazinyl-2-[(2-methylbenzimidazol-1-yl)methyl]-1,3,4-oxadiazole (**196**) was refluxed with various aromatic aldehydes in ethanol and in the presence of a catalytic amount of acetic acid to afford the corresponding Schiff's bases **200** - **202** in 89-95 yields. The structures of **201** and **202** were confirmed by mass spectra which revealed the presence of the characteristic signals corresponding to their molecular formulas. The structures of **200**, **201**, and **202** were confirmed by  $^1\text{H}$  NMR spectra which showed a signal corresponding to the  $\text{CH}_2$  group as a singlet at  $\delta$  4.51-4.65 ppm, signals of the aromatic protons at  $\delta$  7.22-8.64 ppm, a



signal corresponding to the CH group (H-4-furan and H-3-furan) as a singlet at  $\delta$  6.24 and 6.79 ppm respectively, NH signal at  $\delta$  9.93-9.96 ppm, in addition to the CH=N signals as tow singlet at  $\delta$  7.80,8.8.00 and 8.22 ppm (Scheme 4). **196** was allowed to react with D-(+)-xylose and D-(+)- galactose in an aq. ethanolic solution with a catalytic amount of acetic acid. The structures of these compounds were confirmed by spectral data. The IR spectra of **205**, **206** showed the presence of characteristic absorption bands corresponding to (OH) group in the region  $\nu$  3445–3448  $\text{cm}^{-1}$ .

The  $^1\text{H}$  NMR spectra showed signals of the sugar chain protons at  $\delta$  3.48-4.44 ppm, the C-1 methine proton as a doublet in the range  $\delta$  8.10-8.44 ppm, in addition to the aromatic protons in the region  $\delta$  7.45-7.86 ppm. Treatment of the sugar hydrazones **205**, **206** with acetic anhydride in pyridine at room temperature gave the corresponding per-O-acetyl derivatives **209**, **210** in 72, 78% yields. The  $^1\text{H}$  NMR spectra of **209**, **210** showed the signals of the O-acetyl-methyl protons at  $\delta$  1.93-2.10 ppm. The rest of the sugar protons appeared in the range  $\delta$  3.98-5.24 ppm (Scheme 5).



## References

- Ohta S, Osaki T, Nishio S, Furusawa A, Yamashunta M, et al. (2000) Serial double nucleophilic addition of amines to the imidazole nucleus. *Tetrahedron Lett* 41: 7503-7506.
- Kubicki M (2004) Two tautomers in one crystal: 4(5)-nitro-5(4)-methoxyimidazole. *Acta Crystallogr B* 60: 191-196.
- Maeki J, Taehinen P, Kronberg L, Klika KD (2005) *J Phys Org Chem* 18: 240.
- Zhong YL, Lee J, Reamer RA, Askin D (2004) New method for the synthesis of diversely functionalized imidazoles from N-acylated alpha-aminonitriles. *Org Lett* 6: 929-931.
- Bossio R, Marcaccini S, Pepino R, Torroba T (1996) Studies on Isocyanides and Related Compounds. Synthesis of 1-Aryl-2-(tosylamino)-1H-imidazoles, a Novel Class of Imidazole Derivatives. *J Org Chem* 61: 2202-2203.
- Markwalder JA, Pottorf RS, Seitz SP (1997) One-pot conversion of alpha-ureido and alpha-thioureido esters to imidazol-2-ones and imidazole-2-thiones. *Synlett* 521-522.
- Reader VA (1998) An efficient synthesis of 2-(methylaminomethyl)-4,5-dialkyl-1H imidazoles. *Synlett* 1077-1078.
- Munk SA, Harcourt DA, Arasasingham PN, Burke JA, Kharlamb AB, et al. (1997) Synthesis and evaluation of 2-(arylamino)imidazoles as alpha 2-adrenergic agonists. *J Med Chem* 40: 18-23.
- Khanna I, Yu Y, Huff R, Weier R, Xu X, et al. (2000) Selective cyclooxygenase-2 inhibitors: heteroaryl modified 1,2-diarylimidazoles are potent, orally active antiinflammatory agents. *J Med Chem* 43: 3168-3185.
- Habersaat N, Froehlich R, Wuerthwein E (2004) 4H-Imidazoles and Imidazoles from anionic and dipolar electrocyclization reactions of 2,4-diazapenta-1,3-dienes. *Eur J Org Chem* 2567-2581.
- Yamada M, Yura T, Morimoto M, Harada T, Yamada K, et al. (1996) 2-[(2-Aminobenzyl)sulfinyl]-1-(2-pyridyl)-1,4,5,6-tetrahydrocyclopent[d]imidazoles as a novel class of gastric H<sup>+</sup>/K<sup>+</sup>-ATPase inhibitors. *J Med Chem* 39: 596-604.
- Marwaha A, Singh P, Mahajan MP, Velumurugan D (2004) An efficient unprecedented synthesis of novel functionalized imidazoles from secondary amino-N-carbothioic acid (phenyl-p-tolylimino-methyl)amides and dimethyl acetylenedicarboxylate. *Tetrahedron Lett* 45: 8945-8947.
- Derstine CW, Smith DN, Katzenellenbogen JA (1996) Trifluoromethyl-Substituted Imidazolines: Novel Precursors of Trifluoromethyl Ketones Amenable to Peptide Synthesis. *J Am Chem Soc* 118: 8485-8486.
- Derstine CW, Smith DN, Katzenellenbogen JA (1997) Trifluoromethyl-Substituted  $\Delta^3$ -Imidazolines: Synthesis and Reactivity. *Tetrahedron Lett* 38: 4359-4362.
- Sparks RB, Combs AP (2004) Microwave-assisted synthesis of 2,4,5-triaryl-imidazole; a novel thermally induced N-hydroxyimidazole N-O bond cleavage. *Org Lett* 6: 2473-2475.
- Siamaki AR, Arndtsen BA (2006) A direct, one step synthesis of imidazoles from imines and acid chlorides: a palladium catalyzed multicomponent coupling approach. *J Am Chem Soc* 128: 6050-6051.
- Bon RS, Hong C, Bouma MJ, Schmitz RF, de Kanter FJ, et al. (2003) Novel multicomponent reaction for the combinatorial synthesis of 2-imidazolines. *Org Lett* 5: 3759-3762.
- Bon RS, van Vliet B, Sprenkels NE, Schmitz RF, de Kanter FJ, et al. (2005) Multicomponent synthesis of 2-imidazolines. *J Org Chem* 70: 3542-3553.
- Pridgen LN, Mokhallati MK, McGuire MA (1997) A Stereospecific Synthesis of Both Enantiomers of 2-(1'-Amino-2'-Methylpropyl) Imidazole, a Key Synthon in the Synthesis of SB 203386; a Potent Protease Inhibitor. *Tetrahedron Lett* 38: 1275-1278.
- Le Bihan G, Rondou F, Pelé-Tounian A, Wang X, Lidy S, et al. (1999) Design and synthesis of imidazoline derivatives active on glucose homeostasis in a rat model of type II diabetes. 2. Syntheses and biological activities of 1,4-dialkyl-, 1,4-dibenzyl-, and 1-benzyl-4-alkyl-2-(4',5'-dihydro-1'H-imidazol-2'-yl)piperazines and isosteric analogues of imidazoline. *J Med Chem* 42: 1587-1603.
- McKenna JM, Halley F, Souness JE, McLay IM, Pickett SD, et al. (2002) An algorithm-directed two-component library synthesized via solid-phase

- methodology yielding potent and orally bioavailable p38 MAP kinase inhibitors. *J Med Chem* 45: 2173-2184.
22. Bilodeau MT, Cunningham AM (1998) PEG-Supported Synthesis of 3,5-Disubstituted-1,2,4-Triazoles. *J Org Chem* 63: 2800.
23. Peddibhotla S, Jayakumar S, Tepe JJ (2002) Highly diastereoselective multicomponent synthesis of unsymmetrical imidazolines. *Org Lett* 4: 3533-3535.
24. Kitbunnadaj R, Hashimoto T, Poli E, Zuiderveld OP, Menozzi A, et al. (2005) N-substituted piperidinyll alkyl imidazoles: discovery of methimepip as a potent and selective histamine H3 receptor agonist. *J Med Chem* 48: 2100-2107.
25. Kitbunnadaj R, Zuiderveld OP, De Esch IJ, Vollinga RC, Bakker R, et al. (2003) Synthesis and structure-activity relationships of conformationally constrained histamine H(3) receptor agonists. *J Med Chem* 46: 5445-5457.
26. De Esch IJ, Vollinga RC, Goubitz K, Schenk H, Appelberg U, et al. (1999) Characterization of the binding site of the histamine H3 receptor. 1. Various approaches to the synthesis of 2-(1H-imidazol-4-yl)cyclopropylamine and histaminergic activity of (1R,2R)- and (1S,2S)-2-(1H-imidazol-4-yl)-cyclopropylamine. *J Med Chem* 42: 1115-1122.
27. Neipp CE, Ranslow PB, Wan Z, Snyder JK (1997) Dienophilicity of Imidazole in Inverse Electron Demand Diels-Alder Reactions; Intramolecular Reactions with 1,2,4-Triazines. *Tetrahedron Lett* 38: 7499-7502.
28. Li HY, Drummond S, DeLucca I, Boswell GA (1996) Singlet oxygen oxidation of pyrroles - synthesis and chemical-transformations of novel 4,4-bis(trifluoromethyl)imidazoline analogs. *Tetrahedron* 52: 11153-11162.
29. Hrabie JA, Citro ML, Chmurny GN, Keefer LK (2005) Carbon-bound diazeniumdiolates from the reaction of nitric oxide with amidines. *J Org Chem* 70: 7647-7653.
30. Larter ML, Phillips M, Ortega F, Aguirre G, Somanathan R, et al. (1998) Synthesis of racemic cis and trans 2,4,5-triprydylimidazolines. *Tetrahedron Lett* 39: 4785-4788.
31. Mohammadpoor-Baltork I, Zolfigol MA, Abdollahi-Alibeik M (2004) Novel, mild and chemoselective dehydrogenation of 2-imidazolines with trichloroisocyanuric acid. *Synlett* 2803-2805.
32. Mohammadpoor-Baltork I, Zolfigol MA, Abdollahi-Alibeik M (2004) Novel and chemoselective dehydrogenation of 2-substituted imidazolines with potassium permanganate supported on silica gel. *Tetrahedron Lett* 45: 8687-8690.
33. Abdollahi-Alibeik M, Mohammadpoor-Baltork I, Zolfigol MA (2004) Alumina supported potassium permanganate: an efficient reagent for chemoselective dehydrogenation of 2-imidazolines under mild conditions. *Bioorg Med Chem Lett* 14: 6079-6082.
34. Mohammadpoor-Baltork I, Abdollahi-Alibeik M (2005) Mild, efficient, and chemoselective dehydrogenation of 2-imidazolines, bis-imidazolines, and N-substituted-2-imidazolines with potassium permanganate supported on montmorillonite K-10. *Can J Chem* 83: 110-114.
35. Anastassiadou M, Baziard-Mouysset G, Payard M (2000) New one-step synthesis of 2-aryl-1H-imidazoles: Dehydrogenation of 2-aryl-Delta 2-imidazolines with dimethylsulfoxide. *Synthesis* 1814-1816.
36. Giorgio E, Minichino C, Viglione RG, Zanasi R, Rosini C, et al. (2003) Assignment of the molecular absolute configuration through the ab initio Hartree-Fock calculation of the optical rotation: can the circular dichroism data help in reducing basis set requirements? *J Org Chem* 68: 5186-5192.
37. Lovely CJ, Du H, Dias HVR (2003) The Regioselective Preparation of 1-Benzyl- and 1-Methyl-4-vinylimidazole and their reactions with N-Phenylmaleimide. *Heterocycles* 60: 1-7.
38. Muri E, Mishra H, Avery M, Williamson J (2003) Design and Synthesis of Heterocyclic Hydroxamic Acid Derivatives as Inhibitors of Helicobacter pylori Urease. *Synth Commun* 33: 1977-1995.
39. Hayakawa Y, Kimoto H, Cohen L, Kirk K (1998) Syntheses of (Trifluoromethyl) imidazoles with Additional Electronegative Substituents. An Approach to Receptor-Activated Affinity Labels. *J Org Chem* 63: 9448-9454.
40. Shafiee A, Rastkary N, Foroumadi A (1998) Syntheses of 2-(2-arylethyl) imidazoles. *J Heterocycl Chem* 35: 607-610.
41. Cruickshank R, Duguid JP, Marion BP, Swain RHA (1975) *Medicinal Microbiology*. 12th edn. Churchill Livingstone, London. p: 196.
42. Dahiya R (2008) Synthesis, characterization and antimicrobial studies on some newer imidazole analogs. *Sci Pharm* 76: 217-239.
43. Su HC, Ramkissoon K, Doolittle J, Clark M, Khatun J, et al. (2010) The development of ciprofloxacin resistance in Pseudomonas aeruginosa involves multiple response stages and multiple proteins. *Antimicrob Agents Chemother* 54: 4626-4635.
44. Poyner TF, Dass BK (1996) Comparative efficacy and tolerability of fusidic acid/hydrocortisone cream (Fucidin H cream) and miconazole/hydrocortisone cream (Daktacort cream) in infected eczema. *J Eur Acad Dermatol Venereol* 7: S23-S30.

**Citation:** Amer HH, El-Kousy SM, Salama WM, H-H Sheleby A (2016) Synthesis and Antimicrobial Activity of New Synthesized Benzimidazole Derivatives and their Acyclic Nucleoside Analogues. *Organic Chem Curr Res* 5: 157. doi:10.4172/2161-0401.1000157

### OMICS International: Publication Benefits & Features

#### Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

#### Special features:

- 700 Open Access Journals
- 50,000 Editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus, Google Scholar etc.
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsgroup.org/journals/submission>