

# Synthesis, DNA-binding Properties and Quantitative Structure-Activity Relationships on Ruthenium(II) Complexes with Calf-Thymus DNA

Hao Zhang<sup>1</sup>, Fan Yang<sup>1</sup>, Qiong Wu<sup>2</sup>, Si-Yan Liao<sup>3</sup>, Ping Liu<sup>4</sup>, Wen-Jie Mei<sup>2\*</sup>, Li Li<sup>3</sup>, Shuang-Yan Zhang<sup>3</sup> and Xi-Cheng Wang<sup>1\*</sup>

<sup>1</sup>The First Affiliated Hospital of Guangdong Pharmaceutical University, Guangzhou, 510006, PR China

<sup>2</sup>The Pharmacy Department of Guangdong Pharmaceutical University, Guangzhou, 510224, PR China

<sup>3</sup>School of Pharmacy, Guangzhou Medicine University, Guangzhou 510275, PR China

<sup>4</sup>The Chemistry Department of Hu'nan Arts and Philosophy Science University, Changde, 415000, PR China

## Abstract

A series of novel ruthenium(II) complexes with electron-donor or electron-acceptor groups in intercalative ligands,  $[\text{Ru}(\text{phen})_2(o\text{-MOPIP})]^{2+}$  (1),  $[\text{Ru}(\text{phen})_2(o\text{-MPIP})]^{2+}$  (2),  $[\text{Ru}(\text{phen})_2(o\text{-CPIP})]^{2+}$  (3) and  $[\text{Ru}(\text{phen})_2(o\text{-NPIP})]^{2+}$  (4) have been synthesized and characterized with elementary, ES-MS, <sup>1</sup>H NMR, electronic absorption and emission spectra. The binding properties of these complexes to CT-DNA have been investigated by spectroscopy and viscosity experiments. It's illustrated that these complexes bind to DNA in a non-classical intercalation mode and their intrinsic binding constants ( $K_b$ ) for 1, 2, 3 and 4 are calculated as 1.1, 0.35, 0.53 and  $1.7 \times 10^5 \text{ M}^{-1}$ , respectively. The Quantitative Structure-Activity Relationships (QSAR) of these ruthenium complexes, as well as some other ruthenium complexes congers has been investigated, and a linearity equation have been obtained:  $\log K_b = 0.2429\pi + 0.0429\pi^2 + 0.2907\sigma + 0.63891 + 4.3491$  ( $n=12$ ;  $R=0.9338$ ;  $F=11.9134$ ;  $p=0.0030$ ). This results show that the electron-acceptor group and a large hydrophobic group will enhance the DNA binding affinity of ruthenium complexes.

**Keywords:** Ruthenium; DNA; Quantitative structure-Activity relationships

**Abbreviations:** Phen: 1,10-phenanthroline; *o*-MOP: *o*-(2-methoxyphenyl) imidazo [4,5-*f*][1,10] phenanthroline; *o*-MP: *o*-(2-methylphenyl) imidazo [4,5-*f*][1,10] phenanthroline; *o*-CP: *o*-(2-chlorophenyl) imidazo [4,5-*f*][1,10] phenanthroline; *o*-NP: 2-(2-nitrophenyl) imidazo [4,5-*f*][1,10] phenanthroline

## Introduction

For years, many attentions have been focused on the interaction of octahedral Ru (II) complexes with DNA owing to their potential utility as DNA probes, molecular light switches and chemotherapy drugs and photodynamic therapy for tumors [1-13]. For one thing, DNA has long been considered the main target for anticancer drugs. In general, Ru (II) complexes can bind to DNA in three non-covalent modes: intercalation binding, groove binding and electrostatic binding. It's know that complexes with an enlarged aromatic ligand (intercalating ligand) can bind to DNA with high affinity (104~106), while those complex such as  $\text{Ru}(\text{bpy})_3^{2+}$  can bind to DNA mostly in electrostatic [14].

For the last decade, a number of ruthenium complex with 2-phenylimidazo [4,5-*f*][1,10]-phenanthroline (PIP) and its derivates as intercalating ligand have been synthesized and their DNA binding properties have been investigated thoroughly. Ji et al. indicate that the binding affinity of these complexes depended not only on the conformations of DNA, but also on the structures of intercalative ligands [15-18]. The factors included the enlarged aromatic ring, intramolecular hydrogen bond and the planarity properties of intercalative ligand will enhance the binding affinity of these ruthenium to DNA, and the electronic effects (the donor/acceptor electron properties of substituent group on intercalative ligand) is also one factor influencing the bind of complexes to DNA [19,20].

A recently studies try to explain the DNA-binding affinity by computational calculations with density functional theory (DFT). It's shown that the energy of these complex' frontier molecular orbit, that is the highest occupied molecular orbit (HOMO) and the lowest unoccupied molecular orbit (LUMO) is varied and when they intercalating in the DNA base pairs, the energy of the transition

conformation will different. According to the frontier molecular theory, an electron will transfer more easily from a high HOMO to a lower LUMO, and resulting those complex have the lowest LUMO (in generally, the HOMO of DNA is higher than that of ruthenium complexes) will bind to DNA the strongest [21-23]. But this is still confused since the binding energy of complexes to DNA is not known and the optimal conformation of the supermolecular complex-DNA is also not been illuminated.

In 1930's, Hammett indicated that the activity of an organic reaction is in relating to the substituent effects [24-26]. Based on these, Fujita and Hansch developed a linear Hansch equation to elucidate the relationship between the bioactivity/physical activity and the structure of organic molecules [27], and which is so called the linear free energy relationships and thus have been utilized extensively in agrochemistry, pharmaceutical chemistry, toxicology [28] for its excellent predictable ability. More recently, Prasanna S et al. successful discerned the structural and physicochemical requirements for selective COX-2 over COX-1 inhibition among the fused pyrazole ring systems by Hansch method [29]. However, there are still no reports focused on quantity structure-activity relationship on DNA-binding properties of ruthenium(II) complexes [30,31].

In this paper, a series of Ru (II) complexes with electron-donor or electron- acceptor substituents in the intercalative ligands,

**\*Corresponding authors:** Xi-Cheng Wang, The First Affiliated Hospital of Guangdong Pharmaceutical University, Guangzhou, 510006, PR China, Tel: 86-20-61321397; E-mail: [13902400598@126.com](mailto:13902400598@126.com)

Wen-Jie Mei, The Pharmacy Department of Guangdong Pharmaceutical University, Guangzhou, 510224, PR China, Tel: 86-20-39352129; E-mail: [wenjimei@126.com](mailto:wenjimei@126.com)

**Received** February 25, 2016; **Accepted** March 08, 2016; **Published** March 10, 2016

**Citation:** Zhang H, Yang F, Wu Q, Si-Yan L, Liu P, et al. (2016) Synthesis, DNA-binding Properties and Quantitative Structure-Activity Relationships on Ruthenium(II) Complexes with Calf-Thymus DNA. Med chem (Los Angeles) 6: 137-142. doi:[10.4172/2161-0444.1000338](https://doi.org/10.4172/2161-0444.1000338)

**Copyright:** © 2016 Zhang H, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

[Ru(phen)2(*o*-MOP)]<sup>2+</sup>1, [Ru(phen)2(*o*-MP)]<sup>2+</sup>2, [Ru(phen)2(*o*-CP)]<sup>2+</sup>3 and [Ru(phen)2(*o*-NP)]<sup>2+</sup>4 (Scheme 1) were synthesized and characterized. The DNA-binding properties of these complexes have been investigated by the spectroscopic and viscosity experiments. The quantity structural-activity relationship of these ruthenium complexes, as well as some other analogues has also been investigated.

## Experiment Section

### Chemicals


CT-DNA was purchased from the Sino-American Biotechnology Company. All reagents and solvents were purchased commercially (AR, Acros Inc., and Sigma Inc., etc.) and used without further purification unless otherwise noted. Doubly distilled water was used to prepare buffers. The concentration of calf thymus DNA was determined spectrophotometrically using the molar absorptivity 6600 M<sup>-1</sup>.cm<sup>-1</sup> (260 nm) (The ratio of UV absorbance at 260 and 280 nm is in the range of 1.8-1.9:1).

### Synthesis and characterization

[Ru(phen)<sub>2</sub>Cl<sub>2</sub>].2H<sub>2</sub>O were prepared following the literature procedure [24]. Ru(II) complexes 1, 2, 3 and 4 were synthesized by refluxing Ru(phen)<sub>2</sub>Cl<sub>2</sub> and *o*-MOP (*o*-MP, *o*-CP or *o*-NP) in ethylene glycol under an argon atmosphere with high yield. Each complex was obtained as a PF<sub>6</sub><sup>-</sup> salt and purified with column chromatography.

Electrospray mass spectrometry (ES MS) has recently been shown to be a powerful tool for measuring the molecular mass of non-volatile and thermally unstable compounds [26]. The ES-MS for 1, 2, 3 and 4 exhibits a fragment ion peak of (M+1PF<sub>6</sub>)<sup>+</sup> at 933.0, 917.0, 936.9 and 947.9 (*m/z*), respectively (Table 1). The fragment ion peaks of M<sup>2+</sup> of these complexes appear at 394.3, 386.4, 396.4 and 401.8 for 1, 2, 3 and 4, respectively, and the resolution of this peak for these complexes shows that the species is doubly charged and the isotopic distribution corresponds to the calculated one.

The electronic absorption spectra of these Ru (II) complexes in Tris-buffer are characterized by an intense ligand-centered transition in the UV region and a metal-to-ligand charge transfer transition (MLCT) in the visible region. The lowest-energy absorption bands ascribed to the MLCT transitions for 1, 2, 3 and 4 are 455, 452, 453 and 453 nm respectively. The intense and sharp bands at 263, 263, 263 and 261 nm in UV region for 1, 2, 3 and 4, respectively, are attributed to the intraligand π→π\* transition via comparison with the spectra of [Ru(bpy)<sub>3</sub>]<sup>2+</sup>. Little variation in the energy of the MLCT bands with the electronic effect in intercalative ligand was observed.



No.	Substituent Group			Physical Parameter			logK <sub>b</sub>
	X	Y	Z	σ <sup>a</sup>	π <sup>b</sup>	I <sup>c</sup>	
1	-OCH <sub>3</sub>	-H	-H	0.04	-0.02	1	5.04
2	-CH <sub>3</sub>	-H	-H	-0.15	0.56	0	4.54
3	-Cl	-H	-H	0.4	0.71	0	4.72
4	-NO <sub>2</sub>	-H	-H	1.05	-0.28	1	5.23
5	-Br	-H	-H	0.44	0.86	0	4.57 <sup>d</sup>
6	-H	-OCH <sub>3</sub>	-H	-0.27	-0.02	1	4.84
7	-H	-Cl	-H	0.23	0.71	0	4.49
8	-H	-NO <sub>2</sub>	-H	0.78	-0.28	1	5.19
9	-H	-OH	-H	-0.37	-0.67	1	4.83
10	-H	-Br	-H	0.23	0.86	0	4.77 <sup>d</sup>
11	-H	-OC <sub>6</sub> H <sub>5</sub>	-H	-0.03	2.08	0	5.04 <sup>e</sup>
12	-H	-OH	-OCH <sub>3</sub>	-0.33	-0.69	1	4.63 <sup>f</sup>

a. σ is the abbreviation of electron parameter. All data are cited from reference [27].

b. π is the abbreviation of hydrophobic parameter. All data are cited from reference [27].

c. I is the abbreviation of an inductive parameter. The data is 1 when there is a hydrogen bond in the intercalative ligand, or it will be 0.

d. These data is cited from reference [39].

e. These data is cited from reference [40].

f. These data is cited from reference [41].

g. These data is cited from reference [31].

**Table 1:** The intrinsic binding constant and the physical parameters of Ruthenium(II) complex.

Ru (II) complexes 1, 2 and 3 emit fluorescence in Tris-buffer in the range of 500 – 700 nm at room temperature, with the maximum at 589, 588 and 589, respectively, and only a very weak fluorescence was observed for complexes 4 at the same conditions (the maximum is at 588 nm).

**2-(2-methoxyphenyl) imidazo [4,5-*f*][1,10] phenanthroline(*o*-MOP) (1a):** The ligand 2-(2-methoxyphenyl) imidazo [4,5-*f*][1,10] phenanthroline (*o*-MOP) was prepared by the method similar to that in reference [32], and with some modification.

A solution of phenanthraquinone (0.26 g, 1.2 mmol), *o*-anisaldehyde (0.24 g, 1.8 mmol) and ammonium acetate (1.9 g, 25 mmol) in 10 cm<sup>3</sup> glacial acetic acid was refluxed for 2 hour. The cooled deep red solution was diluted with 25 cm<sup>3</sup> water, and neutralized with ammonium hydroxide. Then the mixture was filtered and the precipitates were washed with water and acetone, then dried and purified by chromatography over 60-80 mesh SiO<sub>2</sub> using methanol as an eluent, yields: 0.35 g, 84%. Calculated for C<sub>20</sub>H<sub>14</sub>N<sub>4</sub>O·H<sub>2</sub>O (%): C: 69.7; H: 4.69; N: 16.3; Found(%): C: 69.3; H: 4.66; N: 16.2. ES-MS (in DMSO, *m/z*): 326.7 (calc. 326.4).

**2-(2-methylphenyl) imidazo [4,5-*f*][1,10] phenanthroline(*o*-MP) (2a):** *O*-MP was synthesized by the same method as above, but with phenanthraquinone (0.26 g, 1.2 mmol) and *o*-tolualdehyde (0.22 g, 1.8 mmole), yield: 0.31 g, 78%. Calculated for C<sub>20</sub>H<sub>14</sub>N<sub>4</sub>·H<sub>2</sub>O(%): C: 73.2; H: 4.91; N: 17.1; Found(%): C: 73.6; H: 4.97; N: 17.4. ES-MS (in DMSO, *m/z*): 310.6 (calc. 310.4).

**2-(2-chlorophenyl) imidazo [4,5-*f*][1,10] phenanthroline(*o*-CP) (3a):** *O*-CP was synthesized by the same method as above, but with phenanthraquinone (0.26 g, 1.2 mmol) and 2-chlorobenzaldehyde (0.25 g, 1.8 mmole), yield: 0.32 g, 76%. Calculated for C<sub>19</sub>ClH<sub>11</sub>N<sub>4</sub>·H<sub>2</sub>O(%): C: 65.4; H: 3.76; N: 16.1; Found(%): C: 65.6; H: 3.82; N: 16.4. ES-MS (in DMSO, *m/z*): 331.0 (calc. 330.8).

**2-(2-nitrophenyl) imidazo [4,5-*f*][1,10] phenanthroline(*o*-NP) (4a):** *O*-NP was synthesized by the same method as above, but with phenanthraquinone (0.26 g, 1.2 mmol) and 2-nitrobenzaldehyde (0.27

g, 1.8 mole), yield: 0.38 g, 87%. Calculated for  $C_{19}H_{11}N_5O_2 \cdot H_2O$ (%): C: 63.5; H: 3.65 N: 19.5; Found(%): C: 63.9; H: 3.62 N: 19.1. ES-MS (in DMSO,  $m/z$ ): 341.6 (calc. 341.3).

**[Ru(phen)<sub>2</sub>(o-MOP)]<sup>2+</sup>(1):** [Ru(phen)<sub>2</sub>(o-MOP)]<sup>2+</sup> was synthesized by the literature [31,33] and with some modification. [Ru(phen)<sub>2</sub>Cl<sub>2</sub>·2H<sub>2</sub>O (0.09 g, 0.17 mmol) and 1a (0.058 g, 0.17 mmol) were added to 10 cm<sup>3</sup> ethylene glycol. The mixture was refluxed for 2 h under an argon atmosphere. The cooled reaction mixture was diluted with water (20 cm<sup>3</sup>) and filtered to remove solid impurities. The complex was then separated from soluble impurities by precipitation with NH<sub>4</sub>PF<sub>6</sub>. The precipitated complex was dried, dissolved in a small amount of acetonitrile, and purified by chromatography over alumina oxide using MeCN-toluene (2:1, v/v) as an eluent, yield: 0.16 g, 84%. Calculated for C<sub>44</sub>F<sub>12</sub>H<sub>30</sub>N<sub>8</sub>OP<sub>2</sub>Ru·2H<sub>2</sub>O(%): C: 47.4; H: 3.08; N: 10.1; Found(%): C: 47.7; H: 3.11; N: 10.4; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, δ ppm): 9.32 (1H, d); 9.08 (1H, d); 8.77 (4H, d); 8.39 (4H, s); 8.21 (2H, d); 8.14 (2H, t); 8.12 (2H, d); 8.02 (2H, 2d); 7.81 (6H, m); 7.74 (1H, t); 7.35 (1H, d); 7.24 (1H, t); 4.04 (3H, s); ES MS of the PF<sub>6</sub><sup>-</sup> salt in MeCN:  $m/z$  933.0 (M+1PF<sub>6</sub>)<sup>+</sup> (calc: 932.8); 394.3 (M)<sup>2+</sup> (calc: 393.9). Absorption UV-Vis, in water at pH 7.2  $\lambda_{max}(\epsilon/10^4 M^{-1}cm^{-1})$ : 263(8.6), 455(1.7). No corrected emission maximum in water at pH 7.2: 589.4 nm.

**[Ru(phen)<sub>2</sub>(o-MP)]<sup>2+</sup>(2):** [Ru(phen)<sub>2</sub>(o-MP)]<sup>2+</sup> was prepared by the above-mentioned method but with 2a (0.056 g, 0.17 mmol); yield: 0.14 g, 77%. Calculated for C<sub>44</sub>F<sub>12</sub>H<sub>30</sub>N<sub>8</sub>P<sub>2</sub>Ru·2H<sub>2</sub>O(%): C: 48.1; H: 3.12; N: 10.2; Found (%): C: 48.4; H: 3.12; N: 10.6; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, δ ppm): 9.06 (2H, d); 8.78 (4H, d); 8.39 (4H, s); 8.17 (2H, d); 8.08 (2H, d); 8.04 (2H, d); 7.85 (6H, m); 7.51 (3H, m); 7.26 (1H, t); 2.70 (3H, s); ES MS of the PF<sub>6</sub><sup>-</sup> salt in MeCN:  $m/z$  917.0 (M+1PF<sub>6</sub>)<sup>+</sup> (calc: 916.8); 386.4 (M)<sup>2+</sup> (calc: 385.9). Absorption UV-Vis, in water at pH 7.2  $\lambda_{max}(\epsilon/10^4 M^{-1}cm^{-1})$ : 264(9.9), 452(1.7). No corrected emission maximum in water at pH 7.2: 587.8 nm.

**[Ru(phen)<sub>2</sub>(o-CP)]<sup>2+</sup>(3):** [Ru(phen)<sub>2</sub>(o-CP)]<sup>2+</sup> was prepared by the above-mentioned method but with 3a (0.059 g; 0.17 mmol); yield: 0.15 g, 79%. Calculated for C<sub>43</sub>F<sub>12</sub>ClH<sub>27</sub>N<sub>8</sub>P<sub>2</sub>Ru·4H<sub>2</sub>O(%): C: 44.8; H: 3.06; N: 9.71; Found (%): C: 44.5; H: 3.10; N: 9.81; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, δ ppm): 9.03 (2H, d); 8.77 (4H, d); 8.38 (4H, s); 8.13 (2H, d); 8.07 (2H, d); 8.00 (2H, d); 7.93 (1H, d); 7.77 (6H, m); 7.64 (2H, m); 7.26 (1H, m); ES MS of the PF<sub>6</sub><sup>-</sup> salt in MeCN:  $m/z$  936.9 (M+1PF<sub>6</sub>)<sup>+</sup> (calc: 937.2); 396.4 (M)<sup>2+</sup> (calc: 396.2). Absorption UV-Vis, in water at pH 7.2  $\lambda_{max}(\epsilon/10^4 M^{-1}cm^{-1})$ : 263(11.6) 454(2.1). No corrected emission maximum in water at pH 7.2: 589.4 nm.

**[Ru(phen)<sub>2</sub>(o-NP)]<sup>2+</sup>(4):** [Ru(phen)<sub>2</sub>(o-NP)]<sup>2+</sup> was prepared by the above-mentioned method but with 4a (0.061 g; 0.17 mmol); yield: 0.14 g, 74%. Calculated for C<sub>43</sub>F<sub>12</sub>H<sub>27</sub>N<sub>9</sub>O<sub>2</sub>P<sub>2</sub>Ru·2H<sub>2</sub>O (%) : C: 45.8; H: 2.77; N: 11.2; Found (%): C: 45.5; H: 2.81; N: 11.8; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, δ ppm): 8.87 (2H, d); 8.77 (4H, d); 8.38 (4H, s); 8.21 (1H, d); 8.12 (2H, d); 8.10 (2H, d); 7.85 (2H, d); 7.76 (6H, m); 7.68 (3H, m); ES MS of the PF<sub>6</sub><sup>-</sup> salt in MeCN:  $m/z$  947.9 (M+1PF<sub>6</sub>)<sup>+</sup> (calc: 947.8); 802.3 (M-H<sup>+</sup>)<sup>+</sup> (calc: 801.8); 401.8 (M)<sup>2+</sup> (calc: 401.4). Resolution of the peak 401.8 shows that the species is double charged and the isotopic distribution corresponds to the calculated one. Absorption UV-Vis, in water at pH 7.2  $\lambda_{max}(\epsilon/10^4 M^{-1}cm^{-1})$ : 262(12.2) 454(2.2). No corrected emission maximum in water at pH 7.2: 588.0 nm.

## Physical measurements

Microanalyses were carried out on an Elementar Vario EL elemental analyser. Electrospray mass spectra (ESI-MS) were recorded on a LCQ system (Finnigan MAT, USA). The spray voltage, tube lens offset, capillary voltage and capillary temperature were set at 4.50 kV, 30.00 V, 23.00 V and 200 °C, respectively, and the quoted  $m/z$  values

are for the major peaks in the isotope distribution. Emission spectra were measured on a Shimadzu RF-5000 spectrofluorophotometer and UV-Visible absorption was recorded on a Shimadzu UVPC-3000 spectrophotometer. Viscosity experiments were performed on an Ubbelohde viscometer, immersed in a thermostatted water-bath maintained at 30.0 ± 0.1°C. Data were presented as  $(\eta/\eta_0)^{1/3}$  vs. the concentration of [Ru]/[DNA]. Viscosity values were calculated from the observed flow time of DNA-containing solutions ( $t > 100$  s) corrected for the flow time of buffer alone ( $t_0$ ), i.e.,  $\eta=t-t_0$ .

## Results and Discussion

### DNA-binding properties of Ru (II) complexes

**Electronic absorption spectra:** In general, the complex binding to DNA in an intercalation mode exhibits a red and hypochromism shift in the absorption spectra, and the extents of spectral change are closely correlative to the DNA-binding affinities of these complexes. The spectral shifts in an intercalation mode are usually greater than those in groove binding mode. In the presence of double helix calf thymus DNA (CT-DNA), the electronic absorption spectra for all of these complexes exhibit obviously hypochromism, and the hypochromism values for 1, 2, 3 and 4 at MLCT absorption band (452~455 nm) are 12, 9, 9 and 21%, respectively.

In order to clarify the DNA-binding affinities of these complexes, the intrinsic binding constants were calculated according to equation (1) [34], through a plot of

$$[DNA]/\Sigma a-\Sigma f \text{ vs. } [DNA]$$
$$[DNA]/\Sigma a-\Sigma f = [DNA]/\Sigma b-\Sigma f + 1/K_b(\Sigma a-\Sigma f) \quad (1)$$

where [DNA] is the concentration of DNA in base pairs,  $\Sigma a$ ,  $\Sigma f$  and  $\Sigma b$  are respectively the apparent extinction coefficient ( $A_{obsd}/[M]$ ), the extinction coefficient for free metal (M) complex and the extinction coefficient for the metal (M) complex in the fully bound form. In plots of  $[DNA]/\Sigma a-\Sigma f$  versus [DNA],  $K_b$  is given by the ratio of slope to intercept. The calculated values for 1, 2, 3 and 4 at MLCT absorption band are 1.1, 0.35, 0.53 and  $1.7 \times 10^5 M^{-1}$ , respectively. These values are smaller than those for [Ru(bpy)<sub>2</sub>dppz]<sup>2+</sup> ( $> 10^6 M^{-1}$ ) [35] and [Ru(ip)<sub>2</sub>dppz]<sup>2+</sup> ( $2.1 \times 10^7 M^{-1}$ ) [36]. Such DNA-binding constants suggest that the interaction of these complexes with DNA should be in an intercalation mode.

**Emission spectra:** The interaction of Ru (II) complexes with double helix CT-DNA was monitored via luminescence. All ruthenium complexes 1-4 emit luminescence in the range 500-700 with the maximum near 600 nm at room temperature. Upon the addition of CT-DNA, the emission spectra of all of these complexes enhanced obviously (Figure 1). The emission of complex 4 exhibits pronounced enhancement (Figure 2), and its emission intensity increases steadily to 8.5 times relative to that of the original and reaches saturation at  $ca. [DNA]/[Ru]=8:1$ . However, the emission intensities increase by 2.2, 1.7 and 1.5 for complexes 1, 2 and 3, respectively (Figure 2). The enhancement of emission intensities of these complexes can be attributed to the hydrophobic environment inside the DNA helix, which reduces the accessibility of water molecules and makes the mobility of the complexes be restricted at the binding site.

**Viscosity behaviors:** The viscosity experiments, being sensitive to the change of length of double helix DNA, were considered as one of the most unambiguous methods to determine the binding mode of complex to DNA in absence of crystal data [37]. In general, the relative viscosity of DNA in presence of complex in an intercalation mode will be increased, because the intercalative ligand will separate the base




Figure 1: Emission spectra of Ru (II) complexes 1(a), 2(b), 3(c) and 4(d) in absence and in presence of increasing amount of CT-DNA.




Figure 2: Relative emission intensity ( $I/I_0$ ) of Ru (II) complexes 1(■), 2(●), 3(▲) and 4(▼) in absence and in presence of increasing amount of CT-DNA.




Figure 3: Relative viscosity of CT-DNA in absence and in presence of increasing amount of Ru (II) complexes 1(■), 2(●), 3(▲), 4(▼) and  $[Ru(bpy)_3]^{2+}$ (◆) at  $30 \pm 0.1^\circ C$ .

pairs of DNA, and thus lengthen the DNA helix. On the contrary, a partial and/or non-classical intercalation of complex will reduce the relative viscosity of DNA, since the binding ligand may bend (or kink) the DNA helix and reduce its effective length [38]. The experiments on relative viscosity of rod-like CT-DNA in the presence of complexes 1, 2, 3 and 4, as well as  $[Ru(bpy)_3]^{2+}$ , were carried out and the results were shown in Figure 3.

The viscosity of DNA remains almost unchanged upon addition of  $[Ru(bpy)_3]^{2+}$ , which is consistent with an electrostatic association. However, in the presence of Ru (II) complexes 1, 2, 3 and 4 respectively, the relative viscosity of rod-like DNA was increased (Figure 3), because the stacking interaction of these complexes with the base pairs of DNA lengthens the DNA helix, indicating these complexes can bind to DNA in intercalation mode.

## Quantitative structure-activity relationships on ruthenium(II) complexes

Quantitative Structure-Activity Relationships was carried out on MatLab 6.5 for these newly synthesized ruthenium(II) complexes, as well as some congeners from references. The data of intrinsic binding constant and the physical parameters were listed in Table 1.

Firstly, we try to draw a plot with the  $\log K_b$  versus electronic parameter ( $\sigma$ ) on Origin 6.0, as shown in Figure 4, and the corresponding equation is as follows:

$$\log K_b = 0.6726 (\pm 0.1751)\sigma + 4.5001 (\pm 0.09799) \quad (2)$$

$$n=7; r=0.8643; SD=0.1683; p=0.0121$$

Outliner: X=-OCH<sub>3</sub>, Y=-H; X=-CH<sub>3</sub>, Y=-OH; X=-H, Y=-Phen; X=-H, Y=-OH; X=-H, Y=-OCH<sub>3</sub>

Considering there is hydrogen bond exists in some of these ruthenium complexes, we import indicative variable ( $I_H$ -bonding) to indicate hydrogen bond exists in the intercalative ligand. The value of  $I_H$ -bonding is 1 if there is hydrogen bond, regardless it's intramolecular or intermolecular hydrogen bond, and the value of  $I_H$ -bonding is 0 if there is not, thus we get model 2:

$$\log K_b = 0.2770\sigma + 0.2818I_H + 4.6366 \quad (3)$$

$$n=12; R=0.7441; F=5.5824; p=0.0265$$

It's obviously the hydrogen bond contribute to the DNA binding of these ruthenium complexes, since the coefficient  $I_H$ -bonding is positive. Encouraged, we considered that the hydrophobic parameter may also contribute to the DNA-binding properties of these complexes, thus we obtained the model 3:

$$\log K_b = 0.2429\pi + 0.0429\pi^2 + 0.2907\sigma + 0.6389I_H + 4.3491 \quad (4)$$

$$n=12; R=0.9338; F=11.9134; p=0.0030$$

In this model 3, it's obviously the coefficient of electronic parameter ( $\sigma$ ) is positive, indicating the electron acceptor group on intercalative ligand of ruthenium complexes will enhance the binding affinity of ruthenium complexes to DNA, while an electron donor group will decrease the binding affinity. The positive coefficient for the hydrophobic parameter ( $\pi$ ) indicate a hydrophobic group in the intercalative ligand will increase the DNA-binding affinity of ruthenium complexes, while hydrophilic group will decrease the DNA-g affinity.




Figure 4: The relationship of the binding affinity ( $K_b$ ) of ruthenium complexes on the electronic parameter ( $\sigma$ ).

## Conclusion

A series of ruthenium(II) have been synthesized, and the binding-behavior of these ruthenium(II) complexes with calf-thymus DNA have been investigated, and the results show that these complexes can bind to DNA in intercalating mode. The further studies on the quantity structure-activity relationship of these ruthenium complexes, as well as some from reference was investigated, and a QSAR equation was obtained:  $\log K_b = 0.2429\pi + 0.0429\pi^2 + 0.2907\sigma + 0.6389I_H + 4.3491$  ( $n=12$ ;  $R=0.9338$ ;  $F=11.9134$ ;  $p=0.0030$ ). It's shown that the DNA-binding affinity of ruthenium complexes in studied depended on the electronic effect, hydrophobic effect and hydrogen bond, and an electron withdraw group in the intercalative ligand will increase the DNA-binding affinity of ruthenium complexes, while an electron donor group will decrease the DNA-binding affinity. In addition, hydrogen bond is important to obtain a high DNA-binding ruthenium complex.

## Acknowledgements

This work was supported by the National Nature Science Foundation of China (81572926), the Provincial Major Scientific Research Projects in Universities of Guangdong Province (2014KZDXM053), the Science and Technology Project of Guangdong Province (2014A020212312), the Guangzhou City Science and Technology Plan (2013J4100072), the Science and Technology Projects of Yuexiu District (2014-WS-039), and the Joint Natural Sciences Fund of the Department of Science and Technology and the First Affiliated Hospital of Guangdong Pharmaceutical University (GYFYLH201309).

## References

- Caiping T, Shouhai W, Sensen L (2011) Synthesis, structures, cellular uptake and apoptosis-inducing properties of highly cytotoxic ruthenium- Norharman complexes. Dalton Trans 40: 8611-8621.
- Jiang CW, Chao H, Li H, Ji LN (2003) Syntheses, characterization and DNA-binding studies of ruthenium(II) terpyridine complexes:  $[Ru(tpy)(PHBI)]^{2+}$  and  $[Ru(tpy)(PHNI)]^{2+}$ . J Inorg Biochem 93: 247-255.
- Yau HCM, Chan HL, Yang M (2002) Determination of mode of interactions between novel drugs and calf thymus DNA by using quartz crystal resonator. Sensor Actuat B Chem 81: 283-288.
- Patel KK, Plummer EA, Darwish M, Rodger A, Hannon MJ (2002) Aryl substituted ruthenium bis-terpyridine complexes: intercalation and groove binding with DNA. J Inorg Biochem 91: 220-229.
- Vashisht Gopal YN, Konuru N, Kondapi AK (2002) Topoisomerase II antagonism and anticancer activity of coordinated derivatives of  $[RuCl(2)(C(6)H(6))(dmsO)]$ . Arch Biochem Biophys 401: 53-62.
- Bouma M, Nuijen B, Jansen MT, Sava G, Bult A, et al. (2002) Photostability profiles of the experimental antimetastatic ruthenium complex NAMI-A. J Pharm Biomed Anal 30: 1287-1296.
- Novakova O, Hofr C, Brabec V (2000) Modification of natural, double-helical DNA by antitumor cis- and trans- $[Cl(2)(Me(2)SO(4))(4)Ru]$  in cell-free media. Biochem Pharmacol 60: 1761-1771.
- Mishra L, Yadaw AK, Srivastava S (2000) Synthesis, spectroscopic, electrochemical and antibacterial studies of new Ru (II) ,10-phenanthroline complexes containing aryl diazopentane-2,4-dione as co-ligand. New J Chem 24: 505-510.
- Panneerselvam A, Nataraj C, Yoon JJ, Periasamy V (2013) Synthesis, characterization, DNA interaction, antioxidant and anticancer activity of new ruthenium(II) complexes of thiosemicarbazone/semicarbazone bearing 9,10-phenanthrenequinone. J Photoch Photobio B: 17-26.
- Kamatchi TS, Chitrapriya N, Kim SK, Fronczek FR, Natarajan K (2013) Influence of carboxylic acid functionalities in ruthenium(II) polypyridyl complexes on DNA binding, cytotoxicity and antioxidant activity: synthesis, structure and in vitro anticancer activity. Eur J Med Chem 59: 253-264.
- Shobha Devi C, Anil Kumar D, Singh SS, Gabra N, Deepika N, et al. (2013) Synthesis, interaction with DNA, cytotoxicity, cell cycle arrest and apoptotic inducing properties of ruthenium(II) molecular "light switch" complexes. Eur J Med Chem 64: 410-421.
- Huang HL, Li ZZ, Liang ZH, Liu YJ (2011) Cell Cycle Arrest, Cytotoxicity, Apoptosis, DNA-Binding, Photocleavage, and Antioxidant Activity of Octahedral Ruthenium(II) Complexes. Eur J Inorg Chem 2011: 5538-5547.

13. Deepika N, Kumar YP, Devi CS, Reddy PV, Srishailam AS (2013) Synthesis, characterization, and DNA binding, photocleavage, cytotoxicity, cellular uptake, apoptosis, and on-off light switching studies of Ru (II) mixedligand complexes containing 7-fluorodipyrido[3,2-a:2',3'-c] phenazine. *J Biol Inorg Chem* 18: 751-766.
14. Zheng CG, Ma CL, Yu XW, Qian L, Song Y, et al. (2011) Electronic effect of substituents on the DNA intercalation of ruthenium(II) polypyridyl complexes. *Chem Biodiversity* 8: 1486-1496.
15. Chen W, Turro C, Friedman LA, Barton JK, Turro NJ (1997) Resonance raman investigation of ru(phen)2(dppz)2+ and related complexes in water and in the presence of DNA. *J Phys Chem B* 101: 6995-7000.
16. Lasey RC, Banerj SS, Ogawa MY (2000) Synthesis and characterization of a equence-specific DNA-binding protein that contains ruthenium polypyridyl centers. *Inorg Chim Acta* 300-302: 822-828.
17. Moucheron C, Kirsch-De Mesmaeker A, Kelly JM (1997) Photoreactions of ruthenium(II) and osmium(II) complexes with deoxyribonucleic acid (DNA). *J Photochem Photobiol B* 40: 91-106.
18. Goll JG, Thorp HH (1996) Oxidation of DNA by trans-dioxoruthenium(VI) complexes: self-inhibition of DNA cleavage by metal complexes. *Inorg Chim Acta* 242: 219-223.
19. Ji LN, Zou XH, Liu JG (2001) Shape- and enantioselective interaction of ru(II)/co(III) polypyridyl complexes with DNA. *Coord Chem Rev* 216: 513-536.
20. Shobha Devi C, Nagababu P, Natarajan S, Deepika N, Venkat Reddy P, et al. (2014) Cellular uptake, cytotoxicity, apoptosis and DNA-binding investigations of Ru(II) complexes. *Eur J Med Chem* 72: 160-169.
21. Zheng KC, Wang JP, Peng WL, Liu XW, Yun FC (2001) Studies on 6,6'-disubstitution effects of the dpq in [Ru(bpy)2(dpq)]2+ with DFT method. *J Phys Chem A* 105: 10899-10905.
22. Zheng KC, Liu XW, Wang JP, Yun FC, Ji LN (2003) DFT studies on the molecular orbitals and related properties of [Ru(phen)2(9,9' -2R- dpq)]2+(R=NH2,OH,H and F). *J Mol Struct (Theochem)* 637: 195-203.
23. Mei WJ, Liu J, Zheng KC, Lin LJ, Chao H, et al. (2003) Experimental and theoretical study on DNA-binding and photocleavage properties of chiral complexes - and -[Ru(bpy)2L] (L o-hpip, m-hpip and p- hpip). *Dalton Trans* 7: 1352-1359.
24. Sullivan BP, Salmon DJ, Meyer TJ (1978) Mixed phosphine 2,2'-bipyridine complexes of ruthenium. *Inorg Chem* 17: 3334-3341.
25. Arakawa R, Tachiyashiki S, Matsuo T (2002) Detection of reaction intermediates: photosubstitution of (polypyridine) ruthenium(II) complexes using on-line electrospray mass spectrometry. *Anal Chem* 67: 4133-4138.
26. Hammett LP (1934) Some relations between reaction rates and equilibrium constants. *Chem Rev* 17: 125-136.
27. Hansch C, Leo AS, Heller RE (1995) Exploring QSAR. fundamentals and applications in chemistry and biology, American Chemical Society, Washington, DC.
28. Hansch C, Leo A (1979) Substituent constants for correlation analysis in chemistry and biology. John Wiley & Sons, New York.
29. Prasanna S, Manivannan E, Chaturvedi SC (2005) QSAR analyses of conformationally restricted ,5-diaryl pyrazoles as selective COX-2 inhibitors: application of connection table representation of ligands. *Bioorg Med Chem Lett* 15: 2097-2102.
30. Gill MR, Thomas JA (2012) Ruthenium(II) polypyridyl complexes and DNA--from structural probes to cellular imaging and therapeutics. *Chem Soc Rev* 41: 3179-3192.
31. Liu J, Zheng W, Shi S, Tan C, Chen J, et al. (2008) Synthesis, antitumor activity and structure-activity relationships of a series of Ru(II) complexes. *J Inorg Biochem* 102: 193-202.
32. Steck EA, Day AR (1943) Reactions of phenanthraquinone and retenequinone with aldehydes and ammonium acetate in acetic acid solution 1. *J Am Chem Soc* 65: 452-456.
33. Hage R, Haasnoot JG, Reedijk J, Wang R, Ryan EM, et al. (1990) Synthesis, spectroscopic and electrochemical properties and X-ray studies of bis(2,2'-bipyridyl)(3-(2-hydroxy-phenyl)-5-(pyridin-2-yl)-,2,4-triazole)ruthenium(II) hexafluorophosphate. *Inorg Chim Acta* 174: 77-85.
34. Wolfe A, Shimer GH Jr, Meehan T (1987) Polycyclic aromatic hydrocarbons physically intercalate into duplex regions of denatured DNA. *Biochemistry* 26: 6392-6396.
35. Satyanarayana S, Dabrowiak JC, Chaires JB (1993) Tris(phenanthroline) ruthenium(II) enantiomer interactions with DNA: mode and specificity of binding. *Biochemistry* 32: 2573-2584.
36. Shi HJ, Chen Y, Gao F, Yub HJ, Lib GY, et al. (2008) Synthesis, DNA-binding and DNA- photocleavage properties of ruthenium(II) mixed-polypyridyl complex [Ru(tbz)2(dppz)]2+. *J Mol Struct* 892: 485-489.
37. Satyanarayana S, Dabrowiak JC, Chaires JB (1992) Neither delta- nor lambda-tris(phenanthroline)ruthenium(II) binds to DNA by classical intercalation. *Biochemistry* 31: 9319-9324.
38. Liu JG, Ye B H, Li H, Jia L, Li RH, et al. (1999) Synthesis, characterization and DNA- binding properties of novel dipyridophenazine complex of ruthenium(II): [Ru(IP)2(DPPZ)]2+. *J Inorg Biochem* 73: 117-122.
39. Xu H, Zheng KC, Lin LJ, Li H, Gao Y, et al. (2004) Effects of the substitution positions of Br group in intercalative ligand on the DNA-binding behaviors of Ru(II) polypyridyl complexes. *J Inorg Biochem* 98: 87-97.
40. Tan LF, Chao H, Li H, Liu YJ, Sun B, et al. (2005) Synthesis, characterization, DNA-binding and photocleavage studies of [Ru(bpy)2(PPIP)]2+ and [Ru(phen)2(PPIP)]2+. *J Inorg Biochem* 99: 513-520.
41. Xu H, Deng H, Zhang QL, Huang Y, Liu JZ, et al. (2003) Synthesis and spectroscopic RNA binding studies of [Ru(phen)2MHPIP]2+. *Inorg Chem Com* 6: 766-768.

**Citation:** Zhang H, Yang F, Wu Q, Si-Yan L, Liu P, et al. (2016) Synthesis, DNA-binding Properties and Quantitative Structure-Activity Relationships on Ruthenium(II) Complexes with Calf-Thymus DNA. *Med chem (Los Angeles)* 6: 137-142. doi:10.4172/2161-0444.1000338

### OMICS International: Publication Benefits & Features

#### Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

#### Special features:

- 700 Open Access Journals
- 50,000 editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, DOAJ, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission>