

**Open Access** 

# Synthesis of Bioactive Imidazoles: A Review

### Poonam Gupta\* and Jitendra K. Gupta

School of studies in Chemistry, Jiwaji University, Gwalior-474011, India

#### Abstract

Heterocyclic compounds are acquiring more importance in recent years because of their pharmacological activities. The imidazole nucleus is an important synthetic strategy in drug discovery. Imidazole is a planar fivemember ring system with N atom in 1 and 3 positions. The systemic name for the compound is 1, 3 diazole, one of the N bear an H atom and other to be regarded as a pyrrole type N. Imidazole was first named as glyoxaline. It is amphoteric in nature, susceptible to electrophilic and nucleophilic attack. It also occurs in the purine nucleus & amino acid histidine, 4-amino-imidazole-5-carboxamide occurs naturally as a riboside. This interesting group of heterocyclic compound has diverse biological activities such as antimicrobial, anticancer, analgesic, anti-inflammatory, antiviral, anthelmintic, anticonvulsant, antiulcer, anti-allergic activity etc. Numerous methods for the synthesis of imidazoles and also their various structure reactions offer enormous scope in the field of medicinal chemistry. This articles aims to review the work reported, their chemistry and pharmacological activities of imidazole during past years.

**Keywords:** Imidazole; Heterocyclic; Antibacterial; Antiinflammatory; Antifungal and antitumor.

#### Introduction

Heterocyclic compounds are also used in pharmacy and agriculture. Analysis of scientific papers in the last two decades revealed that there is a general trend in research for new drugs involving modification of existing biologically active matrices and molecular design of the structures of compounds. The imidazoles nucleus is an important synthetic strategy in drug discovery. Imidazole derivatives exhibited antimicrobial, anti-inflammatory, analgesic, anti-tubercular and anticancer activity. One of the most important application of imidazole derivatives is their use as material for treatment of denture stomatities. The high therapeutic properties of the imidazole related drugs have encouraged the medicinal chemists to synthesize a large number of novel chemotherapeutic agents. Imidazole drugs have broadened scope in clinical medicines. Medicinal properties of imidazoles include anticancer, anticoagulants, anti-inflammatory, antibacterial, antifungal, antiviral, antitubercular, antidiabetic and antimalarial [1-7]. Imidazole and its derivatives are reported to be physiologically and pharmacologically active and find applications in the treatment of several diseases. Imidazole is an organic compound with the formula (CH), N (NH) CH. It is a colourless solid that dissolves in water to give mildly basic solution. In chemistry, it is an aromatic heterocycle, classified as a diazole and as an alkaloid. Imidazoles are a common component of a large number of natural products and pharmacologically active molecules (Figure 1) Imidazole was first synthesized by Heinrich Debus in 1858, but various imidazole derivatives [8-11] have been discovered as early as the 1840s, it used glyoxal [12] and formaldehyde [13, 14] in ammonia to form imidazole. This synthesis, while producing relatively low yields, is still used for creating C-substituted imidazoles [15].

Imidazoles containing free imino hydrogen and a substituent in the 4- and 5- position, or two dissimilar substituents in these positions, might be expected to occur in the isomeric forms. These isomers differ in the position of the imino hydrogen which may be attached to either of the two nitrogen atoms.

Over the years, the imidazole nucleus has attracted the attention of the scientific community due to its chemical and biological properties [16, 17]. For example, this nucleus is present in the structures of several natural products in the form of the essential amino-acid histidine or in alkaloids exhibiting anti-tumoral, anti-cancer (dacarbazine), antihistaminic (cimetidine), anti-parasitic (metronidazole), and antihypertensive (losartan) and anti-bacterial activities [18-20]. A great numbers of medicines contain the imidazole nucleus, including ketoconazole which are used to treat fungal infections, bacterial infections, and gastric ulcers, respectively (Figure 2) [21, 22]. Due to their importance, it has become an attractive target for the synthetic and medicinal chemist. There are many synthetic methodologies that have been developed for assembling and decorating the imidazole ring with diverse functional groups.

### Synthesis of imidazoles

Bunev et al. [23] have been synthesized a new series of 1,4,5-trisubstituted imidazoles 3 containing trifluoromethyl group has been developed using van Leusen reaction, which incorporates two-component condensation reaction trifluoroacetimidoyl chlorides 1 with tosylmethylisocyanide 2. This protocol provides a novel and improved method for obtaining trifluoromethyl containing 1, 4, 5-trisubstituted imidazoles in good yields (Scheme 1).

Sharma et al. [24] reported two novel series of 2-(substituted phenyl)-1H-imidazole 7 and (substituted phenyl)-[2-(substituted phenyl)-imidazol-1-yl]-methanone 10 analogues it is achieved by the reaction of substituted aniline 5 in HCl/water mixture were diazotized using solution of sodium nitrite. Imidazoles were added in intermediate 6. Compound 10 were synthesized by the reaction of compound 7 in diethyl ether was added to a solution of corresponding benzoic acid 8 with substituted benzoyl chloride 9 (Scheme 2).

Pandya et al. [25] also reported a simple and concise route for the synthesis of highly substituted imidazole derivatives 12 have been

\*Corresponding author: Poonam Gupta, School of studies in Chemistry, Jiwaji University, Gwalior-474011, India. Tel: 1800-233-1964; E-mail: poonamgupta\_001@yahoo.com

Received April 07, 2015; Accepted April 28, 2015; Published May 04, 2015

Citation: Gupta P, Gupta JK (2015) Synthesis of Bioactive Imidazoles: A Review. Chem Sci J 6: 91. doi:10.4172/2150-3494.100091

**Copyright:** © 2015 Gupta P, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



developed by the reaction of 10 with aromatic aniline 11 via coppermediated oxidative C–H functionalization in good to high yields. The advantage of the reaction lies in its mild reaction conditions and readily available starting materials (Scheme 3).

Liu et al. [26] have been synthesized a series of 2-(4-(2-(substituted-1-yl) ethoxy) phenyl)-1H-phenanthro [9, 10-d] Imidazole 15 and 2-(4-(4-(substituted-1-yl) butoxy) phenyl)-1H-phenanthro [9, 10-d] imidazole 16 by multicomponent reaction method (Scheme 4).

Kathrotiya et al. [27] reported a series of some new quinoline based imidazole-5-one derivatives 19, 21 have been synthesized by the fusion of oxazol-5-ones 17, with various p-substituted anilines 18, 20 and zeolite in pyridine (Scheme 5).

Mungra et al. [28] also reported another series of some new tetrazolo[1,5-a]quinoline based tetrasubstituted imidazole derivatives 26 have been synthesized by a reaction of tetrazolo[1,5-a]quinoline-4-carbaldehyde 22, benzyl 23, aromatic amine 25 and ammonium acetate 24 in the presence of iodine through one-pot multi-component reaction (MCR) approach (Scheme 6).

Desai et al. [29] reported the synthesis of N-(4-((2-chloroquinolin-3-yl) methylene)-5-oxo-2-phenyl-4, 5-dihydro-1H-imidazol-1-yl) (aryl) amides 30 by the reaction of 2-chloroquinoline- 3-carbaldehyde 27 and N-amino arylcarboxamides 28 in pyridine. They were react with 4-((2-Chloroquinolin-3-yl) methylene)-2-phenyloxazol-5(4H)-one 29 was heated with again an N-amino arylcarboxamides in pyridine (Scheme 7).

Li et al. [30] have been synthesized trisubstituted imidazoles 33 by the reaction of 1, 2-di (furan-2-yl)-2-oxoethyl carboxylates 31 in

presence of RCOCl they form an intermediate 32 then it is converted into the synthesized compound (Scheme 8).

Chen et al. [31] reported a series of 1-R1-2-R-4, 5-di (furan-2-yl)-1H-imidazole derivatives 39 were synthesized in multisteps. Reaction start with furan-2-carbaldehyde 34 and vitamin B1 to gave 1, 2-di (furan-2-yl)-2-hydroxyethanone 35. 35 react with benzyl chloride or allyl chloride and pyridine to form an intermediate 36, it were react with sodium acetate and gave substituted difuran imidazole 37. This was followed by NaH/THF to form the final compound 38 (Scheme 9).

Ziarani et al. [32] also reported  $SiO_2$ -Pr-SO<sub>3</sub>H catalyst based 1,2,4,5-tetrasubstituted imidazoles 43 by reaction with fourcomponent, one-pot reaction of 1,2-diketones 39, aryl aldehydes 40, ammonium acetate 41 and substituted aromatic amines 42 in excellent yields under solvent free conditions (Scheme 10).

Jawaharlal et al. [33] reported tetrasubstituted imidazole 48 by the refluxing of 9, 10-phenenthraquinone 44 with aryl aldehyde 46, primary amines 47 and ammonium acetate 45 in the presence of glacial acetic acid (Scheme 11).

Lavanya et al. [34] also reported the synthesis of 6-bromo-2substitutedphenyl-1H-imidazo [4, 5-b] pyridine derivatives 51 were prepared with 5-Bromopyridine-2, 3-diamine 49 underwent facile condensation with various aromatic carboxylic acid derivatives 50 in the presence of Etan's reagent (Scheme 12).

Prabhu et al. [35] reported another series of some novel aryl imidazole derivatives 57 were prepared by the condensation of compounds containing primary aromatic amine 52 and aryl aldehydes 53 to give respective Schiff's bases 54, which was further reacted with ammonium acetate 55 and isatin 56 in the presence of glacial acetic acid (Scheme 13).

Sathe et al. [36] have synthesized 4-Fluoro-3-chloroanilline 58 treated with potassium thiocyanate in presence of glacial acetic acid and bromine was converted into 2-amino-6-fluoro-7-chlorobenzothiazole 59, resulting into 2-amino benzothiazole. The synthesized compound in presence of 2-phenyl-4-benzylidine-5-oxazolinone 60 refluxed in pyridine to obtained 2- (2- Phenyl - 4 - benzylidenyl - 5 - oxo - imidazolin - 1 - ylamino) - 6 - fluoro -7-substituted (1, 3) benzothiazoles 62, 61 (Scheme 14).

Stella et al. [37] reported an efficient and practical synthesis of imidazolyl derivatives 65 were achieved through thiocyanation of aniline derivatives 63 to gave the intermediate 64 which followed by the reaction with ethylene diamine in the presence of carbondisulphide (Scheme 15).

Lakshmanan et al. [38] also reported the synthesis of 1-(4-substitutedphenyl)-2-(2-methyl-1H-imidazol-1-yl) ethanone 67 and synthesis of 1-(4-substituted phenyl)-2-(1H-imidazol-1-yl) ethanone 68 by the reaction of para substituted phenacyl bromides 66 with imidazoles (Scheme 16).

Husain et al. [39] reported a series of 1,2,4-trisubstituted-1 Himidazoles 72 were synthesized by the 2,4-disubstituted-1 H- imidazoles 71 and the title compounds were synthesized from 4-methoxyphenyl glyoxal 70 with following multistep synthesis (Scheme 17).

## Pharmacological Profile of Imidazoles

Imidazole and their derivatives are still the most widely used in the therapeutic areas and have shown a broad spectrum of activity against various pathogens. Since the discovery of various drugs







Volume 6 • Issue 2 • 100091

Chem Sci J ISSN: 2150-3494 CSJ, an open access journal















H₌Ča

Scheme 14: Synthesis of 2 - (2 - Phenyl - 4 - benzylidenyl - 5 - oxo - imidazolin - 1 - ylamino) - 6 -fluoro-7-substituted (1, 3) benzothiazoles.

62

HaC

61

Page 6 of 12



Scheme 17: Synthesis of 1, 2, 4 trisubstituted 1-H-imidazoles.

72

Page 8 of 12

S.No.	Chemical Structure	Chemical Name	Activity
1	Ph CH <sub>3</sub> NH ON	2-Cyano-3-(4-fluorophenyl)- <i>N</i> '-[1-(5-methyl-2- phenyl-1 <i>H</i> -imidazol- 4-yl)ethylidene]acrylohydrazie	Antimicrobial, antioxidant, anti- hemolytic and cytotoxic [40].
2	$R^{1}$ $R^{1}$ $R^{2}$ $R^{1}$ $R^{1}$ $R^{2}$ $R^{1}$ $R^{1$	1-[2-(1 <i>H</i> -imidazol-1-yl)acetyl]-3-methyl-2,6- diphenylpiperidin-4-one	Antibacterial and antifungal [41].
3		2,4-Dichloro- <i>N-</i> (4-(4-chloro-1 <i>H-</i> imidazol-1-yl)-3- methoxyphenyl)benzamide	Antimicrobial and Antitubercular [42].
4	$H_{2N}$	transition metal(II) complexes of imidazole-2- carbaldehyde semicarbazone (H <sub>2</sub> L)	Antimicrobial [43].
5	$R^{1}$ $R^{2}$ $R^{1}$ $R^{2}$ $R^{1}$ $R^{2}$ $R^{3}$ $R^{3}$ $R^{3}$ $R^{3}$ $R^{3}$ $R^{3}$ $R^{3}$ $R^{3}$	(2-aryl-1 <i>H-</i> imidazol- 4-yl)methanone	Antiproliferative [44].
6		3-(4-((4-(1 <i>H</i> -phenanthro[9,10-d]imidazol-2-yl) phenoxy) methyl)-1 <i>H</i> -1,2,3-triazol-1-yl)propan-1-amine	Alzheimer's disease [21].

Page 9 of 12

7		2,4,5-triphenyl-1 <i>H-</i> imidazole	Antimicrobial [45].
8		2-[4-(4,5-Diphenyl-1 <i>H-</i> imidazol-2-yl)- phenyl]-6-(40- mathoxy-biphenyl-4-yl)-pyridine (MPBI)	Antibacterial, Antifungal [46].
9	S = N = N = N = N $HN = N = N = N$ $Ar$ $HN = Ar$ $HN = Ar$ $HN = Ar$	methyl(2Z)-[3- ({(E)-[3-aryl-1 <i>H</i> -pyrazol-4-yl]methylidene}amino)- 5-oxo-2- thioxoimidazolidin-4-ylidene]ethanoate	Antibacterial [47].
10	H <sub>3</sub> C N O N O H <sub>3</sub> C O R	3-(1-(4-methoxybenzyl)-2-butyl-4-chloro-1 <i>H-</i> imidazol-5-yl-1-arylprop-2-en-1-one	antibacterial, antifungal [48].
11	R N N CH <sub>3</sub> R CH <sub>3</sub>	(5Z)-5-[4- (dimethylamino)benzylidene]-3-(5- substituted -1,3,4-oxadiazol -2-yl)-2-phenyl- 3,5-dihydro-4 <i>H</i> -imidazol-4-one	Anthelmintic [49].
12		4-[2-(2-hydroxyphenyl)imidazo[4,5-b]indol-3(4 <i>H</i> )- yl]benzenesulfonamide	antibacterial and antihelmintic [35].
13	$F \xrightarrow{\mathbf{N}} S \xrightarrow{\mathbf{N}} H_{5} \mathbf{C}_{6}$	2[2'- Phenyl - 4'- benzidinyl- 5'- oxo- imidazoline- 1yl- amino] -6 fluoro- 7- chloro (1,3) benzothiazole	anti-inflammatory [36].

Page 10 of 12

14	4-(4methoxyphenyl)-1,2-diphenyl-1 <i>H-</i> imidazole	Anti-inflammatory, antifungal [39].
15	2,3-dihydroimidazo[1,2-b][1,4,2]benzodithiazines	Anti-HIV [50].
16	N-(2,4-dihydroxybenzylidene)- 2-(2-(phenylthiomethyl)-1 <i>H</i> -benzo[d]-imidazol-1- yl)acetohydrazide	Antitumor [51].
17	3-bromo-3-deazaneplanocin	Antiviral [52].

Table 1: Various imidazole derivatives show various pharmacological activities.

contain the imidazole nucleus, including ketoconazole, metronidazole and cimetidine, which are used to treat fungal infections, bacterial infections and gastric ulcers, respectively. On the basis of various literature surveys imidazole derivatives shows various pharmacological activities (Table 1).

#### Conclusion

Imidazole is a five membered heterocyclic compound. There were so many different conventional methods to synthesize imidazole and its derivatives. On the basis of the literature it was found that imidazole was synthesized under solvent free condition and refluxing method with the help of efficient and different catalyst and without catalyst with good yield. Imidazole is a base in nature due to nitrogen atom. It under goes electrophilic substitution but nucleophilic substitution is rare one. From the extensive literature survey it was found that it has antimicrobial, anticancer, analgesic, antiinflammatory, anticonvulsant, antiviral, anthelmintic, antiulcer, antiallergic activity etc. So from the above discussion it can be concluded that imidazole is a therapeutically active versatile moiety, which had been exploited in the past years for synthesizing various compounds having diverse pharmacological activities, and still imidazole can be further utilized for the future prospective against various diseases or disorders.

#### Acknowledgements

We are grateful to the Head of the Department, School of studies in Chemistry, Jiwaji University Gwalior and Dean, Birla Institute of Medical Research and College of Life Sciences, Gwalior.

#### References

- Congiu C, Cocco MT, Onnis V (2008) Design, synthesis, and in vitro antitumor activity of new 1, 4-diarylimidazole-2-ones and their 2-thione analogues. Bioorg Med Chem Lett 18: 989–993.
- Siddiqui IR, Singh PK, Srivastava V, Singh J (2010) Facile synthesis of acyclic analogues of carbocyclic nucleoside as potential anti-HIV pro-drug. Indian Journal of Chemistry 49B: 512-520.
- Lin YI, Peterson PJ, Yang Y, Weiss WJ, Shales DM, et al. (2008) 5,5,6-Fused tricycles bearing imidazole and pyrazole 6-methylidene penems as broadspectrum inhibitors of ß-lactamases. Bioorg Med Chem Lett 16: 1890–1902.

- Nakamura T, Kakinuma H, Umemiya H, Amada H, Miyata N, et al. (2004) Imidazole derivatives as new potent and selective 20-HETE synthase inhibitors. Bioorg Med Chem Lett 14: 333–336.
- Roman G, Riley JG, Vlahakis JZ, Kinobe RT, Brien JF, et al. (2007) Heme oxygenase inhibition by 2-oxy-substituted 1-(1H-imidazol-1-yl)-4phenylbutanes: effect of halogen substitution in the phenyl ring. Bioorg Med Chem 15: 3225-3234.
- Babizhayev MA (2006) Biological activities of the natural imidazole-containing peptidomimetics n-acetylcarnosine, carcinine and L-carnosine in ophthalmic and skin care products. Life Sci 78: 2343-2357.
- Nantermet PG, Barrow JC, Lindsley SR, Young M, Mao S, et al. (2004) Imidazole acetic acid TAFIa inhibitors: SAR studies centered around the basic P(1)(') group. Bioorg Med Chem Lett 14: 2141-2145.
- 8. Brown EG (1998) Ring Nitrogen and Key Biomolecules, Kluwer Academic Press.
- 9. Pozharskii AF (1997) Heterocycles in Life and Society, John Wiley & Sons.
- Rondu F, Le Bihan G, Wang X, Lamouri A, Touboul E, et al. (1997) Design and synthesis of imidazoline derivatives active on glucose homeostasis in a rat model of type II diabetes. 1. Synthesis and biological activities of N-benzyl-N'-(arylalkyl)-2-(4', 5'-dihydro-1'H-imidazol-2'-yl) piperazines. J Med Chem 40: 3793-3803.
- Hayashi T, Kishi E, Soloshonok VA, Uozumi Y (1996) Erythro-Selective aldoltype reaction of N-sulfonylaldimines with methyl isocyanoacetate catalyzed by gold(I). Tetrahedron Letters 37: 4969-4972.
- Jung ME, Huang A (2000) Use of optically active cyclic N, N-dialkyl aminals in asymmetric induction. Org Lett 2: 2659-2661.
- Corey EJ, Grogan MJ (1999) Enantioselective synthesis of alpha-amino nitriles from N-benzhydryl imines and HCN with a chiral bicyclic guanidine as catalyst. Org Lett 1: 157-160.
- Isobe T, Fukuda K, Araki Y, Ishikawa T (2001) Modified guanidines as chiral superbases: the first example of asymmetric silylation of secondary alcohols. Chemical Communication 243-244.
- Fujioka H, Murai K, Ohba Y, Hiramastu A, Kita YA (2005) A mild and efficient one-pot synthesis of 2-dihydroimidazoles from aldehydes. Tetrahedron Letters 46: 2197-2199.
- Narasimhan B, Deepika S, Kumar P (2011) Biological importance of imidazole nucleus in the new millennium. Med Chem Research 20: 1119–1140.
- Botta M, Corelli F, Gasparrini F, Messina F, Mugnaini CJ et al. (2000) Chiral Azole Derivatives. Enantiomers of Bifonazole and Related Antifungal Agents: Synthesis, Configuration Assignment, and Biological Evaluation. Organic Chemistry 65: 4736–4739.
- Jin Z (2006) Imidazole, oxazole and thiazole alkaloids. Nat Prod Rep 23: 464-496.
- Molina P, Tárraga A, Otón F (2012) Imidazole derivatives: a comprehensive survey of their recognition properties. Org Biomol Chem 10: 1711-1724.
- 20. Jin Z (2013) Muscarine, imidaozle, oxazole and thiazole alkaloids. Nat Prod Rep 30: 869-915.
- Kuma JR (2010) Review of Imidazole heterocyclic ring containing compounds with their biological activity. Pharmacophore 1: 167–177.
- 22. Bendaha H, Yu L, Touzani R, Souane R, Giaever G (2011) New azole antifungal agents with novel modes of action: Synthesis and biological studies of new tridentate ligands based on pyrazole and triazole. Eur J Med Chem 46: 4117-4124.
- Alexander SB, Maksim AV, Vladimir ES, Gennady IO, Alexander SP (2014) Synthesis of 1-aryl-4-tosyl-5-(trifluoromethyl)-1H-imidazoles. Journal of Fluorine Chemistry 163: 34-37.
- Sharma D, Narasimhan BS, Kumar P, Judge VJ, Narang R (2009) Synthesis, antimicrobial and antiviral evaluation of substituted imidazole Derivatives. European Journal of Medicinal Chemistry 44: 2347–2353.
- Pandya AN, Agrawal DK2 (2014) A concise synthesis of highly substituted imidazoles via copper-mediated oxidative C-H functionalization. See comment in PubMed Commons below Tetrahedron Lett 55: 1835-1838.
- 26. Liu J, Qiu J, Wang M, Wang L, Su L, et al. (2014) Synthesis and characterization

of 1H-phenanthro[9,10-d]imidazole derivatives as multifunctional agents for treatment of Alzheimer's disease. See comment in PubMed Commons below Biochim Biophys Acta 1840: 2886-2903.

- Kathrotiya HG, Patel NA, Patel RG, Patel MP (2012) An efficient synthesis of 3-quinolinyl substituted imidazole-5-one derivatives catalyzed by zeolite and their antimicrobial activity. Chinese Chemical Letters 23: 273–276.
- Mungra DC, Kathrotiya HG, Ladani NK, Patel MP, Patel RG(2012) Molecular iodine catalyzed synthesis of tetrazolo[1,5-a]-quinoline based imidazoles as a new class of antimicrobial and antituberculosis agents. Chinese Chemical Letters 23:1367–1370.
- Desai NC, Maheta AS, Rajpara KM, Oshi VVJ, Vaghani HV et al.(2011) Green synthesis of novel quinoline based imidazole derivatives and evaluation of their antimicrobial activity. Journal of Saudi Chemical Society.
- 30. Li B, Gu Q, He Y, Zhao T, Wang T, et al.(2012) Facile synthesis of trisubstituted imidazoles from 1,2-di(furan-2-yl)-2-oxoethyl carboxylates and their chemiluminescence. Comptes Rendus Chimie 15: 784-792.
- Chen Y, Gu Q, Li B, Chen Q, Chen X et al.(2013)Efficient synthesis of 1-R1-2-R-4,5-di(furan-2-yl)-1H-imidazoles and their luminescence properties. Comptes Rendus Chimie 16: 1103–1110.
- Ziarani GM, Dashtianeh Z, Nahad MS, Badiei A(2014)One-pot synthesis of 1,2,4,5-tetra substituted imidazoles using sulfonic acid functionalized silica (SiO2-Pr-SO3H), Arabian Journal of Chemistry.
- Jawaharmala HS, Lambab SN, Singh G, Sainid DR, Kaur AD, Narwal NS(2012) Synthesis of Imidazole Compounds and Evaluation of Their Antimicrobial Activity. Indo Global Journal of Pharmaceutical Science 2: 147-156.
- 34. Lavanya P, Suresh M, KotaiahY, Harikrishna N, Rao CV(2011) Synthesis, antibacterial, antifungal and antioxidant activity studies on 6-bromo-2substituted phenyl-1H-imidazo[4,5-b] pyridine. Asian Journal of Pharmaceutical and Clinical Research 4: 69-73.
- Prabhu M, Radha R (2012) Synthesis, characterization and evaluation of antibacterial and antihelmintic activity of some novel aryl imidazole derivatives. Asian Journal of Pharmaceutical and Clinical Research 5 : 154-159.
- 36. Sathe BS Jagtap VA, Deshmukh SD, Jain BV (2011) Screening of in-vitro antiinflammatory activity of some newly synthesized fluorinated benzothiazolo imidazole compounds. International Journal of Pharmacy and Pharmaceutical Sciences 3: 220-222.
- Stella CR, Rajam S, Venkatraman BR (2012) Characterization and antimicrobial evaluation of newly synthesized imidazole derivatives. International Journal of Chem Tech Research 4:1447-1450.
- Lakshmanan B, Mazumder PM, Sasmal D, Ganguly S (2010) Biologically Active Azoles: Synthesis, Characterization and Antimicrobial Activity of Some 1-substituted imidazoles. Der Pharmacia Lettre, 2: 82-89.
- Husain A, Drabu S, Kumar N, Alam MM, Bawa S (2013) Synthesis and biological evaluation of di- and tri-substituted imidazoles as safer anti-inflammatoryantifungal agents. J Pharm Bioallied Sci 5: 154-161.
- Abdel-Wahab BF, Awad GEA, Badria FA (2011) Synthesis, antimicrobial, antioxidant, anti-hemolytic and cytotoxic evaluation of new imidazole-based heterocycles. European Journal of Medicinal Chemistry 46: 1505-1511.
- Ramachandran R, Rani M, Senthan S, Jeong YT, Kabilan S (2011) Synthesis, spectral, crystal structure and in vitro antimicrobial evaluation of imidazole/ benzotriazole substituted piperidin-4-one derivatives. Eur J Med Chem 46: 1926-1934.
- 42. Ranjith PK, Pakkath R, Haridas KR, Kumari SN (2014) Synthesis and characterization of new N-(4-(4-chloro-1H-imidazol-1- yl)-3-methoxyphenyl) amide/sulfonamide derivatives as possible antimicrobial and antitubercular agents. European Journal of Medicinal Chemistry 71: 354-365.
- 43. Rodríguez-Argüelles MC, Mosquera-Vázquez S, Sanmartín-Matalobos J, García-Deibe AM, Pelizzi C et al. (2010) Evaluation of the antimicrobial activity of some chloro complexes of imidazole-2-carbaldehyde semicarbazone: X-ray crystal structure of cis-NiCl2(H2L) (H2O) Polyhedron 29: 864–870.
- 44. Chen J, Ming Li C, Wang J, Ahn S, Wang Z et al. (2011) Synthesis and antiproliferative activity of novel 2-aryl-4-benzoyl-imidazole derivatives targeting tubulin polymerization. Bioorganic & Medicinal Chemistry 19: 4782–4795.
- 45. Jourshari MS, Mamaghani M, Shirini F, Tabatabaeian K, Rassa M et al. (2013) An expedient one-pot synthesis of highly substituted imidazoles using supported

ionic liquid-like phase (SILLP) as a green and efficient catalyst and evaluation of their anti-microbial activity. Chinese Chemical Letters, 24: 993–996.

- 46. Nagarajan N, Vanitha J, Ananth DA, Rameshkumar S, Sivasudha T, Renganathan R(2013) Bioimaging, antibacterial and antifungal properties of imidazole-pyridine fluorophores: Synthesis, characterization and solvatochromism. Journal of Photochemistry and Photobiology B Biology 127: 212–222.
- Vijesh AM, Isloor AM, Telkar S, Arulmoli T, Hoong-Kun F (2013) Molecular docking studies of some new imidazole derivatives for antimicrobial properties. Arabian Journal of Chemistry, 6: 197–204.
- Maheta HK, Patel AS, Naliapara YT (2012) Synthesis and microbial study of some novel cyanopyrans and cyanopyridines containing imidazole nucleus. International Journal of Chemical Science, 10: 1815-1829.
- 49. Patel K, Jayachandran E, Shah R, Avali V, Sreenivasa GM (2010) Synthesis characterization and antihelmintic activity (perituma posthuma) of new oxadiazole incorporated with imidazole and pyrazole. International Journal of Pharma and Bio Sciences 1: 1-13.
- 50. Brzozowski Z, Saczewski F, Neamati N (2006) Synthesis and anti-HIV-1 activity of a novel series of 1, 4, 2-benzodithiazine-dioxides. Bioorg Med Chem Lett 16: 5298-5302.
- 51. Liu T, Sun C, Xing X, Jing L, Tan R, et al. (2012) Synthesis and evaluation of 2-[2-(phenylthiomethyl)-1H-benzo[d] imidazol-1-yl)acetohydrazide derivatives as antitumor agents. Bioorg Med Chem Lett 22: 3122-3125.
- 52. Liu C, Chen Q, Schneller SW (2012) 3-Bromo-3-deazaneplanocin and 3-bromo-3-deazaaristeromycin: synthesis and antiviral activity. Bioorg Med Chem Lett 22: 5182-5184.