


Synthesis of the New 2-[6-Nitro-2-Benzothiazolylazo]-4-Hydroxy Benzoic Acid Organic Reagent for Spectrophotometric Determination of Copper(II)

Aqeel Mahdi Jreo*

Department of Chemistry, College of Education, Kufa University, Iraq

Abstract

A new 2-[6-Nitro-2-benzothiazolylazo]-4-hydroxy benzoic acid (NO₂BTAHB) organic reagent was synthesized. A sensitive and selective spectrophotometric method was proposed for the rapid determination of Cu(II) using (NO₂BTAHB) reagent. The reaction between Cu(II) and (NO₂BTAHB) reagent is instantaneous at pH=6.0 and the absorbance remains stable for over 24 hrs. The Method allows for the determination of Cu(II) over the range (0.1-6.0) µg.ml⁻¹, with molar absorptivity of (7.45 × 10³)l.mol⁻¹.cm⁻¹ and a detection limit of 0.0245 µg.ml⁻¹. Recovery and relative error values of precision and accuracy of method were found to be R.S.D=1.7%, Re=98.6%, and Erel=-1.4%. The properties of complex was studied and show; (M:R) ratio was 1:2 at pH=6.0, and the stability constant of 7.796 × 10⁻⁹ L².mol⁻². The interferences of ions (Ni²⁺, CrO₄²⁻, Ca²⁺, pb²⁺, Cu²⁺, WO₄²⁻, MO₄²⁻, Co²⁺, Mg²⁺, Cd²⁺, Ba²⁺, Bi³⁺) and masking agents effect on absorbance were studied.



Keywords: Ligand; Ion; Cu; Determination; Reagent

Introduction

Copper plays a different role in the human body. It is an essential nutrient or a toxic element for human beings, depending on the concentration level [1,2]. Some chromogenic reagents have been used in spectrophotometric methods of determination of copper such as Acetophenone-p-chlorophenylthiosemicarbazone [3], Hydrazinecarboxymide-2-[(2-hydroxyphenyl) methylene] (HC22HPM) [4], Piperazine [5], Chloro(phenyl) glyoxime [6], sodium(I) diethyldithiocarbamate [7], brilliant cresyl blue(BCB)[8], cefixime [9], 4-(6-Bromobenzothiazolylazo)Orcinol [10], 3-(2'-thiazolylazo)-2,6-diaminopyridine [11], 2-[2-(4-methylbenzothiazolyl)azo]-5-dimethylaminobenzoic acid [12], 2[2-(5-bromo thiazolyl) azo]-4-methoxy phenol [13], and 2,6-dichlorophenolindophenol [14]. Thiazolylazo compounds have attracted the attention, as they are sensitive chromogenic reagents in addition to being important complexing agents. These dyes are useful in spectrophotometric determinations due to their good selectivity over a wide range of pH and they are relatively easy to synthesize and purified [15]. In this paper, a new (NO₂BTAHB) chromogenic reagent was synthesized for spectrophotometric determination of Cu(II).


Reagents

All reagents were of analytical grade. Freshly distilled and deionized water was used for solutions preparations

Preparation of reagent [16]

To a mixture of {(4.3 gm of para nitro aniline and 3.8 gm of ammonium thiocyanate) in 70 ml glacial acetic acid}, was added drop by drop from burette (1.2 ml Br₂ + 15 ml glacial acetic acid) keeping at temperature >10°C. After 15 minutes alkaline solution was added to precipitate the thiazole derivative, 1.145 gm of thiazole and in 50 ml

glacial acetic acid then add (5 ml conc. HCl + 25 ml water) to the solution. After that drop by drop from burette a solution (0.690 gm NaNO₂ + 50 ml H₂O) with stirring at 0-5°C to diazonium salt, then (0.1.390 gm of parahydroxy benzoic acid + 50 ml ethanol) is added to diazonium salt and 2-[6-nitro-2-benzothiazolyl azo]-4- hydroxy benzoic acid (NO₂BTAHB)organic reagent was formed.



*Corresponding author: Aqeel Mahdi Jreo, Department of Chemistry, College of Education, Kufa University, Iraq, Tel: +96433 340952; E-mail: aqeel-me79@gmail.com

Received: March 20, 2015; Accepted: March 28, 2015; Published April 06, 2015

Citation: Jreo AM (2015) Synthesis of the New 2-[6-Nitro-2-Benzothiazolylazo]-4-Hydroxy Benzoic Acid Organic Reagent for Spectrophotometric Determination of Copper(II). Biochem Anal Biochem 4: 167. doi:10.4172/2161-1009.1000167

Copyright: © 2015 Jreo AM. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Standard solutions

Stock Cu(II) solution; A solution of Cu(II) ($100 \mu\text{g}\cdot\text{ml}^{-1}$) was prepared by dissolving (0.0392) gm of $\text{CuSO}_4\cdot 5\text{H}_2\text{O}$ in (100 ml) distilled water. Other standard solutions of Cu(II) were prepared by dilution of stock solution with distilled water. $1 \times 10^{-3}\text{M}$ (NO_2BTAHB) standard solution was prepared by dissolving (0.088) g in 250 ml of absolute ethanol. -Buffer solution [17] ($\text{pH}=6.0$) was prepared by mixing 12.63 ml of (0.2)M Na_2HPO_4 (which was prepared by dissolving 2.83 gm in 100 ml distilled water) and 7.37 ml of (0.1) M Citric acid (which was prepared by dissolving 1.92 gm in 100 ml distilled water).

Apparatus

Spectrophotometric measurements were made with a Shimadzu (UV-Vis.) scientific equipment with 1.0 cm cell for plot spectra. The pD-303. Spectrophotometer, APEL Japan, was used in the other measurements. The pH-meter, 720 WTW, Germany and FT-IR Spectrophotometer shimadzu, Japan, were used in this work.

Procedure

To an aliquot containing $\leq 10 \mu\text{g}\cdot\text{ml}^{-1}$ of Cu(II) in a 10 ml volumetric flask, was added 2 ml of buffer solution, and 3.5 ml of ($2 \times 10^{-4}\text{M}$) of (NO_2BTAHB) solution. The solution was diluted to the mark with distilled water, and absorbance was measured at 25°C and wave length of 618 nm against the reagent solution as a blank solution prepared under the same conditions.


Results and Discussion

FT-IR spectrum of reagent (NO_2BTAHB)

The following table shows the main vibration frequencies of main absorption bands characteristic of reagent (Table 1 and Figure 1).

Properties of the (NO_2BTAHB)

(NO_2BTAHB) reagent is slightly soluble in water, red powder, orange and stable solution for suitable period time, but in basic medium $\text{pH} \geq 8.0$ the solution being pink. Such behavior may be interpreted by the following equilibria (Figure 2).



Study of Cu(II)-(NO_2BTAHB)complex

Absorption spectra: a-Ultra violet-visible absorption spectra of (NO_2BTAHB) reagent, and Cu(II)-(NO_2BTAHB)complex solution are shown in Figure 3. The reagent showed an absorption maximum at 439 nm, and the complex at 618 nm.

FT-IR spectrum of Cu-(NO_2BTAHB) complex

Changing in intensities, shift in peaks positions, and fission in azo peak were seen which indicate to formation of complex as in following Figure 4.

Effect of pH

The effect of pH was studied over the rang (2-9) adjusted by means of dilute HCl and NaOH solution. Figure 5 shows the relationship between absorbance and pH, where the maximum absorbance obtained in the range of $\text{pH}=(4.0-7.5)$. At $7.5 < \text{pH} < 4.0$ a decrease in absorbance. Therefore, the optimum pH was 6.0, where the

absorbance was maximum and constant.

Effect of Time

The stability of complex was studied from (0-120) min with 5 minutes intervals up to 24 hrs. The maximum absorbance was reached at 10 minutes Figure 6 after that the absorbance remains constant.

Effect of Temperature

The effect of temperature on absorbance of complex was studied; the study was performed at temperature between ($5-80$) $^\circ\text{C}$. Figure 7 show the maximum absorbance obtained at temperature range ($15-40$) $^\circ\text{C}$ which was regarded as a proper temperature of complex formation. At temperatures higher than 40°C the absorbance decrease due to dissociation of complex gradually.


Determination of Stoichiometry and Formation Constant

The composing of complex was studied by jobs method of continuous variations and mole ratio method [18]. Figures 8 and 9 both methods indicate that the ratio of metal ion to reagent molecules (M:L) was (1:2) at $\text{pH}=6.0$.

The formation constant calculated by applied procedure, was found to be (7.796×10^{19}) $\text{L}^2 \cdot \text{mol}^{-2}$

Suggestion of Structural Formula of Cu(II)-(NO_2BTAHB) Complex

From the obtained results of metal to reagent ratio, and depending on thiazolylazo Compounds properties; the following structure can be suggested;



Analytical Characteristics

Calibration curve

Linear calibration graph through the origin was obtained which obeyed Beers law over the range ($0.1-6.0$) $\mu\text{g}\cdot\text{ml}^{-1}$ of Cu(II). The average molar absorptivity was found to be (7.45×10^3) $\text{l}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$. The sandells sensitivity [19] was (0.0088) μg of Cu(II). cm^{-2} , and correlation coefficient (r) was 0.992 (Figure 10).

Wave number (Cm ⁻¹)	Groups
3300-3400	γO-H , H ₂ O Cryst.
2870	γC-H Aliphatic
3020	γC-H Aromatic
1740	γC=N
1505	γN=N
1422	γC=C
1128	γC-S
1280	γC-O phenolic
1678	γC=O carboxylic
1362	γC-N

Table 1: Main vibration frequencies of main absorption bands characteristic of reagent.




Figure 1: A typical sigmoid curve describing the kinetics of amyloid formation. The use of Equations (5)-(7) for analysis of the kinetics of protein fibrillation.




Figure 2: Uv-Visible spectrum of (NO₂)₂TAHB reagent.




Figure 3: Uv-Visible spectrum of Cu(II)-(NO₂BTAHB) complex.




Figure 4: FT-IR spectrum of Cu(II)-(NO₂BTAHB) complex.




Figure 5: Effect of pH on absorbance Cu(II)-(NO₂BTAHB) complex.




Figure 6: Effect of time on the stability of Cu(II)-(NO₂BTAHB) complex.




Figure 7: Effect of temperature on the stability of Cu(II)-(NO₂BTAHB) complex.




Figure 8: Mole ratio plot, pH =6.0.




Figure 9: Jobs plot, pH=6.0.




Figure 10: Calibration curve of Cu(II)-(NO₂BTAHB) complex.

Precision and Accuracy

The relative standard deviation (R.S.D%), evaluated from seven independent determination of 3.0 $\mu\text{g}\cdot\text{ml}^{-1}$ of Cu(II) was 1.7%, this result show that this method is highly precise. Also the accuracy of this method was

determined by calculated the Erel percentage for 3.0 $\mu\text{g}\cdot\text{ml}^{-1}$ standard solution of Cu(II) which was found to be (-1.4) and Re%=98.6.

Interferences

The effect of the ions (Ni^{2+} , CrO_4^{2-} , Ca^{2+} , pb^{2+} , Cu^{+2} , WO_4^{2-} , MO_4^{2-} , Co^{2+} , Mg^{2+} , Cd^{2+} , Ba^{2+} , Bi^{3+}) which form complex with the reagent during its reaction with Cu(II) were studied. On the other hand, suitable masking agents examined for eliminating the effect of the twelve ions, where the mixture of KI, NaF, $\text{Na}_2\text{S}_2\text{O}_3$, and DMG were found to be a suitable masking agents.

References

1. Wen XD, Yang QL, Yan ZD, Deng QW (2011) Determination of cadmium and copper in water and food samples by dispersive liquid-liquid microextraction combined with UV-vis spectrophotometry. *Microchem J* 97: 249-254.
2. Ganjali MR, Aghabalazadeh S, Khoobi M, Ramazani A, Foroumadi A (2011) Nanocomposite Based Carbon Paste Electrode for Selective Analysis of Copper. *Int J Electrochem Sci* 6: 52-62.
3. Ghazy SE, El-Shazly RM, El-Shahawi MS, Al-Hazmi GAA, El-Asmy AA (2006) Spectrophotometric Determination of Copper (II) in Natural Waters, Vitamins and Certified Steel Scrap Samples Using Acetophenone-p-chlorophenylthiosemicarbazone. *J.Iranian Chemical Society* 3: 140-150.
4. Lokhande RS, Kulkarni S, Pitale S, Patil SK, Janwadkar SP (2011) Extraction and Spectrophotometric Determination of Cu (II) Metal ions using Hydrazinecarboxymide-2-[(2-hydroxyphenyl) methylene (HC22HPM) as an Analytical Reagent. *International Journal of Pharma Sciences and Research (IJPSR)* 2: 184-188.
5. Kavitha C, SarathBabu M, Saraswathi K (2013) Spectrophotometric determination of copper as copper Piperazine. *International Letters of Chemistry, Physics and Astronomy* 8: 205-209.
6. Turkoglu O, Soylak M (2005) Spectrophotometric Determination of Copper in Natural Waters and Pharmaceutical Samples with Chloro(phenyl) glyoxime. *Journal of the Chinese Chemical Society* 52: 575-579.
7. Jankiewicz B, Ptaszyński B, Turek A (1999) Spectrophotometric Determination of Copper (II) in Samples of Soil from Selected Allotment Gardens in Lodz 8: 35-38.
8. Ulusoy HI, Gurkan R, Akcay M (2011) Kinetic spectrophotometric determination of trace copper (II) ions by their catalytic effect on the reduction of brilliant cresyl blue by ascorbic acid. *Turk J Chem* 35: 599-612.
9. Sharma L, Rahman N, Azmi S, Iqbal B, Amburk MI, et al. (2010) UV Spectrophotometric determination of Cu (II) in Synthetic Mixture and Water Samples. *Journal of the Chinese Chemical Society* 57: 622-631.
10. Mouhsen Sh, Hassan K (2012) *Iraqi National Journal of Chemistry* 2: 281-292.
11. Triggler DJ (1996) Ion channels as pharmacologic receptors: the chirality of drug interactions. *Chirality* 8: 35-38.
12. Furukawa M, Katami T, Shibata S (1993) Spectrophotometric determination of copper in aluminium alloys using 2-[2-(4-methylbenzothiazolyl)azo]-5-dimethylaminobenzoic acid. *Journal of Analytical Chemistry* 347: 462-464.
13. Shihad AA (2013) Preparation and Spectrophotometric Study of 2[2-(5-bromo thiazolyl) azo]-4-methoxy phenol.
14. Prodromidis MI, Stalkas CD, Vetsistas PT, Karayannis MI (1994) Spectrophotometric kinetic determination of copper (II) trace amount s based on its catalytic effect on the reduction of the reduced 2,6-dichlorophenolindophenol and hydrogen peroxide. *Talanta* 41: 1645-1649.
15. Texeira LSG, Costa ACS, Ferreira SLC, Carvalho CMS, Freitas ML (1999) Spectrophotometric determination of uranium using 2-[2-thiazolyl) azo]-p-cresol(TAC) in the presence of surfactants. *J Braz Chem Soc* 10: 519.
16. Gusev SI, Zhvakina MV, Kozhevnikov IA (1971) Thiazolylazo dyes and their spectrophotometric applications in analytical chemistry. *Zh Analit Khim* 26: 859.
17. Vogel AI (1953) *Macro and Semimicro Qualitative Inorganic Analysis* 661.
18. Jop (1928) Formation and stability of inorganic complexes in solutions *Ann. Chim* 9: 113.
19. Harvey AE, Manning DL (1950) Methods of establishing empirical formulas of colored complexes in solutions. *J Am Chem Soc* 72: 4488.