

The Effect of Hypothyroidism on Insulin Sensitivity and Their Influence on the Serum Lipid Profile and Renal Function

Abdel-Gayoum AA*

Department of Clinical Laboratory Sciences, Faculty of Applied Medical sciences, University of Hail, Hail, Saudi Arabia

*Corresponding author: Abdel-Gayoum AA, Department of Clinical Laboratory Sciences, Faculty of Applied Medical sciences, University of Hail, Hail, Saudi Arabia, Tel: +966-565714884; E-mail: aabdelgayoum@hotmail.com

Received date: September 06, 2016; Accepted date: October 05, 2016; Published date: October 10, 2016

Copyright: © Abdel-Gayoum AA. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

The overt and subclinical hypothyroidism is more prevalent in patients with diabetes mellitus than in general population. Recently, more studies have been warranted to elucidate the relationship between thyroid hormones disorders and the insulin activity. The present study aims to investigate the correlations between the thyroid hormone levels in subclinical and overt hypothyroid patients with the insulin resistance and their impact on serum lipid profiles and kidney function.

Methods: A total of fifty newly diagnosed hypothyroid patients were recruited for the study and classified into: 1) Subclinical hypothyroid (SH) group: n=26; 2) Overt hypothyroid (OH) group: n=24, and control (C) group: n=18. Fasting blood was collected and serum was used for biochemical analysis.

Results: The fasting serum insulin, glucose levels and the estimated insulin resistance index (HOMA) of the SH and OH groups were significantly ($P<0.001$) elevated compared to control. The regression analysis revealed a significant negative correlation between FT_4 and insulin ($r=-0.32$, $P=0.04$) and significant positive correlations between TSH and insulin ($r=0.57$, $P=0.002$), between TSH and HOMA ($r=0.51$, $P=0.001$), between HOMA and uric acid ($r=0.37$, $P=0.02$), and between TSH and TG ($r=0.47$, $P=0.002$). The serum creatinine, urea and uric acid concentrations were significantly ($P<0.001$) elevated in the OH group but not the SH. The serum total cholesterol, TG and LDL-cholesterol were significantly elevated in both SH and OH.

Conclusion: Hypothyroidism is associated with insulin resistance, renal impairment, hyperurecemia and dyslipidemia, which are atherosclerotic risk indicators. The TSH had maximum impact on the changes.

Subclinical and overt hypothyroid patients with elevated TSH are at high risk of developing atherosclerosis, thus may need close monitor to contain the rise in plasma TSH.

Keywords: Hypothyroidism; Insulin resistance; Renal function; Dyslipidemia

Introduction

Thyroid disorders are common in the general population, with hypothyroidism being the predominant disorder in the adult population [1,2]. The disorder is associated with impaired cardiac contractility, endothelial dysfunction, atherosclerosis and increased carotid intima-media thickness [3,4]. Several studies have shown overt and subclinical hypothyroidism being more prevalent in patients with diabetes mellitus than in general population [5], and women with subclinical hypothyroidism are believed to be at more risk to develop gestational diabetes [6]. The thyroid hormones are known to have a stimulating effect on maturation of the insulin secreting beta cells, and thyroid hormone receptors have been detected in these cells [7]. The hormones enhance gluconeogenesis and glycogenolysis in an opposing effect to insulin [8], whereas, they are known to facilitate the cellular glucose uptake by expressing the glucose transporter-4 (GLT-4) isozyme [9]. Recently researchers have become more concerned to elucidate the complex relationship between thyroid hormones and the insulin activity and more studies have been warranted to clarify the

underlying pathogenic mechanisms [10]. In the light of these reports, we planned to evaluate the correlations between the thyroid function parameters (free T_3 , free T_4 and TSH) in subclinical and overt hypothyroid patients with the insulin resistance and their impact on the serum lipid profiles and kidney function in patients from Hail region, Saudi Arabia.

Patients and Methods

Protocol of study

The study was conducted on newly diagnosed patients with thyroid disorders who were visiting the Diabetic and Endocrinology Out-Patient Unit of King Khalid Hospital, Hail, Saudi Arabia. The patients clinical data entered in the record book of the laboratory were used in this study. A total of 50 newly diagnosed hypothyroid patients were recruited for the study. Patients suffering from chronic liver or kidney diseases were excluded. Depending on their serum TSH, free T_3 (FT_3) and free T_4 (FT_4) levels, the patients were classified into:

- Subclinical hypothyroid (SH) group: n=26 patients (M=9, F=17) with serum TSH>4.0 mU/L and with normal FT_4 and FT_3 levels;

- Overt hypothyroid (OH) group: n= 24 patients (M=10, F=14) with TSH>4.0 mU/L and FT₄<9.0 pmol/L;
- The control (C) group: n= 18, (M=8, F=10) were normal subjects visiting the clinic for routine check-up. All patients had an age range of 18-77 years (mean age was 43.47 ± 11.20 years).

The Inclusion criteria were newly diagnosed subclinical or overt hypothyroid, male and females age ≥ 18 Yrs. Exclusion criteria were cardiovascular diseases, diagnosed diabetes mellitus on insulin or other hypoglycemic medication, chronic liver disease or known kidney disease. A sample of 5 ml venous blood was collected from each patient after an overnight fast and serum was separated and used for the biochemical analysis. The protocol of experiment was explained and the consent was obtained from the participants. The study was approved by the ethical committee, Faculty of Applied Medical Sciences, University of Hail, Hail, Saudi Arabia.

Biochemical assays

The concentrations of serum FT₃, FT₄, TSH and insulin were assayed by Autoanalyzer (ELecsys 2010, Cobas E 411-Mannheim Germany). The serum glucose, urea, creatinine, uric acid, total cholesterol, triglyceride (TG), high density lipoprotein cholesterol (HDL-c) and low density lipoprotein cholesterol (LDL-c) were measured by the automated spectrophotometer, Hitachi-717, utilizing commercial kits supplied by Roche Diagnostic, United Kingdom. The very low density lipoprotein (VLDL) fraction was calculated by subtracting HDL+LDL from total cholesterol. The estimated homeostasis model assessment –Insulin resistance (HOMA-IR) index was calculated by the method described by Matthews et al. [11]. It was calculated by multiplying the fasting serum glucose (mmol/L) by the fasting serum insulin (mU/L) divided by 22.5. The values less than 2.5 were considered normal, and the higher values indicated insulin resistance [12].

Statistical analysis

The presented data are means ± SD. The significance of differences between the means was computed by one way analysis of variance, followed by Multiple Comparison Analysis. Spearman's regression analysis was used to study the significance of correlation among the FT₄, FT₃, TSH, HOMA, insulin, glucose, lipids and kidney function parameters. P value less than 0.05 was considered.

Results

Thyroid function

Table 1 summarizes the mean age and thyroid function parameters in the experimental groups. The highest average age was that of the OH group, whereas, the mean age of SH was not different from that of control group. The serum TSH of the OH group was significantly (P<0.001) higher than that of SH, which was in turn significantly (P<0.001) higher than control. The serum FT₃ values of SH and OH groups were not different from that of control and were within normal range. However, the FT₄ of the OH group was significantly (P<0.001) lower than that of the SH and control groups, which were not different from each other.

The serum insulin levels in the SH and OH groups were significantly increased by 2.11-fold, and 2.42-fold, respectively compared to control (Figure 1). However, the fasting serum glucose concentration was

significantly elevated in the OH group by 39.15% compared to control, and was not different from that of the SH group. Moreover, the estimated HOMA-IR was significantly elevated in the OH and SH groups by 3.65-fold and 2.43-fold, respectively compared to control. As shown in Figure 2, the regression analysis showed a significant negative correlation between FT₄ and insulin (r=-0.32, P=0.04) and a trend of negative correlation between FT₄ and HOMA (r=-0.27, P=0.09). It also showed a significant positive correlation between TSH and serum insulin (r=0.57, P=0.002), and between TSH and HOMA (r =0.51, P=0.001). A significant positive correlation was also shown between HOMA and uric acid (r=0.37, P=0.02), and between TSH and TG (r= 0.47, P=0.002).

Effect on kidney function

Table 2 depicts the kidney function parameters in the experimental groups. The serum creatinine, urea and uric acid concentrations were significantly elevated in the OH group by 2.11-fold, 94.39% and 29.61%, respectively compared to control. However, none of the parameters were significantly raised in the SH group.

Effect on serum total lipids and lipoproteins

As shown in Table 3, the serum total cholesterol was significantly increased in the SH and OH by 27.25% and 62.08%, respectively. Similarly, the serum TG was significantly elevated in the SH and OH by 60.78% and 1.01-fold, respectively compared to control.

On the other hand, the HDL-c showed a trend of decrease in the SH and OH groups, but was not statistically significant. However, the LDL-c fractions of the SH and OH groups were significantly elevated by 29.92% and 86.22%, respectively compared to control. Moreover, the VLDL was significantly elevated in the SH and OH by 59.57% and 100%, respectively.




Figure 1: The serum insulin, glucose and estimated insulin resistance index (HOMA-IR) in subclinical hypothyroid (SH), overt hypothyroid (OH) and control (C) groups. Columns and vertical bars represent means ± SD. [†]P<0.001; a: significantly different from C; b: significantly different from SH.




Figure 2: Straight line plot of the regression analysis for ; (A): the serum FT₄ values with serum fasting insulin levels, (B): the serum FT₄ values with the estimated homeostasis model assessment – Insulin resistance (HOMA-IR) index; (C): The serum TSH values with serum insulin levels; (D): The serum TSH values with HOMA-IR index; (E): HOMA-IR index with serum uric acid levels and (F): The serum TSH values with serum triglyceride (TG) levels in pooled control, subclinical and overt hypothyroid patients.

	C	SH	OH
Gender	M=8 F=10	M=9 F=17	M=10 F=14
Age (Yrs)	33.11 ± 10.18	38.94 ± 8.81	48.00 ± 14.66
TSH (mU/L)	2.49 ± 0.81	7.88 ± 3.46a [‡]	14.89 ± 3.48 a [‡] b [‡]
FT ₄ (pmol/L)	14.10 ± 4.47	15.17 ± 3.16	4.84 ± 0.63 a [‡] b [‡]
FT ₃ (pmol/L)	6.17 ± 3.31	5.29 ± 1.01	7.23 ± 3.02

Presented data are mean ± SD; [‡]P<0.001. a: significantly different from C; b: significantly different from SH

Table 1: The mean ages and thyroid function parameters in subclinical hypothyroid (SH), overt hypothyroid (OH) and control (C) groups.

	C	SH	OH
Creatinine (umol/L)	63.46 ± 13.71	70.56 ± 22.36	134.12 ± 15.46a [‡] b [‡]
Urea (mmol/L)	4.46 ± 0.94	5.53 ± 1.16	8.62 ± 2.01a [‡]

Uric acid (umol/L)	232.83 ± 33.15	249.47 ± 25.36	301.79 ± 26.36a [‡] b [‡]
--------------------	----------------	----------------	---

The presented data are means ± SD; [‡]P<0.001; a: significantly different from C; b: significantly different from SH

Table 2: The kidney function parameters in the subclinical hypothyroid (SH), overt hypothyroid (OH) and control (C) groups.

	C	SH	OH
Total cholesterol (mmol/L)	4.22 ± 0.47	5.37 ± 1.28a ^{**}	6.84 ± 1.25a ^b
TG (mmol/L)	1.02 ± 0.41	1.64 ± 0.55a [‡]	2.06 ± 0.72a ^b
HDL-c (mmol/L)	1.40 ± 0.21	1.35 ± 0.21	1.26 ± 0.17
LDL-c (mmol/L)	2.54 ± 0.54	3.30 ± 1.22a [‡]	4.73 ± 0.99a ^b
VLDL-c (mmol/L)	0.47 ± 0.18	0.75 ± 0.25a [‡]	0.94 ± 0.33a ^b

The presented data are means ± SD; ^{*}P<0.05, ^{**}P<0.01, [‡]P<0.001; a: significantly different from C; b: significantly different from SH

Table 3: The serum lipids and lipoprotein levels in the subclinical hypothyroid (SH), overt hypothyroid (OH) and control (C) groups.

Discussion

The present study revealed that our patients of the overt hypothyroid group were older in age with the percentage of females being higher than males. This is a commonly observed trend in thyroid disorders, where hypothyroidism is shown to be more prevalent among older ages and particularly in women [13]. Although the mean value of FT₄ in the overt hypothyroid group was below normal, but the mean FT₃ value was within the normal range (normal range 3.5-7.8 pmol/L). It is known that about 85% of the circulating T₃ is synthesized in extra-thyroid tissues by deiodination of T₄ where metabolic factors prevailing in these tissues may influence its plasma levels. This probably explains its maintenance within normal range in spite of fall in the levels of FT₄. In our study we addressed the possible influence of the thyroid hormones and TSH on insulin sensitivity. In a cohort study, Gronich et al. [4] indicated that overt and subclinical hypothyroidism were related to increased risk for the development of diabetes mellitus, whereas, treatment with thyroxine replacement therapy alleviated that risk.

In the present study there was a several fold increase in the fasting insulin level of the hypothyroid patients with significant increases in the serum glucose concentrations. The estimated insulin resistance index (HOMA) exhibited a significant increase in both subclinical and overt hypothyroid patients. Our results were in agreement with those of Kapadia et al. [10] who found significantly increased fasting serum insulin level with lower insulin sensitivity in the hypothyroid patients. Our data also indicated a significant negative correlation between FT₄ and serum insulin values (r=-0.32, P=0.04), and a more significant positive correlation between TSH and the serum insulin values (r=0.57, P=0.0002). Similarly, TSH exhibited a highly significant positive correlation with HOMA (r= 0.518, P=0.001). This indicates a strong influence of TSH on the fasting insulin levels and the insulin resistance. Some studies have indicated the effect of TSH on insulin action and that even a subtle increase in plasma TSH levels within the normal range can affect insulin secretion [14,15] and may cause insulin resistance and metabolic syndrome [16,17]. Moreover,

hypothyroid patients are known to experience a decrease in glucose transporters GLUT₄ leading to a reduction of glucose uptake and promoting insulin resistance [18,15]. The relationship between thyroid hormonal status and insulin levels in the pathogenesis of insulin resistance is complex. The higher fasting serum insulin concentrations were believed to develop as a compensation result of the insulin resistance [19]. Many authors accept the concept that a patient suffering from an autoimmune disorder is more prone to be affected by an autoimmune disorder of insulin resistance. Tina et al. [20] observed more prevalence of elevated fasting insulin levels among a group of hypothyroid patients with highly elevated levels of anti-thyroid peroxidase antibodies (above 1000 IU/ml) which might support the concept of autoimmunity role in insulin resistance. Other investigators relate the occurrence of insulin resistance among hypothyroid patients to the high prevalence of obesity and the high fat deposits in this population [21]. All proposed underlying causes of insulin resistance among the hypothyroid patients are commonly present, where the majority of these patients are obese and the circulating abnormal antibodies are normally detected.

The present data exhibited significant impairment of kidney function in the overt hypothyroid group as evidenced by elevations of the serum creatinine, urea and uric acid concentrations. As frequently observed in the diabetic population, there is also high prevalence of hypothyroidism among the kidney disease patients [22,23]. Some authors suggest that hypothyroidism leads to impairment of kidney function through alterations in the renal hemodynamics and tubular structure [24]. However, our data exhibited a significant correlation between HOMA and uric acid but not creatinine. It has been indicated that hyperinsulinemia may contribute to hyperuricemia by blocking uric acid excretion, whereas, hyperuricemia has also been reported to precede the development of hyperinsulinemia. In the experimental animals high plasma uric acid levels were shown to directly induce insulin resistance *in vivo* and *in vitro* by inhibiting insulin receptor substrate-1. The generation of reactive oxygen species is believed to play a key role in hyperuricemia-induced insulin resistance [25]. Hyperuricemia was also shown to inhibit the insulin signaling and induce insulin resistance in the cardiomyocytes, which was suggested to be a mechanism of hyperuricemic-related cardiovascular disease [26].

The present results also revealed significant elevations in serum total and LDL-cholesterol levels as well as the serum TG levels in the overt hypothyroid group. Interestingly, the TSH exhibited a significant positive correlation with TG but not with cholesterol. However, Xu et al. [27] found a significant positive correlation between TSH and the serum total cholesterol levels. Other investigators have reported that hypothyroidism is associated with increased serum total and LDL-c, the small dense lipoprotein (a) and TG levels [28]. Moreover, the plasma total cholesterol and LDL-C were also reported to be elevated in the subclinical hypothyroid patients with elevated atherogenic lipid parameters [1,21]. This was observed particularly in subclinical hypothyroid patients with TSH levels ≥ 10 showing increased risk of cardiovascular disease and mortality [29]. The investigators attributed these changes to decreased clearance of LDL particles from circulation due to reductions in the hepatic LDL receptors, and decreased hepatic cholesterol catabolism by the T3-regulated cholesterol 7- α -hydroxylase enzyme [30]. Moreover hypothyroidism is associated with increased oxidation of LDL particles. Such modification of the LDL particles can impair their receptor-mediated uptake causing accumulation [31]. Such elevated levels of circulating oxidized-LDL in hypothyroidism are believed to increase the vascular systemic and pulmonary resistances

causing pulmonary hypertension [32]. Results of the present study have emphasized the significant impact of TSH to increase the TG and possibly total cholesterol levels independent of the thyroid hormones. It highlights the importance of maintaining an appropriate TSH level in patients susceptible to atherosclerotic diseases.

Conclusion

Hypothyroidism is associated with insulin resistance, renal impairment, hyperuricemia and dyslipidemia, all of which are risk factors for cardiovascular diseases.

TSH showed more influence on these changes independent of the thyroid hormones.

Subclinical and overt hypothyroid patients with elevated TSH are at high risk of developing atherosclerosis, thus may need close monitor to contain the rise in plasma TSH level.

Acknowledgement

The author is most grateful to Mr. Nader A. Alaazmi and Mr. Munif H. Al-Shammari from Clinical laboratory of King Khalid Hospital for their assistance in the biochemical analysis of the patients' blood samples.

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

1. Rizos CV, Elisaf MS, Liberopoulos EN (2011) Effects of thyroid dysfunction on lipid profile. *Open Cardiovasc Med J* 5: 76-84.
2. Wu P (2000) Thyroid disease and diabetes. *Clin Diabetes* 18: 38-40.
3. Rhee CM, Brent GA, Kovesdy CP, Soldin OP, Nguyen D, et al. (2015) Thyroid functional disease: an under-recognized cardiovascular risk factor in kidney disease patients. *Nephrol Dial Transplant* 30: 724-737.
4. Ciccone MM, De Pergola G, Porcelli MT, Scicchitano P, Caldarola P, et al. (2010) Increased carotid IMT in overweight and obese women affected by Hashimoto's thyroiditis: an adiposity and autoimmune linkage? *BMC Cardiovasc. Disord* 10: 22.
5. Gronich N, Deftereos SN, Lavi I, Persidis AS, Abernethy DR, et al. (2015) Hypothyroidism Is a Risk Factor for New-Onset Diabetes: A Cohort Study. *Diabetes Care* 38: 1657-1664.
6. Tudela CM, Casey BM, McIntire DD, Cunningham FG (2012) Relationship of subclinical thyroid disease to the incidence of gestational diabetes. *Obstet Gynecol* 119: 983-988.
7. Zinke A, Schmoll D, Zachmann M, Schmoll J, Junker H, et al. (2003) Expression of thyroid hormone receptor isoform alpha1 in pancreatic islets. *Exp Clin Endocrinol Diabetes* 111: 198-202.
8. Raboudi N, Arem R, Jones RH, Chap Z, Pena J, et al. (1989) Fasting and postabsorptive hepatic glucose and insulin metabolism in hyperthyroidism. *Am J Physiol* 256: E159-E166.
9. Weinstein SP, O'Boyle E, Fisher M, Haber RS. (1994) Regulation of GLUT2 glucose transporter expression in liver by thyroid hormone: Evidence for hormonal regulation of the hepatic glucose transport system. *Endocrinol* 135: 649-654.
10. Kapadia KB, Bhatt PA, Shah JS (2012) Association between altered thyroid state and insulin resistance. *J Pharmacol Pharmacother* 3: 156-160.
11. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, et al. (1985) Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 28: 412-419.

12. Gutch M, Kumar S, Razi SM, Gupta, KK, Gupta A (2015) Assessment of insulin sensitivity/resistance. *Indian J Endocrinol Metab* 19: 160-164.
13. Rehman SU, Cope DW, Senseney AD, Brzezinski W (2005) Thyroid Disorders in Elderly Patients. *South Med J* 98: 543-549.
14. Ortega E, Koska J, Pannacciulli N, Bunt JC, Krakoff J (2008) Free triiodothyronine plasma concentrations are positively associated with insulin secretion in euthyroid individuals. *Eur J Endocrinol* 158: 217-221.
15. Maratou E, Hadjidakis DJ, Kollias A, Tsegka K, Peppas M, et al. (2009) Studies of insulin resistance in patients with clinical and subclinical hypothyroidism. *Eur J Endocrinol* 160: 785-790.
16. Raterman HG, van Eijk IC, Voskuyl AE, Peters MJL, Dijkmans BAC, et al. (2010) The metabolic syndrome is amplified in hypothyroid rheumatoid arthritis patients: a cross sectional study. *Ann Rheum Dis* 69: 39-42.
17. Heima NE, Eekhoff EM, Oosterwerff MM, Lips PT, van Schoor NM, et al. (2013) Thyroid function and metabolic syndrome in older persons: a population based study. *Eur J Endocrinol* 168: 59-65.
18. Shepherd PR, Kahn BB (1999) Glucose transporters and insulin action--implications for insulin resistance and diabetes mellitus. *N Engl J Med* 341: 248-257.
19. Kahn CR (1994) Insulin action, diabetogenesis, and the cause of type II diabetes. *Diabetes* 43: 1066-1084.
20. Tina M, Faranak S, Koorosh K (2014) Insulin resistance in hypothyroid patients under Levothyroxine therapy: a comparison between those with and without thyroid autoimmunity. *J Diabetes Metab Disord* 13: 1-7.
21. Jung CH, Sung KC, Shin HS, Rhee EJ, Lee WY, et al. (2003) Thyroid dysfunction and their relation to cardiovascular risk factors such as lipid profile, hsCRP, and waist hip ratio in Korea. *Korean J Intern Med* 18: 146-153.
22. Bando Y, Ushioji Y, Okafuji K, Toya D, Tanaka N, et al. (2002) Non-autoimmune primary hypothyroidism in diabetic and non-diabetic chronic renal dysfunction. *Exp Clin Endocrinol Diabetes* 110: 408-415.
23. Chonchol M, Lippi G, Salvagno G, Zoppini G, Muggeo M, et al. (2008) Prevalence of subclinical hypothyroidism in patients with chronic kidney disease. *Clin J Am Soc Nephrol* 3: 1296-1300.
24. Mariani LH, Berns JS (2012) The renal manifestations of thyroid disease. *J Am Soc Nephrol* 23: 22-26.
25. Zhu Y, Hu Y, Huang T, Zhang Y, Li Z, et al. (2014) High uric acid directly inhibits insulin signalling and induces insulin resistance. *Biochem Biophys Res Commun* 447: 707-714.
26. Zhi L, Yuzhang Z, Tianliang H, Hisatome I, Yamamoto T, et al. (2016) High Uric Acid Induces Insulin Resistance in Cardiomyocytes In Vitro and In Vivo. *PLoS One* 11: e0147737.
27. Xu C, Yang X, Liu W, Yuan H, Yu C, et al. (2012) Thyroid stimulating hormone, independent of thyroid hormone, can elevate the serum total cholesterol level in patients with coronary heart disease: a cross-sectional design. *Nutr Metab* 9: 54-55.
28. Duntas LH (2002) Thyroid disease and lipids. *Thyroid* 12: 287-293.
29. Rodondi N, den Elzen WP, Bauer DC, Cappola AR, Razvi S, et al. (2010) Subclinical hypothyroidism and the risk of coronary heart disease and mortality. *JAMA* 304: 1365-1374.
30. Drover VA, Agellon LB (2004) Regulation of the human cholesterol 7 α -hydroxylase gene (CYP7A1) by thyroid hormone in transgenic mice. *Endocrinology* 145: 574-581.
31. Ittermann T, Baumeister SE, Völzke H, Wasner C, Schminke UH, et al. (2012) Are serum TSH levels associated with oxidized low-density lipoprotein? Results from the Study of Health in Pomerania. *Clin Endocrinol (Oxf)* 76: 526-532.
32. Scicchitano P, Dentamaro I, Tunzi F, Ricci G, Carbonara S, et al. (2016) Pulmonary hypertension in thyroid diseases. *Endocrine* pp: 1-10.