

The Effect of Maize Germ on the Presence of Aflatoxins in Corn Flours Treated with a Thermo-Alkaline Process

Lineth J Vega-Rojas^{1,3}, Magda Carvajal-Moreno^{2*}, Isela Rojas-Molina³, Francisco Rojo-Callejas⁴, Silvia Ruiz-Velasco⁵ and Mario E Rodríguez-García⁶

¹Posgrado en Ciencias Químico Biológicas, Facultad de Química, Universidad Autónoma de Querétaro, Centro Universitario C.P-76010 Querétaro, Qro, México

²Departamento de Botánica, Instituto de Biología, Universidad Nacional Autónoma de México (UNAM), Ciudad Universitaria, 04510 Ciudad de México, México

³Laboratorio de Investigación Química y Farmacológica de Productos Naturales, Facultad de Química, Universidad Autónoma de Querétaro, Cerro de las Campanas S/N, C.P. 76010 Querétaro, Qro., México.

⁴Departamento de Química Analítica, Facultad de Química, Universidad Nacional Autónoma de México, 04510 Ciudad de México, México

⁵Departamento de Probabilidad y Estadística, Instituto de Investigaciones en Matemáticas Aplicadas y en Sistemas, UNAM, 04510 Ciudad de México.

⁶Departamento de Nanotecnología, Centro de Física Aplicada y Tecnología Avanzada, Universidad Nacional Autónoma de México, Querétaro, Qro, CP 76230, México

Abstract

Background: Aflatoxins are important and frequent teratogens, mutagens and carcinogens of maize, and the germ of the maize seed has linoleic acid, which can control aflatoxins. An aflatoxin analysis of maize flours with and without germ can show the role of the germ in the plant control of these toxins.

Methods: The samples were cooked with the traditional nixtamalization process with varying calcium hydroxide content (from 0 to 2.1 w/w of corn) and steeping times of 0 and 9 h. The aflatoxin purification was performed with immunoaffinity columns, and the quantification was performed using HPLC.

Results: It was found that the presence of the germ and the concentrations of 1.4 and 2.1% w/w of Ca(OH)₂ had significant effects ($p \leq 0.05$) on the decrease in the AFB₁ and AFG₁ content.

Conclusion: The linoleic acid of the germ inhibited AFB₁ and AFG₁. However, the aflatoxin content of the experimental samples was higher than 12 µg kg⁻¹, which is the tolerance limit permitted by NOM-247-SSA1-2008.

Keywords: Aflatoxin; Corn; Germ; Nixtamalization

Abbreviations

CAN: Acetonitrile; AFB₁: Aflatoxin B₁; AFB₂: Aflatoxin B₂; AFG₁: Aflatoxin G₁; AFG₂: Aflatoxin G₂; AFS: Aflatoxins; ANOVA: Analysis of variance; Ca(OH)₂: Calcium hydroxide; CaO: Calcium oxide; HCl: Hydrogen Chloride; pH: Concentration of hydrogen ions in units of moles per liter; R²: Correlation coefficient; DNA: Deoxyribonucleic acid; H₂O_d: Distilled water; ≤: Equal or less than; ≥: Equal or more than; EC: European Community; g: Gram; HPLC: High Performance Liquid Chromatography; h: Hours; kg: Kilogram; LOD: Limit of detection; LOQ: Limit of quantification; w/w: Mass/mass; MeOH: Methanol; µg kg⁻¹: Micrograms per kilogram; µL: Microliters; mL(s): Milliliter(s); mL/min: Milliliters per minute; mm: Millimeter; min: Minutes; ng g⁻¹: Nanograms per gram; ng mL⁻¹: Nanograms per milliliter; nm: Nanometers; PBS: Phosphate buffer saline; p: Probability; Rec %: Recovery percentage; RT: Retention Time; rpm: Revolutions per minute; NaCl: Sodium Chloride; NaOH: Sodium hydroxide; TFA: Trifluoroacetic Acid; UV-Vis: Ultraviolet-Visual

Introduction

Maize grains, in their different cooking forms, are the staple food in the diet of the Mexican population, with an average per capita intake of 342 g/day [1]. In Mexico and some countries in Central and North America, most corn-based products are subjected to a thermo-alkaline process called nixtamalization [2]. Most of the maize designated for the miller-tortilla industry and flour is frequently contaminated by insects, mites, bacteria, and fungi, which attack the grain in the field and in storage [3]. Aflatoxins (AFs) are secondary metabolites that chemically correspond to the bis-dihydro-furanocoumarins produced mainly by the fungi *Aspergillus flavus*, *A. parasiticus* and *A. nomius*, which are associated with the stress of the plant in the development stage and stress associated with drought, competition with weeds, poor

fertility, and insect pests [4]. The contamination and propagation of the fungi occur during tillage, growth and especially in the post-harvest period [4,5]. Aflatoxins are the most frequently ingested mutagens in food, and the International Agency for Research on Cancer (IARC), in 2002, classified them as Group I carcinogens for humans [6]. The fungal colonization of the grain starts in the tip cap continues in the embryo and finishes in the endosperm. The fungal spores enter the maize flower instead of a pollen grain, and contamination occurs in the maize grain from the inside [7]. *A. flavus* and *A. parasiticus* are capable of invading seeds, are pathogenic to insects, and grow as saprophytes on crop debris in the field and soil. The sclerotia of *A. flavus* can be another source of primary inoculum, which forms in damaged and intact maize kernels and may overwinter in kernels dispersed onto the soil [7]. Insect wounds facilitate the entrance of the fungi directly to the pericarp and endosperm, facilitating AF contamination, although these tissues are not preferred. The presence of *A. flavus* in the germ was detected after 72 h of inoculation with the fungus [8,9]. The colonization of the grain with *A. flavus* and *A. parasiticus* starts in the tissues rich in oil, from the embryo to the endosperm [10]. The germ of

*Corresponding author: Magda Carvajal-Moreno, Departamento de Botánica, Instituto de Biología, Universidad Nacional Autónoma de México (UNAM), Ciudad Universitaria, 04510 Ciudad de México, México, Tel: +(5255) 5622 9138; Fax: +(5255) 5550 1760; E-mail: prashantagarwal2@gmail.com

Received August 01, 2016; Accepted August 29, 2016; Published September 09, 2016

Citation: Vega-Rojas LJ, Carvajal-Moreno M, Rojas-Molina I, Rojo-Callejas F, Ruiz-Velasco S, et al. (2016) The Effect of Maize Germ on the Presence of Aflatoxins in Corn Flours Treated with a Thermo-Alkaline Process. J Microb Biochem Technol 8: 408-414. doi: 10.4172/1948-5948.1000317

Copyright: © 2016 Vega-Rojas LJ, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

maize is the reproductive part that germinates to grow into a plant [11]; it is the embryo of the seed. A survey of the AFs in maize tortillas from shops and supermarkets in Mexico City was carried out and 13% of the samples had higher amounts of AFs than the tolerance limit allowed by Mexican legislation ($12 \mu\text{g kg}^{-1}$) [12]. However, the analysis of other nixtamalized products such as flours has not been carried out.

Maize germ oil [13] may be used as a source for oil extraction or used directly as a food ingredient. The germ is retained as an integral part of whole-grain foods. However, in countries such as Colombia and Venezuela, the grain of maize used to make arepas, empanadas and snacks is fractioned by a mechanical process that intentionally removes the pericarp and the major part of the germ; this product is known as threshed maize. The flours obtained from threshed maize have lower fiber and fat contents and are therefore enriched to improve their nutritional value [14]. *Milling with the removal of the cereal germ is the method most used today to ensure better preservation* [15].

Non-whole grain methods of milling are intended to isolate the endosperm, which is ground into flour, with the removal of both the husk (bran) and the germ. The removal of the bran is aimed at producing flour with a white rather than a brown color and eliminating fiber; neither of these objectives is desirable from a nutritional perspective.

The germ is rich in polyunsaturated fats, primarily omega-6 fatty acids such as linoleic acid, which have a tendency to oxidize and become rancid in storage. The germ removal therefore improves the storage qualities of the flour. Linoleic acid is an essential fatty acid that must be consumed for proper health. During the thermo-alkaline processing of the corn grain with 1 to 3% CaO, 97% of the AFs in the dough are degraded and in the case of tortillas, they are apparently reduced by 98% [5,16].

The aim of the present study is to evaluate the effect of the presence of germ on the aflatoxin content (AFB_1 , AFB_2 , AFG_1 and AFG_2) in nixtamalized corn flours obtained by the traditional method with different concentrations of calcium hydroxide and steeping times.

Materials and Methods

Sample preparation

Ten kg of corn kernels harvested in 2014 (Puma hybrid-Monsanto) used for tortilla making in Mexico were chosen for this experiment. Three hundred grams of corn kernels and 600 mL of distilled water were cooked at 92°C with different $\text{Ca}(\text{OH})_2$ (Fermont, Monterrey, NL, Mexico) concentrations (0, 0.7, 1.4, and 2.1 w/w) with 3 or 9 h of steeping [17].

Two samples, each of 300 g, were lime treated or “nixtamalized” at the same time and with the same process conditions. After cooking and steeping, the germ of one of these samples was manually removed, while the germ was left on in the second sample. The corn samples with and without germ was ground into corn dough and then dehydrated in a furnace (40°C for 8 h). The samples were re-milled using a 0.8-mm hammer mill (Pulvex 200, México D.F. México). Finally, different types of corn flour were obtained.

Aflatoxin extraction from corn flours

Steeping times of 3 and 9 h were chosen for this analysis because these times correspond to the industrial and traditional lime treatment (“nixtamalization”) processes, respectively. Three different concentrations of $\text{Ca}(\text{OH})_2$ (0.7, 1.4 and 2.1) were used [5] to obtain nixtamalized corn flour with and without germ and raw corn flour and cooked corn flours with and without germ.

The AF extraction consisted of blending 25 g of flour with 100 mL of a mixture of MeOH: H_2O_d (80:20 v/v) and 1 g NaCl to clarify. Later, the blended mixture was centrifuged (ALC 4235 Working Cool System, USA) at 4300 rpm for 10 min and the supernatant retained. Four mL of the supernatant was diluted with 16 mL of PBS pH 7.4, and this dilution was passed through an immunoaffinity column (Easi-Extract Aflatoxin R-Biopharm Rhone Ltd., Glasgow, Scotland, UK) previously balanced with 20 mL of PBS [18,19] for the detection of total AF (AFt). After receiving the sample filtrate, the immunoaffinity column was washed with 20 mL of H_2O_d at a flux of 5 mL per min.

The immunoaffinity column was dried by passing air through the agarose gel, and it was eluted with 1.5 mL of HPLC grade MeOH by gravity at one drop per second. Afterwards, 1.5 mL of H_2O_d was applied with refluxing to denature the gel with AFt antibodies and release the pure AF into the 3 mL eluate. Finally, the eluate was dried at 40°C and derivatized [20,21].

Aflatoxin (AF) standards

The AF standards AFB_1 , AFB_2 , AFG_1 and AFG_2 (Sigma-Aldrich, St. Louis, MO, USA) were dissolved in a benzene:acetonitrile (ACN) mixture (98:2 v/v) and stored in amber vials at 4°C to prepare 1000 ng/mL solutions. A UV-Vis spectrophotometer (Genesys 10 UV-Vis, Thermo Electron Corporation, Madison, Wisconsin, USA) was calibrated for the correction factor (Method 971.22B, AOAC 2006) [22]. The absorbance in the spectrophotometer was adjusted to 0 with a blank of HPLC methanol in a quartz cell, and the wavelength of maximum absorbance (360-362 nm) for the four AFs was determined [22].

A stock solution of 1000 ng/mL of each of AFB_1 , AFB_2 , AFG_1 and AFG_2 was used to make 15 dilutions to determine the linearity (calibration curves) of the validation assay method.

Derivatization

The dry AF standards were resuspended in 200 μL acetonitrile (ACN) and 800 μL of derivatizing solution to increase the fluorescence. The derivatizing solution consisted of 5 mL of trifluoroacetic acid (TFA) (Sigma-Aldrich, St. Louis, MO, USA), 2.5 mL of glacial acetic acid (Merck, Naucalpan, Edo Mex, México) and 17.5 mL of deionized water to obtain a final concentration of 20% TFA (v/v). The mixture was vortexed (Vortex G-560, Bohemia, NY, EEUU) for 30 s and the vials were placed in a water bath (Equipos de Laboratorio BG Mod. BM 40T, Mexico) at 65°C for 10 min [20,21]. Finally, 60 μL of the derivatized AFs were injected into the HPLC for chemical quantification.

AF quantification by HPLC

The standard AF HPLC analysis [23] and the eluted samples were quantified with an Agilent Technologies HPLC (Series 1200) with an isocratic pump (G1310A Series DE62957044), fluorescence detector (G1321A Series DE60456380) and autosampler (G1329A Series DE64761666), a chromatographic Agilent Eclipse XDS-C18 column (4.6×250 mm, 5 μm of particle size) and the HPLC program ChemStation 32. The analytical conditions were a $\text{H}_2\text{O}/\text{ACN}/\text{MeOH}$ (65:15:20 v/v/v) mobile phase, 60 μL injection volume per triplicate, 1 mL/min fluid speed for 20 min, excitation at 362 nm and emission at 425 nm for AFB_1 and AFB_2 and 450 nm for AFG_1 and AFG_2 .

Validation of the HPLC quantification method for AFs in corn flour

The validation requirements for the HPLC quantification method for AFs in corn flour have been previously developed [24,25].

Selectivity

This assay was performed to determine the degree to which the method can determine the analyte (AF) without matrix interference. A mixture of the 4 AF standards was analyzed (blank), as were the two matrixes (corn flour with germ and corn flour without germ) independently enriched with the four AFs. The blank and the two corn flour sources with AFs were extracted, concentrated and purified with immunoaffinity total anti-aflatoxin columns. They were then eluted and derivatized, and the resulting chromatograms were compared.

Linearity

This assay was performed to obtain results proportional to the analyte (AF) concentration. Fifteen standard dilutions (0.01, 0.05, 0.5, 1, 2, 4, 5, 8, 10, 16, 30, 32, 64, 100, 128 ng mL⁻¹) of each of the four AFs were tested separately to obtain the calibration curves.

Limits of detection (LOD) and quantification (LOQ)

The LOD was determined by observing the minimal concentration that displayed a chromatographic peak signal on the HPLC equipment. The LOQ was calculated as the LOD multiplied by five, taking into account the sample results.

Recovery percentage (R %)

The recovery percentage is the efficiency of the method in detecting all of the analyte (AFs) present in a sample. Each matrix of cooked corn flour with or without germ was fortified with 100 ng g⁻¹ of each one of the four AFs independently before the analysis. Later, the extraction and derivatization were performed, and the recovery percentage was used for the adjustment of the AF concentrations in the samples.

Statistical analysis

Three factors of variation were considered: the presence of the germ, the concentration of Ca(OH)₂ and the steeping time. A three-way ANOVA and a post hoc Bonferroni procedure for multiple comparisons with a confidence index of 95% were calculated. The tolerance limit for AFt in maize tortillas according to the Mexican Official Rule [26] is 12 µg kg⁻¹.

Results and Discussion

Validation of the method to quantify AF selectivity

The chromatograms after the AF addition to the three matrices showed that the elution order of the analytes and the retention times (RT) were not modified by the different matrices and the four AF analytes did not overlap.

Linearity, LOD and LOQ

The LOD and the coefficient of correlation (R²) of the AF calibration curves for the 4 AFs were AFB₁ (LOD=0.1 ng mL⁻¹, R² = 0.9973); AFB₂ (LOD=0.01 ng mL⁻¹, R²=0.9908); AFG₁ (LOD=0.01 ng mL⁻¹, R₂=0.9969); and AFG₂ (LOD=0.5 ng mL⁻¹ R²=0.9908). The LOD of the equipment was highly efficient and facilitated accurate results.

The RTs were adjusted according to the selectivity and linearity for the two treatments of corn flour with and without germ, and the ranges were for AFB₁, RT=7.195 to 8.983 min; for AFB₂, RT=18.965 to 23.272 min; for AFG₁, RT=7.681 to 9.541 min; and for AFG₂, RT=12.214 to 14.492 min.

Recovery percentages

The recovery percentages were for AFB₁, 91%; for AFB₂, 88%; for AFG₁, 90%; and for AFG₂, 90%.

AF quantification of corn with and without germ samples by HPLC

The AFt contents of the nixtamalized corn flour with germ and the nixtamalized corn flours without germ were higher than 12 µg kg⁻¹, which is the tolerance limit established in NOM-247-SSA1-2008 [26] Figures 1-5.

The germ, which contains linoleic acid, diminishes the amount AFB₁ (Figure 1) and AFG₁ (Figure 3) but does not control AFB₂ (Figure 2) or AFG₂ (Figure 4). The deduction is that the effect of the linoleic acid is on the double bond of the first furan ring, which is the place where AFB₁ attaches to the DNA and this place is related to the mutagenic and carcinogenic effects of AFB₁ and AFG₁. The linoleic acid of the germ had no effect on AFB₂ or AFG₂ because neither have a double bond in the first furan. The effect of the linoleic acid of the germ on the opening of the lactone ring would produce a loss of fluorescence, but in all cases, the AFs were detected due to the presence of their fluorescence. The




Figure 1: Effect of the steeping time and different Ca(OH)₂ concentrations on the amount of Aflatoxin B₁ of corn flour with and without germ.




Figure 2: Effect of the steeping time and different Ca(OH)₂ concentrations on the amount of Aflatoxin B₂ in corn flour with and without germ.

conclusion is that the effect of the linoleic acid of the germ is on the first furan ring and not on the lactone ring. In fact, there were only traces <1 ng of AFG₂, which was more related to the steeping time than to the presence of the germ with linoleic acid (Figure 4). It is very important to note that AFG₂ and AFB₂ have to be converted to AFG₁ and AFB₁ to recover their mutagenicity.

Of all the AFs, aflatoxin G₁ had the highest concentrations (>40 ng) in the corn flour without germ for 1.4% Ca(OH)₂ and a 9 h steeping time, and the linoleic acid of the germ diminishes the AF content (0.54 ng) (Figure 3). It is important to recall that AFG₁ had the highest concentration in all the studied samples. It is important to emphasize that the intake of maize contaminated with AFs for a prolonged time can cause accumulation in the DNA and contribute to AFB₁-DNA adduct formation that can trigger mutation and possibly the development of cancer. The toxicity of the AFs according to their properties of being teratogenic, mutagenic and carcinogenic follow the decreasing order of AFB₁>AFG₁>AFG₂>AFB₂ [27-29].




Figure 3: Effect of the steeping time and different Ca(OH)₂ concentrations on the amount of Aflatoxin G₁ in corn flour with and without germ.




Figure 4: Effect of the steeping time and different Ca(OH)₂ concentrations on the amount of Aflatoxin G₂ in corn flour with and without germ.




Figure 5: Effect of total AF quantity on the different types of aflatoxin in corn flour with and without germ (5A) and the effect of the steeping time with different Ca(OH)₂ concentrations (5B) on the total amount of aflatoxins in corn flour with and without germ.

The AF groups G and B differ in the enzymes involved in their biosynthesis, although they have the same intermediate product; the enzymes implicated in the formation of the AFs of the G group are less stable than those of the B group [30]. The changes in the pH conditions determine the type of metabolites synthesized. Therefore, the AFs of group B are favored at acid pH values below 6, while the AFs of group G are stimulated with neutral to basic pH values above 6 [31]. The different species of fungi produce different types of AFs; *Aspergillus flavus* produces only AFB₁ and AFB₂ and *Aspergillus parasiticus* produces AFB₁, AFB₂, AFG₁ and AFG₂. Therefore, we can deduce that the maize samples were contaminated with both fungi, although they were likely exposed for a longer time to *A. parasiticus*. The biological factors such as the exposure to different species of *Aspergillus*, the physical factors such as the humidity and substrate and the biochemical and environmental factors favor AF production. In the same way, inadequate storage conditions such as a temperature between 20°C and 40°C, humidity between 10 and 20% and relative moisture in the air between 70°C to 90°C increases the AF levels [32,33].

The thermo-alkaline treatment increases the pH of the products and triggers the hydrolysis of the lactone ring of the AFs, forming salts soluble in water. These AF products are lost in the washings, and they can accumulate in the residual water called “nejayote” [5]. The removal of the remaining water could be an alternative for eliminating AFs. Nevertheless, the conditions used in this nixtamalization process did not reduce the levels of aflatoxins B₁, B₂, G₁ or G₂ in the crude, cooking and nixtamalized flours with germ, and there were no significant differences in the conditions. Other studies [34] demonstrated that the AFs zearalenone and deoxynivalenol resist alkaline treatment with a concentration of 2% w/w Ca (OH)₂ and a cooking temperature of 110°C.

The concentrations of AFB₁ and AFG₁ in the corn flours without germ were higher than those in the corn flours with germ in the thermo-alkaline process (Figures 1 and 3). This result shows that compounds in the germ, mainly linoleic acid, exert a protective effect against AFs. However, an opposite effect was observed for AFB₂ for all the conditions tested (Figure 2), where the AFs are found mainly in the corn germ, and they accumulate in the section richest in fatty acids [8].

The AFt content in the corn flours with germ was significantly lower ($p \leq 0.05$) in comparison to that in the flours without germ (Figure 5A). This demonstrates that the germ inhibits the AFs due to its high content of fatty acids, mainly linoleic acid, and other components such as flavonoids, monoterpenes and carotenoids that have the capacity of suppressing the metabolic activation of the AFs, specifically AFB₁ and AFG₁.

The Bonferroni test showed that the presence of the germ and the 1.4 and 2.1% w/w concentrations of Ca(OH)₂ in the nixtamalization process had a significant effect ($p \leq 0.05$) on the contents of AFB₁, AFB₂, AFG₁ and AFt. Uniquely, the steeping time had a significant effect on the contents of AFB₂ ($p=0.0458$) and AFG₁ ($p=0.0445$). The content of AFG₂ is not significantly affected by the three factors analyzed (presence of germ, concentration of Ca(OH)₂ and steeping time).

In some cases, the boiling water in the alkaline process can reduce the detected AF levels in the maize grain from 127 $\mu\text{g kg}^{-1}$ to 68.6 $\mu\text{g kg}^{-1}$ in tortillas [35]. The AF reduction was between 20 to 46% and the residue concentration was higher than the tolerance limit (20 $\mu\text{g kg}^{-1}$).

The AF levels appear to be reduced due to the alkaline pH of the nixtamalization process that opens the lactone ring. The acid pH of the gastric fluid in the stomach during digestion closes the lactone ring and facilitates the reactivation of the AFs in the neutral pH of the pancreatic fluid [36]. A study analyzed the role of pH, temperature, and the frying effect in the AF reduction in tortillas and maize snacks, where only the AFs of the samples treated in pH 9.5 were activated again [5]. The increase in AFs by acidification was much lower, ranging from 0 to 18% higher than the reported values [35]. In alkaline solutions, a slow hydrolysis of the AF lactone ring can be observed. This hydrolysis is reversible, with the reformation of the lactone ring by the environmental acidification [5,35]. In a study of the levels of AFs in peanut beverages artificially contaminated with AFB₁, the role of pH in the inactivation of the AFs was observed at temperatures higher than 100°C. The effects of the pH (5, 8 and 10.2), temperature (121, 130 and 140°C) and heating time (between 5 and 20 s and 15 min) did not significantly ($p < 0.001$) reduce the mutagenic activity. Heating the beverage at 130°C for 20 s or 121°C for 15 min at pH 10.2, however, reduced the mutagenic activity by $79.4 \pm 5.3\%$ and $86.6 \pm 5.2\%$, respectively [37]. This decrease in the mutagenicity is attributed to the transformation of AFB₁ to AFD₁, which is 450 times less mutagenic. The lactone ring hydrolyzes when NaOH is added to adjust the pH. The reduction of the mutagenic effect is attributed to the partial hydration of the lactone ring in the presence of the chloride added to adjust the pH. The AFB₁ is transformed to AFB_{2a}, which is 1000 times less mutagenic. In peanut beverages, the change of pH from 8 to 5 or 10.2 was not significant for the reduction of the AF mutagenicity if there is no heating [37].

The maize grain has developed protection mechanisms against AFs through the synthesis of lipids, flavonoids, and monoterpenes, among other components [32]. Research on corn grains contaminated with AFs showed that linoleic acid (the major fatty acid of the lipid content of the germ) has the capacity to suppress the metabolic activity *in vitro* of AFB₁ [27,38]. The effect of the lipids of the germ on nixtamalized corn flours has not been reported.

The role of linoleic acid is not well known, there are some reports about an antioxidant effect. Linoleic do reduce the aflatoxin content or its fluorescent detection, but it inactivates the mutagenic effect of aflatoxins. Most of the carcinogens are activated by oxidation. The maize plant cannot avoid the presence of the *Aspergillus flavus* or *A. parasiticus* fungi, but it can inhibit its mutagenic effects by avoiding the oxidation of the first furan of the aflatoxin that has the double bond, with the linoleic acid in the germ. The aflatoxin B₁ (Figure 1), AFG₁ (Figure 3) and the hydroxylates AFM₁, AFP₁ and aflatoxicol have a double bond in the first furan where the oxygen links and make the 8,9 exo-epoxide that links to the DNA and can produce a mutagenic effect. In Figures 1 and 3 we can appreciate that the germ (with linoleic and oleic acids) inhibits the aflatoxin B₁ and G₁ production, but the germ (with linoleic and oleic acids) have no effect on Aflatoxin B₂ (Figure 2) and Aflatoxin G₂ (Figure 4) because they have no double bond in the first furan and have to be bio transformed to AFB₁ and AFG₁ respectively to be carcinogenic, so the double bond of the first furan might be the place where the linoleic acid has its effect. Around 90% of the carcinogens are mutagens. Therefore many cereals such as maize, sorghum, rice etc., spices (like pepper and chili peppers), and oilseed plants such as nuts, pistachios, cottonseeds, peanuts, etc. have many antioxidants raised by the presence of the fungi, to inhibit its mutagenicity, but they have also lots of aflatoxins.

It is like a war, the plant produces antioxidants to avoid the oxidation of the first furan of the aflatoxins B₁ and G₁, that is the one that links to N⁷ of the guanine DNA, mainly guanine, to form the adduct (active carcinogen), but these seeds with antioxidants such as linoleic acid in maize are full of inactive and not detectable aflatoxins, because many times the coumarin section of the aflatoxin has the lactone opened and lacks fluorescence. In the case of peanut the antioxidant is resveratrol. It is a dialectic (union of opposites) situation: maize has aflatoxins inactivated by linoleic acid, but 17% of them can be stored in the human DNA during many years. Aflatoxins are “unavoidable toxins” because humans normally eat cereals, spices and oilseeds to live, but these aflatoxins are dangerous because they can produce aflatoxin B₁-formamide pyrimidine (AFB₁-FAPY) adducts that is the active carcinogens. The problem is that the physicians recommend the ingestion of oilseeds to cardiac or gastric disease persons, because they have antioxidants, but they are not aware that they also have “inactivated aflatoxins” that in the animal or human livers can be metabolized and reactivated again.

The AFB₁ mutagenicity could be tested with spiked maize tortilla samples using the Ames test but not with naturally contaminated corn, showing the resistance or control of the maize plant against the AFB₁ mutagenicity [36]. The oils from the maize germ can help to reduce the effect of the AFs.

The conclusion of the present work is that the linoleic acid of the germ inhibited AFB₁ and AFG₁, and its mechanism of action possibly involved the first furan ring and not the lactone ring. There were only traces <1 ng of AFG₂.

The AFt in all studied samples was higher than 12 $\mu\text{g kg}^{-1}$, which is the maximum tolerance limit permitted by NOM-247-SSA1-2008. However, 1.4 and 2.1% w/w concentrations of Ca(OH)₂ and the presence of the germ have significant effects ($p \leq 0.05$) in decreasing the content of AFB₁ and AFG₁.

Acknowledgement

- This work was supported by:
- Consejo Nacional de Ciencia y Tecnología (CONACYT), (“Fondo

Mixto de Fomento a la Investigación Científica y Tecnológica CONACYT- Gobierno del Estado de Querétaro”, Grant No. M0016-2014-03, Project No. QRO-2014-C03-250289 and “Fondo para el Fortalecimiento de la Investigación UAQ-2015” (FOFI-UAQ-2015, Project No. FCQ-2016-16), México.

- CONACYT, for the PhD scholarship given to Lineth J. Vega-Rojas.
- Instituto de Biología, Universidad Nacional Autónoma de México (IBUNAM) for the laboratory facilities, technology, equipment and materials used in this work.
- The University of Querétaro, Mexico for its postgraduate program.

The authors also thank the IBUNAM personnel: Noemí Chávez, from the Technical Secretary (Secretaría Técnica); Joel Villavicencio, Jorge López, Alfredo Wong, and Julio César Montero for computer assistance; and Georgina Ortega Leite and Gerardo Arévalo for library information.

References

1. FAO, Food and Agriculture Organization (2009) La FAO en México. Más de 60 años de cooperación. Cap.3. En: Nutrición y seguridad alimentaria. Agroanálisis AC. Roma, Italia.
2. Rodríguez-García EM, Serna-Saldivar SO, Sánchez-Sinencio F (2008) Cambios nutrimentales en el maíz inducidos por el proceso de nixtamalización. Cap.1. En: Nixtamalización del maíz a la tortilla Aspectos nutrimentales y toxicológicos. Universidad Autónoma de Querétaro, México.
3. Vázquez-Durán A, Díaz-Torres R, Ramírez-Noguera P, Moreno-Martínez E, Méndez-Albores A (2014) Cytotoxic and genotoxic evaluation of tortillas produced by microwave heating during alkaline-cooking of aflatoxin-contaminated maize. *Food Sci* 5: 1024-1029.
4. Marin S, Ramos AJ, Cano-Sancho G, Sanchis V (2013) Mycotoxins: Occurrence, toxicology and exposure assessment. *Food Chem Toxicol* 60: 218-237.
5. Torres P, Guzmán-Ortiz M, Ramírez-Wong B (2001) Revising the role of pH and thermal treatments in aflatoxin content reduction during the tortilla and deep frying processes. *J Agric Food Chem* 49: 2825-2829.
6. IARC International Agency for the Research on Cancer (2012) Chemical agents and related occupations: A review of human carcinogens. *IARC Monogr Eval Carcinog Risks Hum* 100: 225-248.
7. Diener UL, Davis ND (1987) Biology of *Aspergillus flavus* and *A. parasiticus*. In: Zuber MS, Lillehoj EB, Renfro BL eds. Aflatoxin in Maize: A Proceedings of the Workshop. CIMMYT. Mexico DF, pp: 33-40.
8. Payne GA (1983) Epidemiology of aflatoxin formation by *A. flavus*. In: Aflatoxin and *Aspergillus flavus* in Corn. Southern. Diener UL, Asquith RL, Dickens JW Eds. Auburn University: Opelika, AL. USA.
9. Dolezal AL, O'Brien GR, Nielsen DM, Woloshuk CP, Boston RS et al. (2013) Localization, morphology and transcriptional profile of *Aspergillus flavus* during seed colonization. *Mol Plant Pathol* 14: 898-909.
10. Keller NP, Butchko R, Sarr B, Phillips TD (1994) A visual pattern of mycotoxin production in maize kernels by *Aspergillus* spp. *Phytopathology* 84: 483-488.
11. Black MJ, Bewley JD, Halmer P (2006) The Encyclopedia of Seeds - Science, Technology and Uses. CAB International. ISBN 9780851997230.
12. Castillo-Urueta P, Carvajal M, Méndez I, Meza F, Gálvez A (2011) Survey of aflatoxins in maize tortillas from Mexico City. *Food Addit Contam Part B* 4: 42-51.
13. Stellman JM (1998) Encyclopaedia of Occupational Health and Safety. United Nations International Labour Office.
14. Pineda-Gómez P, Acosta-Osorio AA, Coral DF, Rosales-Rivera A, Sánchez Echeverri LA, et al. (2012) Physicochemical characterization of traditional and commercial instant corn flours prepared with threshed white corn. *CyTA J Food* doi:10.1080/19476337.2011.65369.
15. Hui YH, Chandan RC, Clark S, Cross N, Dobbs JC, et al. (2007) Handbook of Food Products Manufacturing: Principles, Bakery, Beverages, Cereals, Cheese, Confectionary, Fats, Fruits and Functional Foods. Wiley-Interscience.
16. De Arriola MC, De Porres E, De Cabrera S, De Zepeda M, Rolz C (1988) Aflatoxins fate during alkaline cooking of corn for tortilla preparation. *J Agric Food Chem* 36: 530-533.
17. Gutiérrez E, Rojas MI, Pons HJ, Guzmán H, Aguas AB, et al. (2007) Study of calcium ion diffusion in nixtamalized quality protein maize as a function of cooking temperature. *Cereal Chem* 84: 186-194.
18. Groopman JD, Donahue KF (1988) Aflatoxin, a human carcinogen: determination in foods and biological samples by monoclonal antibody affinity chromatography. *J Assoc Off Anal Chem* 71: 861-867.
19. Garner RC, Whattam MM, Taylor P.J.L., Stow MW (1993) Analysis of United Kingdom purchased spices for aflatoxins using an immunoaffinity column clean-up procedure followed by high-performance liquid chromatography. *J Chromatogr A* 648: 485-490.
20. Akiyama H, Goda Y, Tanaka T, Toyoda M (2001) Determination of aflatoxins B₁, B₂, G₁ and G₂ in spices using a multifunctional column clean-up. *J Chromatogr A* 932: 153-157.
21. Kok WT. 1994. Derivatization reactions for the determination of aflatoxins by liquid chromatography with fluorescence detection. *J Chromatogr B Biomed Appl* 659: 127-137.
22. Trucksess MW (2006) Natural toxins. In Official Methods of Analysis of AOAC International (18th Edn), Gaithersburg (MD): AOAC International.
23. Jaimez J, Fente CA, Vazquez BI, Franco CM, Cepeda A, et al. (2000) Application of the assay of aflatoxins by liquid chromatography with fluorescence detection in food analysis. *J Chromatogr A* 882: 1-10.
24. IRAM 32 [Norma del Instituto Argentino de Normalización y Certificación] (1997) Quantifying uncertainty in analytical measurement. *Metrología. Vocabulario VIM. Guide Eurachem- Citac Guide*: 2000.
25. García MA, Soberón E, Cortés M, Rodríguez R, Herrera JL, et al (2002) Guía de validación de métodos analíticos, Colegio Nacional de QFB, México DF, México
26. NOM, Norma Oficial Mexicana NOM-247-SSA1-2008. Diario Oficial de la federación (2008). Productos y servicios. Cereales y sus productos. Cereales, harinas de cereales, sémolas o semolinas. Alimentos a base de: cereales, semillas comestibles, de harinas, sémolas o semolinas o sus mezclas. Productos de panificación. Disposiciones y especificaciones sanitarias y nutrimentales. Métodos de prueba. México.
27. Burgos-Hernández A, López-García R, Njapau H, Park DL (2001a) Partial chemical/structural elucidation of anti-mutagenic compounds from corn. *Toxicology* 166: 161-170.
28. Hsieh DPH (1983) Metabolism and transmission of mycotoxins. In: Proceedings of the International Symposium on Mycotoxins. National Research Center, Cairo, Egypt.
29. Carvajal M, Espinosa-Aguirre JJ, Moctezuma MG, Gonsebatt ME, Rojo F et al. (2004) Minimal amount of aflatoxin B1 to produce a mutation in the Ames test with *Salmonella typhimurium* TA 98. *Rev Mex Micol* 19: 71-79.
30. Yabe K, Nakamura M, Hamasaki T (1999) Enzymatic formation of G-group aflatoxins and biosynthetic relationships between G and B group aflatoxins. *Appl Environ Microbiol* 65: 3867-3872.
31. Buchanan RL Jr, Ayres JC (1975) Effect of initial pH on aflatoxin production. *Appl Microbiol* 30: 1050-1051.
32. Burgos-Hernández A, López-García R, Njapau H, Park DL (2001b) Anti-mutagenic compounds from corn. *Food Addit Contam* 9: 797-809.
33. Cotty PJ, García RJ (2007) Influences of climate on aflatoxins producing fungi and aflatoxins contamination. *Int J Food Microbiol* 119: 109-115.
34. Abbas HK, Mirocha CJ, Rosiles R, Carvajal M (1988) Decomposition of zearalenone and deoxynivalenol in the process of making tortillas from corn. *Cereal Chem* 65: 15-19.
35. Price RL, Jorgensen KV (1985) Effects of processing on aflatoxins levels and on mutagenic potential of tortillas made from naturally contaminated corn. *J Food Sci* 50: 347-349.
36. Moctezuma-Zárate MG, Carvajal M, Espinosa-Aguirre JJ, Gonsebatt-Bonaparte ME, Rojo-Callejas F, et al. (2015) Role of pH in the mutagenicity of Aflatoxin B1 in maize tortillas during *in vitro* human digestion model. *J Food Nutr Disord* 4: 1-10.

37. Rustom IYS, López-Leiva MH, Nair BM (1993) Effect of pH and heat treatment on the mutagenic activity of peanut beverage contaminated with aflatoxin. *Food Chemistry* 46: 37-42.

38. Weber EJ (1987) Lipids of the kernel. In: *Corn: Chemistry and Technology*, American Association of Cereal Chemists, St. Paul, MN. USA.

Citation: Vega-Rojas LJ, Carvajal-Moreno M, Rojas-Molina I, Rojo-Callejas F, Ruiz-Velasco S, et al. (2016) The Effect of Maize Germ on the Presence of Aflatoxins in Corn Flours Treated with a Thermo-Alkaline Process. *J Microb Biochem Technol* 8: 408-414. doi: [10.4172/1948-5948.1000317](https://doi.org/10.4172/1948-5948.1000317)

OMICS International: Open Access Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid peer review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled for better prominence and citations
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Best discounts for your subsequent articles

Submit your manuscript at: <http://www.editorialmanager.com/jmbt>