

The First Molecular Detection of Caprine Arthritis Encephalitis Virus (Caev) in Iran

Gholam Ali Kojouri^{1*}, Mojdeh Emami¹ and Hasan Momtaz²

¹Department of Clinical Sciences, School of Veterinary Medicine, Shahrekord University, Iran

²Department of Pathobiology, School of Veterinary Medicine, Shahrekord Azad University, Iran

Abstract

Caprine arthritis encephalitis virus (CAEV) is a retrovirus that naturally infects goats and induces chronic inflammatory and degenerative disease. Infection of goats has a worldwide distribution but it has not been previously recognized by molecular method in Iran. For this reason, 95 blood samples were taken from 4 months to 3 years goats to determine naturally infected animals with caprine arthritis-encephalitis virus (CAEV) by using polymerase chain reaction (PCR) method. The PCR primers were designed to amplify a 433 base pair region of the proviral gag gene. Results showed that 15.7 percent of sampled goats (15 cases) were positive for CAEV. The present study was confirmed the presence of CAEV genome in goats flock of Iran for the first time.

Keywords: Goat; Arthritis; Encephalitis; PCR; Gag Gene

Introduction

Caprine arthritis-encephalitis virus (CAEV) belongs to the lentivirus subfamily of retroviruses [1] which causes chronic progressive arthritis, pneumonia, and mastitis in adult goats and leukoencephalomyelitis in young kids [2,3]. Disease is widespread in populations of domestic sheep and goats throughout the world [4] and its high prevalence especially in industrialized countries (80-95% in the breeding stocks) is a major concern [5,6]. Ingestion of virus-infected colostrums or goat milk [7] is the main way of disease transmission and other routes such as direct contact, bodily secretions, and excretions [8,9] are less abundant. Prevalence of CAE around the world has passed over a wide range of variation among different countries, which may be as low as 1.9% in Turkey and 3.6% in Mexico, or even up to as high as 73% in USA and 82% in Australia [10]. Recent study from Iran showed that the 10.87% of goats were serologically infected for CAEV [11] but unfortunately there is no previous data about the prevalence of this disease by molecular methods in Iran.

To our knowledge, most infected goats are asymptomatic and so far there is no effective treatment and supportive drugs are usually expensive. On the other hand, previous study suggested the high reduced production and significant economic losses due to disease [12]. Therefore, using a rapid, fast and sensitive method is essential for its diagnosis. Recent studies showed the high sensitivity, specificity and accuracy of molecular methods like Polymerase Chain Reaction (PCR) for rapid detection of CAEV in clinical samples [13,14]. Therefore, this current study was carried out for detection of CAEV in slaughtered goats by using PCR method in Iran.

Material and Methods

Animals

Ninety five slaughtered goats were sampled during the experiment. Goats were a member of different flocks and were randomly chosen among slaughter. Flocks were located in different geographic areas of the Chahar Mahal va Bakhtiyari province of Iran.

Samples

After giving an identification number, blood samples were taken and age, sex, and origin of animals were recorded. Additionally, based on dentition and owner's information, the age was determined. Blood

samples were taken from jugular vein and mononuclear cells (PBMC) were isolated from 10 ml EDTA-blood on a Ficoll-Hypaque gradient. They were then destroyed in buffer (100 mM Tris-HCl pH 7.5; 12.5mM EDTA-Na₂; 150 mM NaCl; 0.5% SDS) containing Proteinase K (Sigma) (50 µg per 10⁷ PBMC) at 55° for 2 h. Genomic DNA was subsequently extracted twice with phenol-chloroform-isoamyl alcohol (25:24:1, v/v/v, Sigma). It was then again extracted twice with chloroform-isoamyl alcohol (24:1, v/v, Sigma). Following treatment with 96% ethanol (Sigma) and centrifugation, the precipitate was re-suspended in distilled water (40 µl) and stored at 4°C till the day of the PCR test [15].

Gag gene-PCR

Primers used to amplify the group associated antigen (gag) gene were those described by Clavijo and Thorsen [16], their sequence as follows:

Forward: 5' AGGAGGAGGATTAACAGTGG-3'

Reverse: 5' TCCTGGCCTTAATGCTTG-3'

PCR amplification

The PCR reaction was carried out in primus 96 thermal cycler. DNA fragment were amplified in 25 µL reaction mixtures containing approximately 3 µL of genomic DNA, 1.5 mM MgCl₂, 0.2 mM each Deoxynucleoside Triphosphates (dNTPs), 400 Nm of each primer and 2.5U Taq DNA polymerase and 5% DMSO. The PCR amplification was achieved by 35 cycles each including a denaturation step at 94°C for 1 min, annealing step at 50°C for 30 sec and extension step at 72°C for 1 min. PCR amplified DNA fragments were analyzed by electrophoresis on 1.5% agarose gels stained with ethidium bromide and visualized on a UV trans illuminator [15].

***Corresponding author:** Gholam Ali Kojouri, Department of Clinical Sciences, School of Veterinary Medicine, Shahrekord University, Iran, Tel/Fax: +98 381 4424427; E-mail: drgholam_alikojouri@yahoo.com; kojouri@vet.sku.ac.ir

Received March 29, 2014; Accepted July 25, 2014; Published July 28, 2014

Citation: Kojouri GA, Emami M, Momtaz H (2014) The First Molecular Detection of Caprine Arthritis Encephalitis Virus (Caev) in Iran. J Veterinar Sci Technol 5: 184. doi:10.4172/2157-7579.1000184

Copyright: © 2014 Kojouri GA, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Figure 1: Agarose gel electrophoresis and ethidium bromide staining. Lane M indicates standard DNA marker; 1, negative control; 2, positive control; lanes 3 and 5 positive samples; lanes 4, 6 and 7 negative samples; the 433-bp PCR product has also been indicated.

Results

A total of 95 whole blood samples were taken from apparent healthy goats between June and August 2013. The goats were of various mixed breeds, and their ages ranged from 4 months to 3 years. Peripheral blood mononuclear cells (PBMC) were separated using the Ficoll gradient centrifugation method and total DNA was extracted from PBMC for confirmation by polymerase chain reaction. PCR products were analyzed by gel electrophoresis and ethidium bromide staining. A total of 15 out of 95 crossbred goats had a single amplification band of the expected (433 bp) size (Figure 1). It is also showed that there was no significant difference between male and female goats PCR results using Q-square test ($P > 0.05$). Based on these findings, infection rate among Shahrekord district goats were 15.7%. From 95 samples, 6 of 35 (17.14%) and 9 of 60 (15%) samples that were taken from males and females, respectively, were positive.

Discussion

Our study showed that the PCR method is good compromise for CAEV detection. PCR amplification had been described for detection of CAEV DNA in cultured cell using primers for the pol gene [17] and in PBMC and milk from infected goats using primer for the gag gene [16,18,19,]. DNA from PBMCs has used as templates, with reported sensitivities ranging from 70-95% [14,18]. The PCR assay in the present study showed an overall prevalence of (15.7%). This is in agreement with Karanikolaou et al. [20]. For the first time, our finding has confirmed the presence of CAEV genome in PBMC of apparently healthy Iranian goats.

CAEV is a lentivirus that persistently infects goats and sheep. The finding that CAEV and Maedi-Visna viruses frequently cross the species barrier between goats and sheep, and vice versa, has changed our view of the epidemiology of these viruses that are now referred to as small ruminant lentiviruses (SRLV) [15]. CAEV is transmitted from infected mothers to their offspring, mainly via ingestion of infected colostrum and milk. CAEV induces overt pathology in about one third of the infected animals. The frequency of affected animals varies in different goat families, pointing to an important genetic component in this disease. The principal manifestations are encephalitis and interstitial pneumonia in young animals, whereas arthritis and mastitis predominate in adult goats [21].

Our results showed that the prevalence of CAEV in goats in Iran was 15.7% which was higher than Mexico (3.6%) [22], Italy (4.0%) [23] and Great Britain (10.3%) [24] but entirely lower than USA (73%) [25] and Norway (86%) [26]. Serological study in Hungary showed that 30% of the Hungarian goat population was infected with CAEV [27]. Previous study from Thailand showed that the prevalence of CAEV was 12.4% [28]. Another study from Brazil showed that the total prevalence of CAEV in goats is 8.2% which was lower than our study. In total, the prevalence of CAEV around the world has passed over a wide range of variation which may be as low as 3.6% in Mexico and 15.7% in Iran (our study) until 82% in Australia [10]. Studies indicated that contact with goats from other herds, purchased animals into herd, presence of sheep in the farms and finally herd size are the main risk factors for epidemiology and prevalence of CAEV [29,30].

Comparison between the PCR and an antibody ELISA indicated very strong agreement between the two assays ($\kappa = 0.912$) and it was recognized that four of 7 goats with indeterminate ELISA results were PCR-positive [31]. Unfortunately, studies about the prevalence of CAEV by PCR method are very low throughout the world.

To our knowledge, our study is the first prevalence report of direct detection of CAEV from slaughtered goats by using PCR in Iran. Our results revealed the uses of molecular methods especially PCR for rapid and accurate detection of this virus in clinical samples. The results of this study showed that 17.14% of males and 15% of female goats were positive for presence of CAEV in Iran.

References

1. Al-Ani FK, Vestweber JG (1984) Caprine arthritis-encephalitis syndrome (CAE): a review. *Vet Res Commun* 8: 243-253.
2. Cheevers WP, Knowles DP, McGuire TC, Cunningham DR, Adams DS, et al. (1988) Chronic disease in goats orally infected with two isolates of the caprine arthritis-encephalitis lentivirus. *Lab. Invest*, 58: 510-517.
3. Cork LC, Hadlow WJ, Crawford TB, Gorham JR, Piper RC (1974) Infectious leukoencephalomyelitis of young goats. *J Infect Dis* 129: 134-141.
4. Gufler H, Gasteiner J, Lombardo D, Stifter E, Krassnig R, et al. (2007) Serological study of small ruminant Lentivirus in goats in Italy. *Small Rumin. Res.*, 73: 169-173.
5. Rowe JD, East NE (1997) Risk factors for transmission and methods for control of caprine arthritis-encephalitis virus infection. *Vet Clin North Am Food Anim Pract* 13: 35-53.
6. Pérez G, Bugnard F, Calavas D (1994) Study of a prevention programme for caprine arthritis-encephalitis. *Vet Res* 25: 322-326.
7. De Maar TW, Blumer ES, Sherman DM (1995) Failure of horizontal transmission of caprine arthritis encephalitis virus to non-dairy breeds of goats. *Small Rumin. Res*, 17: 197-198.
8. East NE, Rowe JD, Dahlberg JE, Theilen GH, Pederson NC (1993) Modes of transmission of caprine arthritis-encephalitis virus infection. *Small Rumin. Res.*, 10: 251-262.
9. Leitner G, Krifucks O, Weisblit L, Lavi Y, Bernstein S, et al. (2010) The effect of caprine arthritis encephalitis virus infection on production in goats. *Vet J* 183: 328-331.
10. Al-Qudah K, Al-Majali AM, Ismail ZW (2006) Epidemiological studies on caprine arthritis-encephalitis virus infection in Jordan. *Small Rumin. Res*, 66: 181-186.
11. Sakhaee E, Khalili M, Imani M (2011) Sero-prevalence survey of caprine arthritis-encephalitis virus in Iran. *Online J. Vet. Res*, 15: 256-260.
12. Schoborg RV (2002) Analysis of caprine arthritis encephalitis virus (CAEV) temporal gene expression in infected cells. *Virus Res* 90: 37-46.
13. Al Ahmad MZA, Chebloune Y, Bouzar BA, Baril G, Bouvier F, et al. (2008) Lack of risk of transmission of caprine arthritis-encephalitis virus (CAEV) after an appropriate embryo transfer procedure. *Theriogenology*, 69: 408-415.
14. Eltahir YM, Dovas CI, Papanastassopoulou M, Koumbati M, Giadinis N, et al.

- (2006) Development of a semi-nested PCR using degenerate primers for the generic detection of small ruminant lentivirus proviral DNA. *J Virol Methods* 135: 240-246.
15. Elfahal AM, Zakia AM, El-Hussien AM (2010) First Report of Caprine Arthritis Encephalitis Virus Infection in Sudan. *J. Anim. Vet. Advan*, 9: 736-740.
16. Clavijo A, Thorsen J (1996) Application of polymerase chain reaction for the diagnosis of caprine arthritis-encephalitis. *Small Rumin. Res*, 22: 69-77.
17. Zanoni R, Pauli U, Peterhans E (1990) Detection of caprine arthritis-encephalitis- and maedi-visna viruses using the polymerase chain reaction. *Experientia* 46: 316-319.
18. Reddy PG, Sapp WJ, Heneine W (1993) Detection of caprine arthritis-encephalitis virus by polymerase chain reaction. *J Clin Microbiol* 31: 3042-3043.
19. Rimstad E, East NE, Torten M, Higgins J, DeRock E, et al. (1993) Delayed seroconversion following naturally acquired caprine arthritis-encephalitis virus infection in goats. *Am J Vet Res* 54: 1858-1862.
20. Karanikolaou K, Angelopoulou K, Papanastasiopoulou M, Koumpati-Artopiou M, Papadopoulos O, et al. (2005) Detection of small ruminant lentiviruses by PCR and serology tests in field samples of animals from Greece. *Small Rumin. Res*, 58: 181-187.
21. Park JE, Son SY, Shin HJ (2010) Sequence comparison on gag gene of caprine arthritis encephalitis virus from Korea. *Virus Genes* 41: 99-101.
22. Torres-Acosta JFJ, Gutierrez-Ruiz EJ, Butler V, Schmidt A, Evans J, et al. (2003) Serological survey of caprine arthritis-encephalitis virus in 83 goat herds of Yucatan, Mexico. *Small Rumin Res*, 49: 207-211.
23. Gufler H, Baumgartner W (2007) Overview of herd and CAEV status in dwarf goats in South Tyrol, Italy. *Vet Q* 29: 68-70.
24. Dawson M, Wilesmith JW (1985) Serological survey of lentivirus (maedi-visna/caprine arthritis-encephalitis) infection in British goat herds. *Vet Rec* 117: 86-89.
25. Cutlip RC, Lehmkuhl HD, Sacks JM, Weaver AL (1992) Prevalence of antibody to caprine arthritis-encephalitis virus in goats in the United States. *J Am Vet Med Assoc* 200: 802-805.
26. Nord K, Rimstad E, Storset AK, Loken T (1998) Prevalence of antibodies against caprine arthritis-encephalitis virus in goat herds in Norway. *Small Rumin. Res*, 28: 115-121.
27. Kusza SZ, Bosze ZS, Kukovics S, Javor A (2004) Genetic assay of Caprine arthritis encephalitis in the Hungarian goat herd. *South Africa J. Anim. Sci*, 34: 13-16.
28. Ratanapob N, Rukkamsuk T, Jala S (2009) Seroprevalence of caprine arthritis encephalitis virus infection in goats raised in the central part and western part of Thailand. *Proceed. 47th Kasetsart University Annual Conference*, 62-69.
29. Bandeira DA, de Castro RS, Azevedo EO, de Souza Seixas Melo L, de Melo CB (2009) Seroprevalence of caprine arthritis-encephalitis virus in goats in the Cariri region, Paraiba state, Brazil. *Vet J* 180: 399-401.
30. Lin TN, Ngarmkum S, Oraveerakul K, Virakul P, Techakumphu M (2011) Seroprevalence and risk factors associated with caprine arthritis-encephalitis virus infection in goats in the western part of Thailand. *Thai. J. Vet. Med*, 41: 353-360.
31. Barlough J, East N, Rowe JD, Van Hoosear K, DeRock E, et al. (1994) Double-nested polymerase chain reaction for detection of caprine arthritis-encephalitis virus proviral DNA in blood, milk, and tissues of infected goats. *J. Virol. Methods*, 50: 101-113.

Citation: Kojouri GA, Emami M, Momtaz H (2014) The First Molecular Detection of Caprine Arthritis Encephalitis Virus (Caev) in Iran. *J Veterinar Sci Technol* 5: 184. doi:[10.4172/2157-7579.1000184](https://doi.org/10.4172/2157-7579.1000184)

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 350 Open Access Journals
- 30,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, DOAJ, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: www.editorialmanager.com/lifesciences