

The Generalization of the Stalling's Theorem

Onsory A* and Araskhan M*

Department of Mathematics, Yazd Branch, Islamic Azad University, Yazd, Iran

Abstract

In this paper, we present a relative version of the concept of lower marginal series and give some isomorphisms among $\mathcal{V}G$ -marginal factor groups. Also, we conclude a generalized version of the Stalling's theorem. Finally, we present a sufficient condition under which the order of the generalized Baer-invariant of a pair of finite groups divides the order of the generalized Baer-invariant of its factor groups.

Keywords: Schur-Baer variety; Pair of groups; $\mathcal{V}G$ -marginal series

Introduction

There exists a long history of interaction between Schur multipliers and other mathematical concepts. This basic notion started by Schur [1], when he introduced multipliers in order to study projective representations of groups. It was known later that the Schur multiplier had a relation with homology and cohomology of groups. In fact, if G is a finite group, then $M(G) \cong H^2(G, \mathbb{C}^*) \cong H_2(G, \mathbb{Z})$, where $M(G)$ is the Schur multiplier of G , $H^2(G, \mathbb{C}^*)$ is the second cohomology of G with coefficient in \mathbb{C}^* and $H_2(G, \mathbb{Z})$ is the second internal homology of G [2]. Hopf [3] proved that $M(G) \cong (R \cap F^2) / [R, F]$. He also proved that the Schur multiplier of G is independent of the free presentation of G . Let (G, N) be a pair of groups, where N is a normal subgroup in Ellis [4] defined the Schur multiplier of the pair (G, N) to be the abelian group $M(G, N)$ appears in the following natural exact sequence

$$H_3(G) \rightarrow H_3(G, N) \rightarrow M(G, N) \rightarrow M(G) \rightarrow M(G/N) \\ \rightarrow G/[N, G] \rightarrow (G)_{ab} \rightarrow (G/N)_{ab} \rightarrow 1,$$

where $H_3(-)$ denote the third homology of a group with integer coefficients. He also proved that if the normal subgroup N possess a complement in G , then for each free presentation $1 \rightarrow R \rightarrow F \rightarrow G \rightarrow 1$ of G , $M(G, N)$ is isomorphic with the factor group $(R \cap [S, F]) / [R, F]$, where S is a normal subgroup of F such that $S/R \cong N$. In particular, if $N=G$ then the Schur multiplier of (G, N) will be $M(G) = (R \cap [F, F]) / [R, F]$.

We assume that the reader is familiar with the notions of the verbal subgroup $V(G)$, and the marginal subgroup

$V^*(G)$, associated with a variety of groups \mathcal{V} and a group G [5] for more information on varieties of groups). Let F_∞ be the free group freely generated by the countable set $X = \{x_1, x_2, \dots\}$ and \mathcal{V} and \mathcal{W} be two varieties of groups defined by the sets of laws \mathcal{V} and \mathcal{W} , respectively. Let N be a normal subgroup of a group G , then we define $[NV^*G]$ to be the subgroup of G generated by the elements of the following set:

$$\{v(g_1, g_2, \dots, g_r) v(g_1, g_2, \dots, g_r)^{-1} \mid 1 \leq i \leq r, v \in V, g_1, \dots, g_r \in G, n \in N\}.$$

It is easily checked that $[NV^*G]$ is the least normal subgroup T of G such that N/T is contained in $V^*(G/T)$ [6].

The first to create the generalization of the Schur multiplier to any variety of groups was Baer [7]. It is well known fact that the recent concept is useful in classifying groups into isologism classes. Leedham-Green and McKay [8] introduced the following generalized version of the Baer-invariant of a group with respect to two varieties \mathcal{V} and \mathcal{W} .

Let G be an arbitrary group in \mathcal{W} with a free presentation $1 \rightarrow R \rightarrow F \rightarrow G \rightarrow 1$, in which F is a free group. Clearly,

$1 = W(G) = W(F)R/R$ and hence $W(F) \subseteq R$, therefore,

$$1 \rightarrow R/W(F) \rightarrow F/W(F) \rightarrow G \rightarrow 1$$

is a \mathcal{W} -free presentation of the group G . We call

$$\mathcal{W}\mathcal{V}M(G) = \frac{R/W(F) \cap V(F/W(F))}{[R/W(F)V^*(F/W(F))]} = \frac{W(F)(R \cap V(F))}{W(F)[RV^*F]}$$

the *generalized Baer-invariant* of the group G in \mathcal{W} with respect to the variety \mathcal{V} . Now if N is a normal subgroup of the group G for a suitable normal subgroup S of the free group F , we have $N \cong S/R$. Then we can define the generalized Baer-invariant of the pair of groups with respect to two varieties \mathcal{V} and \mathcal{W} as follows:

$$\mathcal{W}\mathcal{V}M(G, N) = \frac{R/W(F) \cap [S/W(F)V^*(F/W(F))]}{[R/W(F)V^*(F/W(F))]} = \frac{W(F)(R \cap [SV^*F])}{W(F)[RV^*F]}.$$

One may check that $\mathcal{W}\mathcal{V}M(G, N)$ is always abelian and independent of the free presentation of G . In particular, if \mathcal{W} is the variety of all groups and $N=G$ then the generalized Baer-invariant of the pair (G, N) will be

$$\mathcal{V}M(G, G) = \frac{R \cap V(F)}{[RV^*F]} = \mathcal{V}M(G),$$

which is the usual Baer-invariant of G with respect to \mathcal{V} [8].

It is interesting to know the connection between the Baer-invariant of a pair of finite groups (G, N) and its factor groups with respect to the Schur-Baer variety \mathcal{V} . In the next section, we show that under some circumstances there are some isomorphisms among \mathcal{V}_G -marginal factor groups (Theorem 2.2). Also, a sufficient condition will be given such that the order of the generalized Baer-invariant of a pair of finite groups divides the order of the generalized Baer-invariant of the pair of its factor groups (Theorem 2.5).

Variety \mathcal{V} is called a *Schur-Baer* variety if for any group G in which

***Corresponding authors:** Onsory A, Department of Mathematics, Yazd Branch, Islamic Azad University, Yazd, Iran, Tel: 982144865179; E-mail: aliasghar-onsory@yahoo.com

Araskhan M, Department of Mathematics, Yazd Branch, Islamic Azad University, Yazd, Iran, Tel: 982144865179; E-mail: m.araskhan@iauyazd.ac.ir

Received November 28, 2015; **Accepted** December 21, 2015; **Published** December 23, 2015

Citation: Onsory A, Araskhan M (2015) The Generalization of the Stalling's Theorem. J Generalized Lie Theory Appl S2: 003. doi:10.4172/1736-4337.S2-003

Copyright: © 2015 Onsory A, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

the marginal factor group $G / V(G)$ is finite, then the verbal subgroup $V(G)$ is also finite. Schur [9] proved that the variety of abelian groups is a Schur-Baer variety and Baer [10] showed that a variety defined by outer commutator words carries this property. In 2002, Moghaddam et al. [11] proved that for a finite group G , $\mathcal{VM}(G)$ is finite with respect to a Schur-Baer variety \mathcal{V} . In the following lemma we prove similar result for the $\mathcal{WVM}(G, N)$ and $\mathcal{WVM}(G)$ with using another technique.

Lemma 1.1. *Let \mathcal{V} be a Schur-Baer variety and G be a finite group in \mathcal{W} with a normal subgroup N . Then there exists a group H with a normal subgroup K such that*

$$|[NV^*G]| \mathcal{WVM}(G, N) = |[KV^*H]| < \infty.$$

In particular, $|V(G)| \mathcal{WVM}(G) = |V(H)| < \infty$.

Proof. Let $G = F / R$ be a free presentation for the group G and S be a normal subgroup of the free group F such that $N \cong S / R$, then

$$\frac{R}{W(F)[RV^*F]} \subseteq V^* \left(\frac{F}{W(F)[RV^*F]} \right).$$

Let $H = F / W(F)[RV^*F]$ and $K = S / W(F)[RV^*F]$, then $| \frac{H}{V^*(H)} | < |G| < \infty$ and $|[KV^*H]| \leq |V(H)| < \infty$. But

$$|[KV^*H]| = \left| \frac{W(F)[SV^*F]}{W(F)[RV^*F]} \right| = \left| \frac{W(F)[SV^*F]}{W(F)(R \cap [SV^*F])} \right| \left| \frac{W(F)(R \cap [SV^*F])}{W(F)[RV^*F]} \right|.$$

Also, $[NV^*G] = \frac{[SV^*F]R}{R} = \frac{W(F)[SV^*F]R}{R} \cong \frac{W(F)[SV^*F]}{W(F)(R \cap [SV^*F])}$. Thus the result holds.

Stallings' Theorem

In the following lemma we present some exact sequences for the generalized Baer-invariant of a pair of groups and its factor groups.

Lemma 2.1. *Let G be a group with a free presentation $1 \rightarrow R \rightarrow F \rightarrow G \rightarrow 1$ and S, T be normal subgroups of the free group F such that $T \subseteq S, S / R \cong N$ and $T / R \cong K$. Then the following sequences are exact:*

$$(i) 1 \rightarrow \frac{W(F)(R \cap [TV^*F])}{W(F)[RV^*F]} \rightarrow \mathcal{WVM}(G, N) \rightarrow \mathcal{WVM}(G / K, N / K) \rightarrow \frac{K \cap [NV^*G]}{[KV^*G]} \rightarrow 1;$$

$$(ii) \mathcal{WVM}(G, N) \rightarrow \mathcal{WVM}(G / K, N / K) \rightarrow \frac{K}{[KV^*G]} \rightarrow \frac{N}{[NV^*G]} \rightarrow \frac{N}{[NV^*G]K} \rightarrow 1;$$

(iii) Moreover, if K is contained in $V(G)$, then the following sequence is exact:

$$1 \rightarrow \frac{R \cap [SV^*F]}{W(F)[TV^*F] \cap [SV^*F]} \rightarrow \mathcal{WVM}(G / K, N / K) \rightarrow K \rightarrow \frac{N}{[NV^*G]} \rightarrow \frac{N}{[NV^*G]K} \rightarrow 1.$$

Proof. Considering the definition mentioned above we can conclude:

$$\mathcal{WVM}(G / K, N / K) = \frac{W(F)(T \cap [SV^*F])}{W(F)[TV^*F]} \quad \frac{K \cap [NV^*G]}{[KV^*G]} = \frac{(T \cap [SV^*F])R}{[TV^*F]R},$$

$$\mathcal{WVM}(G, N) = \frac{W(F)(R \cap [SV^*F])}{W(F)[RV^*F]}.$$

Now one can easily check that the sequences (i) and (ii) are exact.

(iii) Using the assumption, we have $W(F)[TV^*F] \subseteq R$. Therefore, one can easily check that the following sequence is exact:

$$1 \rightarrow \frac{R \cap [SV^*F]}{W(F)[TV^*F] \cap [SV^*F]} \rightarrow \frac{W(F)(T \cap [SV^*F])}{W(F)[TV^*F]} \rightarrow T / R \rightarrow \frac{S}{[SV^*F]R} \rightarrow \frac{S}{[SV^*F]T} \rightarrow 1.$$

Let N be a normal subgroup of a group G . Then we define a series of normal subgroups of N as follows:

$$N = V_0(N, G) \supseteq V_1(N, G) \supseteq V_2(N, G) \supseteq \dots \supseteq V_n(N, G) \supseteq \dots,$$

where $V_i(N, G) = [V_{i-1}(N, G)V^*G]$ for all $n \geq 1$. We call such a series the lower \mathcal{V}_G -marginal series of N in G . One may also define the upper \mathcal{V}_G -marginal series as in studies of Moghaddam et al. [11].

We say that the normal subgroup N of G is \mathcal{V}_G -nilpotent if it has a finite lower \mathcal{V}_G -marginal series. The shortest length of such series is called the class of \mathcal{V}_G -nilpotency of N in G . If $N = G$, then this is called lower \mathcal{V} -marginal series of G . The group G is said to be \mathcal{V} -nilpotent iff $V_n(G) = 1$, for some positive integer n [12].

Now, we want to show that under some circumstances there are some isomorphisms among \mathcal{V}_G -marginal factor groups. By using Lemma 2.1, we have the following Theorem, which generalizes 7.9.1 of literature of Hilton and Stammbach [13].

Theorem 2.2. *Let $f: G \rightarrow H$ be a group homomorphism and N be a normal subgroup of G and K be a normal subgroup of H such that $f(N) \subseteq K$. Suppose f induces isomorphisms $f_0: G / N \rightarrow H / K$ and $\bar{f}_1: N / [NV^*G] \rightarrow K / [KV^*H]$, and that $f_*: \mathcal{WVM}(G, N) \rightarrow \mathcal{WVM}(H, K)$ is an epimorphism. Then f induces isomorphisms $f_n: G / V_n(N, G) \xrightarrow{\cong} H / V_n(K, H)$ and $\bar{f}_n: N / V_n(N, G) \xrightarrow{\cong} K / V_n(K, H)$ for all $n \geq 0$.*

Proof. At first, we want to mention a point that for making it easier to draw the following diagrams, we would like to introduce $P_n = V_n(N, G)$ and $Q_n = V_n(K, H)$. We proceed by induction. For $n = 0$ the assertion is trivial. For $n = 1$, consider the following diagram:

$$\begin{array}{ccccccc} 1 & \longrightarrow & N / [NV^*G] & \longrightarrow & G / [NV^*G] & \longrightarrow & G / N & \longrightarrow & 1 \\ & & \downarrow \bar{f}_1 & & \downarrow f_1 & & \downarrow f_0 & & \\ 1 & \longrightarrow & K / [KV^*H] & \longrightarrow & H / [KV^*H] & \longrightarrow & H / K & \longrightarrow & 1. \end{array}$$

By the hypothesis \bar{f}_1 and f_0 are isomorphism, hence f_1 is an isomorphism. Assume that $n \geq 2$. By considering Lemma 2.1(ii), we can conclude the following commutative diagram:

$$\begin{array}{ccccccccccc} WVM(G, N) & \rightarrow & WVM(G/P_{n-1}, N/P_{n-1}) & \rightarrow & P_{n-1}/P_n & \rightarrow & N/[NV^*G] & \rightarrow & N/[NV^*G]P_{n-1} & \rightarrow & 1 \\ \downarrow \alpha_1 & & \downarrow \alpha_2 & & \downarrow \alpha_3 & & \downarrow \alpha_4 & & \downarrow \alpha_5 & & \\ WVM(H, K) & \rightarrow & WVM(H/Q_{n-1}, K/Q_{n-1}) & \rightarrow & Q_{n-1}/Q_n & \rightarrow & K/[KV^*H] & \rightarrow & K/[KV^*H]Q_{n-1} & \rightarrow & 1 \end{array}$$

Note that the naturality of the map f induces homomorphisms $\alpha_i, i = 1, 2, \dots, 5$ such that (*) is commutative. By hypothesis α_1 is an epimorphism and α_4, α_5 are isomorphisms. Also, by considering the induction hypothesis and definition of the Baer-invariant of the pair of groups, α_2 is an isomorphism. Hence by five lemma of Rotman's studies [14] α_3 is an isomorphism. Now consider the following diagram and in the same way, f_n is an isomorphism.

Now we obtain the following corollary.

$$\begin{array}{ccccccc}
 1 & \longrightarrow & P_{n-1}/P_n & \longrightarrow & N/P_n & \longrightarrow & N/P_{n-1} & \longrightarrow & 1 \\
 & & \downarrow \alpha_3 & & \downarrow \bar{f}_n & & \downarrow \bar{f}_{n-1} & & \\
 1 & \longrightarrow & Q_{n-1}/Q_n & \longrightarrow & K/Q_n & \longrightarrow & K/Q_{n-1} & \longrightarrow & 1
 \end{array}$$

By the above discussion α_3 is an isomorphism and by induction of hypothesis \bar{f}_{n-1} is an isomorphism, therefore, \bar{f}_n is an isomorphism. Finally, by the following diagram:

$$\begin{array}{ccccccc}
 1 & \longrightarrow & N/P_n & \longrightarrow & G/P_n & \longrightarrow & GN & \longrightarrow & 1 \\
 & & \downarrow \bar{f}_n & & \downarrow f_n & & \downarrow f_1 & & \\
 1 & \longrightarrow & K/Q_n & \longrightarrow & H/Q_n & \longrightarrow & H/K & \longrightarrow & 1
 \end{array}$$

And the same way, f_n is an isomorphism.

Now we obtain the following corollary.

Corollary 2.3. Let $(f, f'): (G, N) \rightarrow (H, K)$ are group homomorphisms satisfy the hypotheses of Theorem 2.2. Suppose further that N and K are \mathcal{V}_G -nilpotent and \mathcal{V}_H -nilpotent, respectively. Then f and f' are isomorphisms.

Proof. The assertion follows from Theorem 2.2 and the remark that there exists $n \geq 0$ such that $V_n(N, G) = \{1\}$ and $V_n(K, H) = \{1\}$.

Now, we have the following theorem, which is a generalization of Stalling's theorem [15].

Theorem 2.4. Let \mathcal{V} be a variety of groups and $f: G \rightarrow H$ be an epimorphism. Let N be a \mathcal{V}_G -nilpotent normal subgroup of G and K be a normal subgroup of H such that $f(N) = K$. If $\ker f \subseteq [NV^*G]$ and $\mathcal{WVM}(H, K)$ is trivial, then f and $f|_N$ are isomorphisms.

Proof. Put $M = \ker f$, then $\frac{N}{[NV^*G]} \cong \frac{K}{[KV^*H]}$, $\frac{G}{N} \cong \frac{H}{K}$ and $\frac{V_n(N, G)M}{M} = V_n(K, H)$ for all $n \geq 0$. Now the result follows from Corollary 2.3.

Finally, a sufficient condition will be given such that the order of the generalized Baer-invariant of a pair of finite groups divides the order of the generalized Baer-invariant of the pair of its factor groups with respect to two varieties of groups. Let $\psi: E \rightarrow G$ be an epimorphism such that $\ker \psi \subseteq V^*(E)$. We denote by $(WV^*)^*(G)$ the intersection of all subgroups of the form $\psi(V^*(E))$. Clearly, $(WV^*)^*(G)$ is a characteristic subgroup of G which is contained in $V^*(G)$. In particular, if \mathcal{W} is the variety of all groups and \mathcal{V} is a variety of abelian groups then this subgroup is denoted by $Z^*(G)$ as in literature of Karpilovsky [2].

Now using the above concept we have the following Theorem.

Theorem 2.5. Let K be a normal subgroup of G contained in $N \cap (WV^*)^*(G)$. Then

$$|\mathcal{WVM}(G, N)| \text{ divides } |\mathcal{WVM}(G/K, N/K)|.$$

Proof. By theorem 3.2 of Neumann [5], natural homomorphism $\mathcal{WVM}(G) \rightarrow \mathcal{WVM}(G/K)$ will be a monomorphism. Now the following commutative diagram

$$\begin{array}{ccc}
 \mathcal{WVM}(G, N) & \xrightarrow{\subseteq} & \mathcal{WVM}(G) \\
 \downarrow & & \downarrow \\
 \mathcal{WVM}(G/K, N/K) & \xrightarrow{\subseteq} & \mathcal{WVM}(G/K)
 \end{array}$$

implies that the natural homomorphism $\mathcal{WVM}(G, N) \rightarrow \mathcal{WVM}(G/K, N/K)$ is also a monomorphism. Thus Lemma 1.2 (i) implies that $\mathcal{WVM}(G, K)$ is trivial. Now we have $|\mathcal{WVM}(G/K, N/K)| = |K \cap [NV^*G]| |\mathcal{WVM}(G, N)|$, which completes the result.

Acknowledgment

The authors wish to thank Yazd Branch, Islamic Azad University for its support of research project under the title the Generalization of the Stalling's theorem.

References

- Schur I (1904) Studies on the representation of finite groups by linear substitutions broken. J Reine Angew Math 132: 85-137.
- Karpilovsky G (1987) The Schur Multiplier. London Math Soc Monographs New Series 2.
- Hopf H (1942) Fundamental gruppe und zweite bettische gruppe. Comment Math Helvetici 14: 257-309.
- Ellis G (1998) The schur multipliers of a pair of groups. Appl categ Structures 6: 355-371.
- Neumann H (1967) Varieties of groups. Bulletin of the American Mathematical Society 73: 603-613.
- Hekster NS (1989) Varieties of groups and isologisms. J Austral Math Soc (Series A) 46: 22-60.
- Baer R (1945) Representations of groups as quotient groups, I-III. Trans Amer Math Soc 58: 295-419.
- Leedham-Green CR, McKay S (1976) Baer-invariant, isologism, varietal laws and homology. Acta Math 137: 99-150.
- Schur I (1904) About the representation of finite groups by linear substitutions broken. J Reine Angew Math 127: 20-50.
- Baer R (1952) Endlichkeitskriterien für Kommutatorgruppen. Math Ann 124: 161-177.
- Moghaddam MRR, Salemkar AR, Rismanchian MR (2002) Some properties of ultra Hall and Schur pairs. Arch Math (Basel) 78: 104-109.
- Fung WKH (1977) Some theorems of Hall type. Arch Math 28: 9-20.
- Hilton PJ, Stammbach U (1970) A course in homological algebra. Springer-Verlag Berlin.
- Rotman JJ (2009) An introduction to homological algebra. Second Edition Universitext Springer New York.
- Stallings J (1965) Homology and central series of groups. J Algebra 2: 170-181.

This article was originally published in a special issue, **Recent Advances of Lie Theory in differential Geometry, in memory of John Nash** handled by Editor. Dr. Princy Randriambolondrantomalala, University of Antananarivo, Madagascar