

The Human Microbiome as a New Source for Antibiotic Discovery

Zhi-Qiang Xiong*

School of Medical Instrument and Food Engineering, University of Shanghai for Science and Technology, Shanghai 200093, China

*Corresponding author: Zhi-Qiang Xiong, School of Medical Instrument and Food Engineering, University of Shanghai for Science and Technology, Shanghai 200093, China, Tel: +86-21-55803272; E-mail: xiongzq@hotmail.com

Received date: December 25, 2016; Accepted date: December 27, 2016; Published date: December 30, 2016

Citation: Xiong ZQ (2016) The Human Microbiome as a New Source for Antibiotic Discovery. Clin Microbiol 5: e137. doi: 10.4172/2327-5073.1000e137

Copyright: © 2016 Xiong ZQ. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Editorial

Bacterial infection especially antibiotic-resistance “superbugs” (e.g., methicillin- and vancomycin-resistant *Staphylococcus aureus*) has caused a huge global threat of public health [1-3]. Bacterial infection currently kills more people than HIV, and will kill more people than cancer in the coming decades [4,5]. Nevertheless, new antibiotics, one of the most important treatment options for bacterial infection, unfortunately continue to be limited because the pace of antibiotic discovery and development is dramatically declining [6]. To meet the urgent need for novel drugs with combating the “superbugs”, researchers are continuing to explore new natural sources such as marine resource [6], metagenome [2], and uncultured microorganisms [1] that were missed in previous screens [4].

The human microbiota (~100 trillion microbes) is the full collection of microorganisms including bacteria, fungi, and archaea that reside inside and on human tissues and biofluids such as skin, lung, and gastrointestinal tracts. The human microbiome (~8 million genes) refers to an aggregate of genes in the human microbiota [7]. Human microbiome has been considered as a counterpart to the human genome, which encodes at least 100 times as many genes as the entire human own genome [8]. Our understanding of the relationship between human and its symbiotic microbiota continues to revolutionize by the persistent studies of the human microbiome [9]. More and more studies demonstrated that the microbiota is essential for human health such as nutrition, neurobiology, immunology, and various diseases like obesity and diabetes [9,10].

With the rapid progress of human microbiome project, numerous studies showed that human microbiome has a hugely diverse and dynamic [5]. Nevertheless, contrary to the immense diversity of human microbiome, human microbiota has only been found to produce bacteriocins with antimicrobial activities against closely related bacteria [11], not produce complex bioactive compounds. In 2014, Donia et al. used a systematic approach by combination of chemistry, genetics, metagenomics, and metatranscriptomics to identify >14,000 biosynthetic gene clusters responsible for 3,118 small-molecule from the genomes of human microbiome [12], suggesting that human microbiome as a new resource has the application potential for finding new antibiotics. More importantly, they discovered that thiopeptides, a class of ribosomally synthesized antibiotics in clinical trials, are widely distributed in the human microbiota. A new thiopeptide antibiotic, lactocillin, isolated from the vaginal microbiota, has potent antibacterial activity against a range of Gram-positive vaginal pathogens [12].

The work by Donia et al. [12] strongly inspired us that human microbiome could harbor rich and diverse bioactive compounds for human health. This opinion was further confirmed this year because of

two novel complex bioactive compounds humimycins [13] and lugdunin [5] recently isolated from human microbiome. Humimycins and lugdunin, both synthesized by NRPS (non-ribosomal peptide synthetase) gene clusters, had broad antibacterial activities against human-associated commensal and antibiotic-resistance “superbugs” [5]. Despite the above works have significant effect on antibiotic discovery from human microbiome, new approaches and techniques are still required for finding more structurally unique antibiotics. I believe, with the better understanding of human microbiome in the next decade, bioprospecting of human microbiome has a great application potential for the discovery of new antibiotics.

References

1. Xiong ZQ (2016) Bioprospecting of Uncultured Microorganisms: The Dawning of Antibiotic Discovery. Clin Microbiol 5: e132.
2. Xiong Z (2013) Metagenomic-Guided Antibiotics Discovery. Clin Microbiol 2: 101.
3. Xiong ZQ, Zhang ZP, Li JH, Wei SJ, Tu GQ (2012) Characterization of *Streptomyces padanus* JAU4234, a Producer of Actinomycin X2, Fungichromin, and a New Polyene Macrolide Antibiotic. Appl Environ Microbiol 78: 589-592.
4. Nothias LF, Knight R, Dorrestein PC (2016) Antibiotic discovery is a walk in the park. Proc Natl Acad Sci USA 113: 14477-14479.
5. Zipperer A, Konnerth MC, Laux C, Berscheid A, Janek D, et al. (2016) Human commensals producing a novel antibiotic impair pathogen colonization. Nature 535: 511-516.
6. Xiong ZQ, Wang JF, Hao YY, Wang Y (2013) Recent advances in the discovery and development of marine microbial natural products. Mar Drugs 11: 700-717.
7. Turnbaugh PJ, Ley RE, Hamady M, Fraser-Liggett CM, Knight R, et al. (2007) The human microbiome project. Nature 449: 804-810.
8. Jones BV, Begley M, Hill C, Gahan CG, Marchesi JR (2008) Functional and comparative metagenomic analysis of bile salt hydrolase activity in the human gut microbiome. Proc Natl Acad Sci USA 105: 13580-13585.
9. Cho I, Blaser MJ (2012) The human microbiome: at the interface of health and disease. Nat Rev Genet 13: 260-270.
10. Jones ML, Tomaro-Duchesneau C, Prakash S (2014) The gut microbiome, probiotics, bile acids axis, and human health. Trends Microbiol 22: 306-308.
11. Kommineni S, Bretl DJ, Lam V, Chakraborty R, Hayward M, et al. (2015) Bacteriocin production augments niche competition by enterococci in the mammalian gastrointestinal tract. Nature 526: 719-722.
12. Donia MS, Cimermanic P, Schulze CJ, Wieland Brown LC, Martin J, et al. (2014) A systematic analysis of biosynthetic gene clusters in the human microbiome reveals a common family of antibiotics. Cell 158: 1402-1414.
13. Chu J, Vila-Farres X, Inoyama D, Ternei M, Cohen LJ, et al. (2016) Discovery of MRSA active antibiotics using primary sequence from the human microbiome. Nat Chem Biol 12: 1004-1006.