

# The profile of mentally ill offenders in Katsina, Northern-Nigeria

The shift towards community care for patients with chronic mental illness has resulted in an increase in the number of forensic patients.<sup>1</sup> This has been associated with other factors such as gender, education, poverty, diagnosis, psychopathology and lack of access to mental health services.<sup>2-4</sup> In Sub-Saharan Africa and other Low and Middle Income Countries (LAMIC) access to mental health facilities is poor and often difficult.<sup>5</sup> A study in Nigeria estimated that 70% of the population had no access to mental health facilities.<sup>6</sup> Moreover, in Nigeria, all the mental health centres are located in urban areas and require out-of-pocket payment for access to their services.<sup>6</sup> The situation is further complicated by the tortuous pathway to mental health services in Nigeria leading to delay in presentation, increased morbidity and poor outcome.<sup>7</sup> Schizophrenia, substance use disorders and organic mental disorders are the commonly reported diagnoses among mentally ill offenders.<sup>4,8,9</sup> These disorders can be managed adequately by providing accessible and affordable mental health services in the community. Such services might likely reduce offending behaviours among mentally ill patients.

This retrospective study was conducted at the Katsina State Psychiatric Hospital in Katsina, Northern Nigeria. Materials reviewed were the clinical records and forensic assessment reports of all the patients referred to the hospital by the criminal justice system over a 2 year period. Information retrieved for analysis included socio-demographic characteristics, diagnoses, offending behaviour and other related indices.

A total of 19 patients were referred to the hospital for forensic assessment during the period under review. The mean age of the patients was  $28.94 \pm 9.34$  years. The mean age of the female patients was  $32.40 \pm 14.82$  years, while that of male patients was  $27.71 \pm 6.85$  years. There was no significant difference between the mean ages of the male and female patients ( $t = .68, p = .53$ ). The mean duration of illness among these patients was  $8.05 \pm 7.78$  years. All the patients were either single or divorced as at the time of presentation. None of the patients was gainfully employed at the time of presentation. Other socio-demographic characteristics of the patients are shown in Table I. Homicide was the most common offending behavior committed by these patients, accounting for 68% of all the offences. There was no significant difference in the mean ages and duration of illness between patients that committed homicide and other offending behaviours respectively ( $t = .016, p = .98$ ;  $t = .48, p = .71$ ). The act of homicide was significantly associated with living in a rural area ( $\chi^2 = 7.36, p = .007$ ), not

having formal education ( $\chi^2 = 4.99, p = .046$ ) and having a diagnosis of schizophrenia ( $\chi^2 = 10.87, p = .003$ ). Relatives of the patients were the most likely victims of offending behavior ( $\chi^2 = 12.06, p = .001$ ). Gender, substance use and previous contact with mental health services were not significantly associated with homicide.

The socio-demographic characteristics of the patients are similar to those reported by other studies of mentally abnormal offenders.<sup>2-4</sup> The majority of the patients are young, unemployed males, single without formal education. Similar observations were made by Ogunlesi et al in Southern Nigeria and Pal in Papua-New Guinea.<sup>8,2</sup>

**Table I: Socio-demographic characteristics of the patients**

Variable	Frequency (N)	Percentage (%)
1 Gender		
Male	14	73.7
Female	5	26.3
2 Place of Residence		
Rural	14	73.7
Urban	5	26.3
3 Type of crime		
Homicide	13	68.4
Non-Homicidal	6	31.6
4 Education		
Formal education	6	31.6
No formal education	13	68.4
5 Diagnosis		
Schizophrenia	13	68.4
Non-Schizophrenic	6	31.6
6 Victim		
Relative	11	57.9
Non Relative	8	42.1
7 Cor-morbid substance use		
Present	11	57.9
Absent	8	42.1
Total		
8 Previous contact with Hospital		
History of contact	3	15.8
No history of contact	16	84.2

The most common psychiatric diagnosis among the studied group of patients was schizophrenia with homicide the most common offending behavior. There were significant relationships between place of residence, education, diagnosis and the act of homicide. The patients from rural areas constituted the majority of mentally ill offenders seen in this study and they are more likely to commit homicide compared to those from urban areas. This might be explained either due to lack of access to mental health services or that they are more easily apprehended by the criminal justice system than those from urban areas. Education not only predisposes to poverty but also influences health seeking behavior. In this study not having formal education was significantly associated with offending behaviour and the act of homicide. Patients with out formal education constituted the majority of the mentally abnormal offenders. Similar observations were made by Ogunlesi et al in Southern Nigeria.<sup>8</sup> The diagnosis of schizophrenia was associated with offending behaviour and homicide in this study which might be explained by the nature of the illness with psychopathologies like command auditory hallucinations and delusions. The study further confirmed that relatives of patients with chronic mental illness are often the victims of the offending behaviour. This might be explained by close associations between the relatives and the patient, which might lead to conflict. Similar observations have been made.<sup>4,9</sup> The study did not find a significant relationship between the act of homicide and previous contact with a mental health facility. This might be because the majority of the patients had no contact with a mental health facility prior to presentation for forensic assessment and treatment.

The provision of accessible and affordable mental health services in the community might reduce the incidence of offending behavior among this group of patients. There is a need to improve provision of mental health facilities, as well as to educate the community about the availability of

effective treatment for mental disorders, in developing countries- especially in the rural areas where most of the population lives.

**AJ Yusuf<sup>1</sup>, FT Nuhu<sup>2</sup>**

<sup>1</sup>Department of Psychiatry Ahmadu Bello University Teaching Hospital, Shika-Zaria, Nigeria

<sup>2</sup>Federal Neuro-Psychiatry Hospital, Barnawa, Kaduna, Nigeria  
email: yusufajika@yahoo.com

### References

1. Kram P, Gabrielsen G. *Crime committed by mentally ill person 1977-1999: development, numbers and cause. Ugeskr Laeger* 2003; 16:2553-2558.
2. Pal S. *Mental disorders in abnormal offenders in Papua New Guinea. Med Law* 1997; 16:87-89.
3. Glancy GD, Regtir C. *The forensic aspect of Schizophrenia. Psychiatr Clin North Am* 1992; 15:579-589.
4. Fazal S, Grann M, Calstrom E, Lichenstien P, Langstrom N. *Risk factors for violent crime in Schizophrenia: a national cohort study of 13,886 patients. J Clin Psychiatry* 2009; 70: 362-369.
5. *World Health Organization. Mental Health Atlas. Geneva: World Health Organization. www.globalatlast.who.int/globalatlast/default.asp (accessed 20 December 2008).*
6. Ayorinde O, Gureje O, Lawal R. *Psychiatric research in Nigeria: bridging tradition and modernisation. Br J Psychiatry* 2004; 184: 536-538.
7. Gureje O, Lasebikan O. *Use of mental health services in a developing country: results from the Nigerian of mental health and wellbeing. Soc Psychiatry Psychiatr Epidemiol* 2006; 41:44-49.
8. Ogunlesi OA, Makanjuola MA, Adelekan ML. *Offenders admitted to the Neuro-Psychiatry Hospital, Aro, Abeokuta: A ten year review. West Afr J Med* 1988; 7:129-135.
9. Mafullul YM. *Criminal homicide in Nigeria: a socio-psychiatric review. Arch Ibadan Med* 2003; 4: 23-26.