

The Use of *Streptomyces coelicolor* in the Removal of Heavy Metals

Ayten Öztürk*

Department of Biotechnology, Art and Science Faculty, University of Niğde, Turkey

Abstract

In recent years, in order to improve the standard of living of human being, outstanding studies have increased on microbial technology. This article focuses on the interactions between secondary metabolites of *Streptomyces* and heavy metals, as they have been presented in literature stemming from two different studies: The removal of heavy metal and the role of heavy metal on secondary metabolites of *Streptomyces* and how these interactions fit into accepted paradigms of response to heavy metal, resistance mechanisms and production of secondary metabolites.

Keywords: *Streptomyces coelicolor*; Removal of heavy metal; Secondary metabolites

Introduction

Organisms known as actinomycetes are the richest source of special metabolites. They are conventionally used to describe filamentous, Gram positive actinobacterium existing in soil, including those of the most prolific genus *Streptomyces*. This genus is home to the best known strain A3 (2) of *Streptomyces coelicolor* which is the genetically best-defined streptomycete [1,2].

Streptomyces coelicolor are used as a model representative of a group of soil-dwelling organisms to understand microbial life in the highly competitive soil environment. It is known that bacteria which produce antibiotics can compete with other organisms for nutrient. These actinobacteria are notable for their production of over most compounds that are pharmaceutically important agents such as natural antibiotics, anti-tumour agents and immunosuppressant. These bacteria have over 20 cluster genes coding for known or predicted secondary metabolites and many pigments. However they have many adaptation strategies and capability in producing spores that can live in even dry and hot in soil [1,3,4].

New developing technologies based on microorganisms contribute to human health, by helping removal of different contaminant from the environment. Kothe et al. [4] have expressed that due to *Streptomyces coelicolor* active secondary metabolism producing some antibiotics and pigments; they could be a good source for the identification of some heavy metal binding components by possibly using biotechnological application [2].

Binding of heavy metals to *Streptomyces*

Although *Streptomyces coelicolor* A3(2) was found to be sensitive to high concentrations of heavy metals, [5,6] they proved to be tolerate higher concentrations and the findings have highlighted a new biologically relevant aspect of the chemical property of vancomycin as a zinc chelator in vitro. Because of some *Streptomyces* strains showed high resistance to some heavy metal such as nickel and zinc, they have been presumed to contribute a large heavy metal binding capacity [7-9].

In our previous study, we have demonstrated the removal of nickel and copper ions effectively from aqueous solution by using dried *Streptomyces coelicolor* A3 (2) [10]. Our results were essential in determining pharmaceutical disposals for bio-remediation and waste water treatment. Effective removal of heavy metals from wastewaters and industrial wastes still remains a major topic in nowadays research and various ways are there to cope with heavy metals discharged from industrial wastewaters to both water and soil

inhibiting microorganisms. However, physiological and biochemical studies in different *Streptomyces* strains have shown that resistance to heavy metals is produced by several mechanisms, acting either alone or in conjunction: an increased reductase activity; production of extracellular quelators; metal efflux pumps; intracellular sequestration and bio-mineralization [11]. Although heavy metals are known to be toxic; metals such as nickel, cobalt, copper, manganese, and others are considered essential micronutrients needed as cofactors in many enzymes [12] and nickel was reported to be necessary for the reactive center in urease and superoxide dismutase of streptomycetes [13].

Actinorhodin production by *Streptomyces coelicolor* A3 (2), and the effects of metals on the growth and antibiotic synthesis has been studied [14-16]. Abbas and Edwards [17] have reported that the reduction in the antibiotic production was observed when some metals were added during the antibiotic-producing phase and little reduction of final actinorhodin titers was observed [17]. However, due to the specific adaptation strategies of some streptomycete strains, they can cope with high heavy metal concentrations [5,18]. Melanin-like pigment produced by some *Streptomyces* during growth in nickel containing media was reported to bind to metal and the capacity of this pigment to form nickel complexes was determined [3,19]. Similarly actinorhodin-related blue pigments produced by *Streptomyces coelicolor* A3 (2) can be used to form new complexes with heavy metals [4,20] that might be very convenient in the removal of heavy metals. The biomasses of *Streptomyces* species were reported to be capable of biosorption of heavy metals [21]. Schütze and Kothe [11] have mentioned that several morphological, physiological and reproductive characteristics coding on major genome of *Streptomyces* would allow its species to occupy extreme environments or contaminated environment [11]. Since these properties are shared by all species of the genus, it can be interpreted as an evidence for the inheritance of resistance to heavy metals.

Conclusion

Streptomyces species have got tolerance to variety of heavy metals.

*Corresponding author: Ayten Öztürk, Department of Biotechnology, Art and Science Faculty, University of Niğde, Turkey, E-mail: aozturk@nigde.edu.tr

Received February 04, 2016; Accepted February 10, 2016; Published February 17, 2016

Citation: Öztürk A (2016) The Use of *Streptomyces coelicolor* in the Removal of Heavy Metals. Adv Tech Biol Med 4: 168. doi: [10.4172/2379-1764.1000168](http://dx.doi.org/10.4172/2379-1764.1000168)

Copyright: © 2016 Öztürk A. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Therefore, *Streptomyces* that are waste products obtained from pharmaceutical manufacturers are vital in improving environmental biotechnology; however bioaccumulation studies of heavy metals and biosorption studies with most heavy metals in *Streptomyces coelicolor* were not studied to date. Bioaccumulation and biosorption studies combined with genomic, biochemical and physiological information will be indispensable for efficient wastewater treatment in future.

References

1. Bentley SD, Chater KF, Cerden AM, Tarraga O, Challis GL, et al (2002) Complete genome sequence of the model actinomycete *Streptomyces coelicolor* A3(2). Nature 417: 141-147.
2. Traxler MF, Kolter R (2015) Natural products in soil microbe interactions and evolution. Nat Prod Rep 32: 956-970.
3. Borodina I, Krabben P, Nielsen J (2005) Genome-scale analysis of *Streptomyces coelicolor* A3(2) metabolism. Genome Res 15: 820-829.
4. Kothe E, Bergmann H, Buchel G (2005) Molecular mechanisms in bio-geo-interactions: From a case study to general mechanisms. Chemie der Erde 65: 7-27.
5. Schmidt A, Haferburg G, Sineriza M, Merten D, Buchel G, et al. (2005) Heavy metal resistance mechanisms in Actinobacteria for survival in AMD contaminated soils. Chemie der Erde 65: 131-144.
6. Zarkan A, Macklyne HR, Truman AW, Hesketh AR (2016) The frontline antibiotic vancomycin induces a zinc starvation response in bacteria by binding to Zn(II). Sci Rep 6: 19602.
7. Schmidt A, Haferburg G, Schmidt A, Lischke U, Metren D, et al. (2009) Heavy metal resistance to the extreme: *Streptomyces* strains from a former uranium mining area. Chemie der Erde 69: 35-44.
8. Alvarez A, Catalano SA, Amoroso MJ (2013) Heavy metal resistant strains are widespread along *Streptomyces* phylogeny. Mol Phylogenet Evol 66: 1083-1088.
9. Daboor SM, Haroon AM, Esmael NAE, Hanona SI (2014) Heavy metal adsorption of *Streptomyces chromofuscus*. Journal of Coastal Life Medicine 2: 431-437.
10. Öztürk A, Artan T, Ayar A (2004) Biosorption of nickel(II) and copper(II) ions from aqueous solution by *Streptomyces coelicolor* A3(2). Colloids Surf B Biointerfaces 34: 105-111.
11. Schütze E, Kothe E (2012) Bio-Geo Interactions in Metal-Contaminated Soils. In: Kothe E, Varma A (Eds) Soil Biology 31. Springer-Verlag, Berlin Heidelberg.
12. Gill M (2014) Heavy metal stress in plants: a review. International Journal of Advanced Research, Volume 2: 1043-1055.
13. Haferburg G, Groth I, Möllmann U, Kothe E, Sattler I (2009) Arousing sleeping genes: shifts in secondary metabolism of metal tolerant actinobacteria under conditions of heavy metal stress. Biometals 22: 225-234.
14. Price B, Adamidis T, Kong R, Champness W (1999) A *Streptomyces coelicolor* antibiotic regulatory gene, *absB*, encodes an RNase III homolog. J Bacteriol 181: 6142-6151.
15. Elibol M (2004) Optimization of medium composition for actinorhodin production by *Streptomyces coelicolor* A3 (2) with response surface methodology. Process Biochemistry 39: 1057-1062.
16. Procópio RE, Silva IR, Martins MK, Azevedo JL, Araújo JM (2012) Antibiotics produced by *Streptomyces*. Braz J Infect Dis 16: 466-471.
17. Abbas AS, Edwards C (1990) Effects of Metals on *Streptomyces coelicolor* Growth and Actinorhodin Production. Appl Environ Microbiol 56: 675-680.
18. Hopwood DA (2006) Soil to genomics: the *Streptomyces* chromosome. Annu Rev Genet 40: 1-23.
19. Kothe E, Dimkpa C, Haferburg G, Schmidt A, Schmidt A, et al. (2010) *Streptomyces* Heavy Metal Resistance: Extracellular and Intracellular Mechanisms. In: Sheramati I, Varma A (Eds) Soil Heavy Metals 19: 225-235.
20. Bystrykh LV, Fernández-Moreno MA, Herrema JK, Malpartida F, Hopwood DA, et al. (1996) Production of actinorhodin-related "blue pigments" by *Streptomyces coelicolor* A3(2). J Bacteriol 178: 2238-2244.
21. Mosbah R, Sahmoune MN (2013) Biosorption of heavy metals by *Streptomyces* species. Cent Eur J Chem 11: 1412-1422.

Citation: Öztürk A (2016) The Use of *Streptomyces coelicolor* in the Removal of Heavy Metals. Adv Tech Biol Med 4: 168. doi: 10.4172/2379-1764.1000168

OMICS International: Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700 Open Access Journals
- 50,000 editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission/>