

Thoracic Anaesthesia in National Cancer Centre of Mongolia

Bolormaa B^{1*}, Ganbold L² and Avirmed D³

¹National Cancer Centre of Mongolia, Mongolia

²Health Sciences University, Ulaanbaatar, Mongolia

³Medical Research Institute of Mongolia, Mongolia

*Corresponding author: Bolormaa B, National Cancer Centre of Mongolia, Mongolia, E-mail: batnasan_bolormaa@yahoo.com

Received date: July 02, 2016; Accepted date: September 26, 2016; Published date: September 30, 2016

Copyright: © 2016 Bolormaa B, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Cancer is the second cause of mortality in the population increasing over last ten years in Mongolia. In 2014 registered 5485 cancer patients and 3530 were new patients. Before 2007, thoracic surgery had used conventional tube in general anaesthesia, then surgical access was not enough, surgery to heart function collapsed load pressing force involved in the hand, which made interior breath from the lungs, heart and respiratory failure, lung injury. In addition, respiratory tract filled with blood sputum, and it takes long time in intensive care due to disorders such as conjunctivitis and content blocking surgical lengthening. During post-surgery complications and several deaths occurred.

Keywords: One lung ventilation; Thoracic anaesthesia

Introduction

Cancer is the second cause of mortality in the population increasing over last ten years in Mongolia. In 2014 registered 5485 cancer patients and 3530 were new patients. Before 2007, thoracic surgery had used conventional tube in general anaesthesia, then surgical access was not enough, surgery to heart function collapsed load pressing force involved in the hand, which made interior breath from the lungs, heart and respiratory failure, lung injury. In addition, respiratory tract filled with blood sputum, and it takes long time in intensive care due to disorders such as conjunctivitis and content blocking surgical lengthening. During post-surgery complications and several deaths occurred.

Goal

When NCC thoracic surgery double lumen tube used as a real possibility right and left bronchial tube separates the lungs, reduce surgical time and postoperative complications.

Study Objectivities

The purpose of thoracic anaesthesia used double lumen tube is adapted Mongolian human characteristics, surgery and postoperative complications and it prevent to increase the death.

Materials and Methods

This study gives in the National Cancer Centre in 2012-2014 during thoracic surgery; double lumen tube anaesthesia department in meeting the study inclusion criteria included in the 2012-2014. 160 patients in treatment groups, III hospital (Shastin's) 160 clinical cardiovascular surgeries, patients took part in the control group.

During the study, pairs and study and control group patients, arterial blood 0.1-0.2ml of oxygen partial pressure (PaO₂), carbon

dioxide pressure (PaCO₂), conducted a study monitoring of oxygen saturation (SaO₂) and acidity (PH).

The study revealed that anaesthesia using mechanical ventilation (CPAP, PEEP, PSV, PCV, ACV, CMV and SIMV) form.

Results

During one lung anaesthesia average in monitor (SpO₂) -95.09% ± 1.07 and blood (SaO₂) -92.65% was ± 5.69 (P<0.032). After surgery, this study has ICU-average 2.2 ± 1.35 days, he had complications 19.65% and 1.64% of death. In 2003 NCC postoperative ICU of stay 6-10 days, of complications -37%, death was 43% (Tables 1 and 2).

Subjects	Result	P утга
Tidal volume of both lungs	7.77 ± 1.07 ml/ kg	
Tidal volume of one lung	5.87 ± 0.46 ml/kg	p<0.014
DLT diameter (Mongolian female)	3.43 ± 2.25 (Fr)	p<0.093
DLT diameter (Mongolian male)	37.09 ± 4.69 (Fr)	
DLT deep (151-160 cm height female)	27.68 ± 2.47 cm	
DLT deep (161-170 cm height male)	28.43 ± 2.6 cm	p<0.004
In noninvasive (SaO ₂)	95.09 ± 1.07%	p<0.032
In arterial blood (SpO ₂)	92.65 ± 5.69%	
Variation of PaCO ₂	37.11 ± 14.6	p<0.028
Variation of PaO ₂	119.15 ± 49.52	

Table 1: The statistical result of double lumen endotracheal tube placement.

Surgery type	N	%
--------------	---	---

<i>Esophagus surgery</i>		
Ivory Lewis	37	23.10%
Other	19	11.90%
<i>Lung surgery</i>		
Pulmectomy	24	15%
Lobectomy, segmectomy ,resections	57	35.60%
To probe thoracotomy	3	1.90%
Other (Tuberculosis, Ehinococcus,... stomy)	20	12.50%

Table 2: Comparison of operation type.

Conclusion

During our study, one lung ventilation blood oxygen levels in the peripheral veins (SpO₂) 95.09 ± 1.07%, arterial blood (SaO₂) 92.65 ± 5.69% (P<0.032). In the study group, Mongolian woman having double lumen tube 35 (Fr) diameters 27.68 ± 2.47 cm depth, of 37 males (Fr) felt that the appropriate place deep in diameter 28.43 ± 2.6 cm.

Other studies this depth is usually between 28-30 cm and 170-190 cm patient. Researchers have every 10cm height double lumen tube placement changed to 1 cm [40].

We significantly had complications 19.65% and 1.64% of death. During one lung anaesthesia 18.75% of patients in the study group used a mechanical ventilation forms are considered a kind of post-surgery complications, one of the most important factor in reducing mortality.

Other researchers performed a retrospective review of all perioperative deaths following esophagectomy for oesophageal cancer at the Mayo Clinic, Rochester from 1993 through 2009. Of 1522 esophagectomies, perioperative mortality occurred in 45 (3.0%) [41].

8 of 16 Dutch cardiothoracic centres participated and collected data on 4066 procedures and 183 surgical site infections, revealing a surgical site infection rate of 2.4% for sternal wounds and 3.2% for harvest sites. 61% of all surgical site infections were recorded after discharge [42-51]. Our study the comparative analysis other researchers, in one lung anaesthesia process felt the safety of patients in Mongolia.

References

- (2012) Report Health Ministry of Mongolia.
- (2013) Statistic National Cancer Centre of Mongolia.
- (2007) Statistic National Cancer Centre of Mongolia.
- Gonchigsuren D (1998) Problem the lung cancer diagnostic for X-ray and CT. Ulaanbaatar pp: 61.
- Brodsky J (1995) Anaesthesia for Thoracic Surgery. A Practice of Anaesthesia. pp: 1148-1155.
- Brodsky J (1995) Anaesthesia for Thoracic Surgery. A Practice of Anaesthesia. (6thedn). pp: 1160-1170.
- Brodsky JB, Tzabazis A, Basarab-Tung J, Shrager JB (2013) Sequential bilateral lung isolation with a single bronchial blocker. A A Case Rep 1: 17-18.
- Morgan EG, Mikhail MS, Murray MJ (2013) Clinical Anesthesiology International. (5thedtn). Wylie and Churchill Davidson.

- Slinger P, Triolet W, Wilson J (1988) Improving Arterial Oxygenation during One-lung Ventilation Anesthesiology 68: 291-295.
- Seymour AH (2003) The relationship between the diameters of the adult cricoids ring and main tracheobronchial tree: a cadaver study to investigate the basis for double-lumen tube selection. J Cardiothorac Vasc Anesth 17: 299-301.
- Ehrenfeld JM, Walsh JL, Sandberg WS (2010) Right- and left-sided Mallinckrodt doublelumen tubes have identical clinical performance. Anesthesia and Analgesia 57: 293-300.
- Campos JH, Hallam EA, Van Natta T, Kernstine KH (2006) Devices for lung isolation used by anesthesiologists with limited thoracic experience: comparison of double-lumen endotracheal tube, Univent torque control blocker, and Arndt wire-guided endobronchial blocker. Anesthesiology 104: 261-266.
- Finlayson GN, Chiang AB, Brodsky JB, Cannon WB (2008) Intraoperative contralateral tension pneumothorax during pneumonectomy. Anesthesia & Analgesia 106: 58-60.
- Knoll H, Ziegeler S, Schreiber JU, Buchinger H, Bialas P, et al. (2006) Airway injuries after one-lung ventilation: a comparison between double-lumen tube and endobronchial blocker: a randomized, prospective, controlled trial. Anesthesiology 105: 471-477.
- Yüceyar L, Kaynak K, Cantürk E, Aykaç B (2003) Bronchial rupture with a left-sided polyvinylchloride double-lumen tube. Acta Anaesthesiol Scand 47: 622-625.
- Brodsky JB, Lemmens HJ (2005) Tracheal width and left double-lumen tube size: a formula to estimate left-bronchial width. J Clin Anesth 17: 267-270.
- Amar D, Desiderio DP, Heerdt PM, Kolker AC, Zhang H, et al. (2008) Practice patterns in choice of left double-lumen tube size for thoracic surgery. Anesth Analg 106: 379-383.
- Campos JH (2005) Progress in lung separation. Thorac Surg Clin 15: 71-83.
- Sucato DJ, Girgis M (2002) Bilateral pneumothoraces, pneumomediastinum, pneumoperitoneum, pneumoretroperitoneum, and subcutaneous emphysema following intubation with a double-lumen endotracheal tube for thoracoscopic anterior spinal release and fusion in a patient with idiopathic scoliosis. J Spinal Disord Tech 15: 133-138.
- Weng W, DeCrosa DJ, Zhang H (2002) Tension pneumothorax during one-lung ventilation: a case report. J Clin Anesth 14: 529-531.
- Brodsky JB (2000) Is bronchoscopy necessary for insertion of double-lumen endotracheal tubes? : Con: Bronchoscopy is Not Necessary. J Bronchology 7: 78-83.
- Brodsky JB, Lemmens HJ (2003) Left double-lumen tubes: clinical experience with 1,170 patients. J Cardiothorac Vasc Anesth 17: 289-298.
- Fortier G, Coté D, Bergeron C, Bussièrès JS (2001) New landmarks improve the positioning of the left Broncho-Cath double-lumen tube-comparison with the classic technique. Can J Anaesth 48: 790-794.
- Brodsky J (1995) Anaesthesia for Thoracic Surgery. A Practice of Anaesthesia (6thedtn) Wylie and Churchill Davidson, pp: 1148-11702.
- Kavanagh BP, Katz J, Sandler AN (1994) Pain control after thoracic surgery. A review of current techniques. Anesthesiology 81: 737-759.
- Guinard JP, Mavrocordatos P, Chiolero R, Carpenter RL (1992) A randomized comparison of intravenous versus lumbar and thoracic epidural fentanyl for analgesia after thoracotomy. Anesthesiology 77: 1108-1115.
- Müller LC, Salzer GM, Ransmayr G, Neiss A (1989) Intraoperative cryoanalgesia for postthoracotomy pain relief. Ann Thorac Surg 48: 15-18.
- Campos JH (2002) Current techniques for perioperative lung isolation in adults. Anesthesiology 97: 1295-1301.
- Campos JH, Gomez MN (2002) Pro: right-sided double-lumen endotracheal tubes should be routinely used in thoracic surgery. J Cardiothorac Vasc Anesth 16: 246-248.
- de Bellis M, Accardo R, Di Maio M, La Manna C, Rossi GB, et al. (2011) Is flexible bronchoscopy necessary to confirm the position of double-

- lumen tubes before thoracic surgery? *Eur J Cardiothorac Surg* 40: 912-916.
31. WebMD does not provide medical advice, diagnosis or treatment.
 32. (2012) Statistics and Epidemiology Cancer Registries. Self-instructional Manual for Tumor Registrars. SEER.
 33. Powell ES, Cook D, Pearce AC, Davies P, Bowler GM, et al. (2011) A prospective, multi centre, observational cohort study of analgesia and outcome after pneumoectomy. *Br J Anaesth* 106: 364-70.
 34. Smith GB, Hirsch NP, Ehrenwerth J (1986) Placement of double-lumen endobronchial tubes. Correlation between clinical impressions and bronchoscopic findings. *Br J Anaesth* 58: 1317-1320.
 35. Seo JH, Bae JY, Kim HJ, Hong DM, Jeon Y, et al. (2015) Misplacement of left-sided double-lumen tubes into the right mainstem bronchus: incidence, risk factors and blind repositioning techniques. *BMC Anesthesiology* 15: 157.
 36. Allman KG, Wilson LH (2006) Oxford handbook of anaesthesia (3rd edn). Oxford university press.
 37. Brodsky JB, Lemmens HJ (2005) Tracheal width and left double-lumen tube size: a formula to estimate left-bronchial width. *J Clin Anesth* 17: 267-270.
 38. Marc S, Orlewicz, Meyers AD (2012) Double-Lumen Endotracheal Tube Placement. 25: 645-651.
 39. Schieman C, Wigle DA, Deschamps C, Nichols FC, Cassivi SD, et al. (2012) Patterns of operative mortality following esophagectomy. *Dis Esophagus* 25: 645-651.
 40. Manniën J, Wille JC, Kloek JJ, van Benthem BH (2011) Surveillance and epidemiology of surgical site infections after cardiothoracic surgery in The Netherlands, 2002-2007. *J Thorac Cardiovasc Surg* 141: 899-904.
 41. Fitzmaurice BG, Brodsky JB (1999) Airway Rupture From Double-Lumen Tubes. *Journal of Cardiothoracic and Vascular Anesthesia* 13: 322-329.
 42. Bellis MD, Accardo R, Maio MD, Manna CL, Rossi GB, et al. (2011) Is flexible bronchoscopy necessary to confirm the position of double-lumen tubes before thoracic surgery?. *Eur J Cardiothorac Surg* 40: 912-916.
 43. Brodsky JB, Lemmens HJ (2005) Tracheal width and left double-lumen tube size: a formula to estimate left-bronchial width. *J Clin Anesth* 17: 267-270.
 44. Morgan EG, Mikhail MS, Murray MJ (2013) *Clinical Anesthesiology (5edn)*. Lange Medical Books/ McGraw-Hill Medical Publishing Companies pp: 487.
 45. Slinger P, Sullivan EA (2013) *A Practical Approach to Cardiac Anesthesia. Anesthetic Management for Surgery of the Lungs and Mediastinum*.
 46. Powell ES, Cook D, Pearce AC, Davies P, Bowler GMR, et al. (2011) A prospective, multicenter, observational cohort study of analgesia and outcome after pneumoectomy. *Br J Anaesth* 106: 364-370.
 47. Kurtz I, Kraut J, Ornekian V, Nguyen MK (2008) Acid-base analysis: a critique of the Stewart and bicarbonate-centered approaches. *Am J Physiol Renal Physiol* 294: F1009-1031.
 48. Shimizu T, Abe K, Kinouchi K, Yoshiya I (1997) Arterial oxygenation during one lung ventilation. *Can J Anaesth* 44: 1162-1166.
 49. Severgnini P, Selmo G, Lanza C, Chiesa A, Frigerio A, et al. (2013) Protective mechanical ventilation during general anesthesia for open abdominal surgery improves postoperative pulmonary function. *Anesthesiology* 118: 1307-1321.
 50. Huang CC, Chou AH, Liu HP, Ho CY, Yun MW, et al. (2005) Tension Pneumothorax Complicated by Double-Lumen Endotracheal Tube Intubation. *Chang Gung Med J* 28: 503-507.
 51. Hemmes SN, Gama de Abreu M, Pelosi P, Schultz MJ (2014) High versus low positive end-expiratory pressure during general anaesthesia for open abdominal surgery (PROVHILO trial): a multicentre randomised controlled trial. *Lancet* 384: 495.