

Thoracic Spinal Cord Atrophy Secondary to Vitamin B12 Deficiency

Frederico Carvalho de Medeiros¹, Lucas Alverne Freitas de Albuquerque^{2*} and Cíntia Alvarenga Pereira¹

¹Department of Neurology, Santa Casa de Belo Horizonte Hospital, Minas Gerais, Brazil

²Department of Neurosurgery, Santa Casa de Belo Horizonte Hospital, Minas Gerais, Brazil.

*Corresponding author: Lucas Alverne Freitas de Albuquerque, Rua Padre Rolim, number 492, apt 101. Zip code 30130-090. Santa Efigênia, Belo Horizonte – MG, Brazil, Tel: +55(85)96279966; Email: lucasalverne@yahoo.com.br

Rec date: Aug 06, 2014, Acc date: Sep 06, 2014, Pub date: Sep 08, 2014

Copyright: © 2014 Medeiros FC, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Keywords: B12 deficiency; MRI; Myelopathy

Description

A 57 year-old woman searched for a neurological Service for the first time when she was 45-year-old, complaining about weakness in

lower limbs for ten months period. The physical examination reported was of sensitive level at T5, reduced proprioception, Romberg's sign and ataxic gait. A Work up discovered a low Vitamin B12 level of 64pg/ml. Vitamin B12 reposition was initiated, however the patient failed to follow up.




Figure 1: Sagittal T2-weighted MRI shows: A) preserved cervical spinal cord and B) important thoracic spinal cord atrophy; C) axial T2-weighted MRI demonstrates a pronounced reduction in the diameter of the thoracic spinal cord.

The patient looked for our Hospital with 57-year-old moving herself using a walker. A cervical and thoracic spinal cord MRI was performed (Figure 1). A sagittal and axial thoracic spinal MRI showed diffuse spinal cord atrophy with preservation of signal intensity at T2-weighted image (T2WI).

The myelopathy of vitamin B12 deficiency is rare and normally affect the lower cervical and upper thoracic regions [1,2]. Extensive

thoracic myelopathy due to vitamin B12 deficiency is very rare [2]. The predominant MRI finding is demyelination involving the dorsal columns, showed by the hyperintense signal at T2 [1-3]. In these cases the clinical and radiological improvements are possible if early therapy is initiated promptly [1].

Spinal cord atrophy secondary to prolonged vitamin B12 deficiency is an extremely rare complication of the myelopathy. We believe that

delayed treatment and prolonged inadequate therapy induces irreversible axonal degeneration with progressive fibrous gliosis, promoting spinal cord atrophy (Figure 1) [1,3]. In this case it was associated to a very poor response to vitamin B12 replacement.

Acknowledgement

The authors would like to acknowledge João Lucas Breder for assistance with preparation of Figure1.

References

1. Pittock SJ, Payne TA, Harper CM (2002) Reversible myelopathy in a 34-year-old man with vitamin B12 deficiency. *Mayo Clin Proc* 77: 291-294.
2. Naidich MJ, Ho SU (2005) Case 87: Subacute combined degeneration. *Radiology* 237: 101-105.
3. Bassi SS, Bulundwe KK, Greeff GP, Labuscagne JH, Gledhill RF (1999) MRI of the spinal cord in myelopathy complicating vitamin B12 deficiency: two additional cases and a review of the literature. *Neuroradiology* 41: 271-274.