

# Title-Combinatory Effects of Diets with Three Protein Levels and Two Fat Levels on Growth Performance and Fillet Composition of Cage Cultured Genetically Improved Farmed Tilapia (GIFT)

De Silva MPKSK<sup>1\*</sup>, Senaarachchi WARK<sup>1</sup> and Liyanage NPP<sup>2</sup>

<sup>1</sup>Department of Zoology, University of Ruhuna, Matara, Sri Lanka

<sup>2</sup>Department of Animal Sciences, Uva-Wellasa University, Badulla, Sri Lanka

## Abstract

Effect of six diets having two lipid levels and three crude protein levels on growth and fillet composition of Genetically Improved Farmed Tilapia (GIFT) reared in cages was evaluated. Diets having Crude Protein levels (CP) 26%, 30% and 36% and Crude Fat levels (CF) (7% and 12%) were formulated. Experiment was set up according to 3 X 2 factorial designs. GIFT fingerlings ( $2.94 \pm 1.47g$ ) were stocked at a density of 75 fish /m<sup>3</sup> in net cages each having a water volume of 2 m<sup>3</sup>. Fish in quadruplicated treatments were fed with six diets for 150 days. Fish fed with 36% CP diets had significantly different ( $P \leq 0.05$ ) values for all growth indices; Final Mean Weight, Specific Growth Rate, Feed Conversion Ratio and Net Yield while no such difference found between 26% CP and 30% CP diets for the same growth indices. Increasing fat level from 7% to 12% had no significant effect on growth performances in GIFT in any protein level diet. Diets having higher CF level (12%) had significantly affected ( $P \leq 0.05$ ) the, CF content in fish filets while no protein sparing seen in fish filets. There was no marked difference in protein content in fish filets of fish fed with 30% and 36% CP diets. Findings reveal that increased fat levels do not contribute for protein sparing of fish and the diet having 36% CP and 7% CF is suitable for the grow out phase and 36/12 for fattening phase of GIFT reared in cages.

**Keywords:** Protein sparing; Fish feeds; Fish growth; Fillet composition; GIFT; Combinatory diets

## Introduction

Fish nutrition and feeding is one of the major requirements for sustainable aquaculture production. Feed is the most expensive component in fish culture and a major part of the total variable cost representing about 30% to 60% is allocated [1,2]. Protein is the principal organic constitute of animal tissues [2,3]. The economical and nutritional aspect of protein utilization in fish is to improve the synthesis of tissues rather than protein use as energy source [4,5]. To balance these two important aspects (nutritional value and cost effectiveness) aquaculture investigators are interested in to reduce protein content in diet without affecting the growth [6].

Protein requirement of tilapia varies with the size/age, protein quality, non-protein energy level, water temperature and salinity, presence of natural food and allowance of feed [7]. If there is insufficient amount of energy in the diet the excess amount of protein will convert in to energy by the fish [8-10].

Dietary lipid is good source of highly digestible energy and only source of essential fatty acids. Lipid also plays a sparing action for protein in fish if the adequate requirement of energy level through diet is provided by lipids [11,12].

Many studies have been conducted on sparing ability of lipid and carbohydrate (energy source) to protein for various fish species in variety of culture conditions [13,14]. Inclusion of higher levels of protein in fish diet also needs high energy levels to maintain desired protein to digestible energy ratio [5].

Increase in dietary lipids in fish feed than the requirement however could increase fat deposition in fish tissues and also negative effect on utilization of other nutrients resulting growth retardation [15,16].

Different dietary protein and oil levels on growth of tilapia had no indication of sparing effect on dietary protein [17]. Feed intake of

juvenile Nile tilapia (*Oreochromis niloticus*) increased with the increase of dietary proteins, while dietary energy seemed to have no influence on the growth [6]. Kim, et al. [5] demonstrated that the effect of different dietary protein (20% and 40%) and lipid levels (9% and 17%) growth and body composition of juvenile Far Eastern catfish, *Silurus asotus*. The protein requirement for the maximum growth of juvenile far eastern catfish is more than 40% and increasing the dietary lipid level from 9% to 17% had improved growth and feed utilization.

As such there is a need of proper understanding on nutritional requirements of a particular fish species and interaction of those nutrients on growth of fish when formulating diets for different fish species.

Fish species are cultured in cages in a wide variety of culture environments. Fish in cages is constrained by limited availability of fish feed of required quality [3]. Genetically Improved Farmed Tilapia (GIFT) a developed strain of *O. niloticus* and considered as a suitable culture fish species in cages [18]. With the aim of popularizing the GIFT cage culture in large number of reservoirs and tanks scattered throughout Sri Lanka, present study was conducted to evaluate the effect of two dietary lipid levels and three crude protein levels on growth and body composition of GIFT raised in cages.

**\*Corresponding author:** De Silva MPKSK, Department of Zoology, University of Ruhuna, Matara, Sri Lanka, Tel: +94713567998/+94 41 2227025; E-mail: kumududs@zoo.ruh.ac.lk

**Received** September 23, 2015; **Accepted** January 04, 2016; **Published** January 06, 2016

**Citation:** De Silva MPKSK, Senaarachchi WARK, Liyanage NPP (2016) Title-Combinatory Effects of Diets with Three Protein Levels and Two Fat Levels on Growth Performance and Fillet Composition of Cage Cultured Genetically Improved Farmed Tilapia (GIFT). J Aquac Res Development S2: 008. doi:10.4172/2155-9546.S2-008

**Copyright:** © 2016 De Silva MPKSK, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Ingredients (%)	Treatments (% Protein/% Lipid )					
	Formulated diets					
	CP/CF					
	26/7	26/12	30/7	30/10	36/7	36/12
Fish meal	20	20	25	25	35	35
Soya bean meal	20	20	27	27	35	35
Coconut seed cake	20	20	20	20	20	20
Maize	26	3	18	3	7	#
Rice bran	13	35	9	23	2	8
Vitamin & Mineral mix	1	1	1	1	1	1
Vitamin E	#	1	#	1	#	1
Nutrient content (dry matter basis)						
CP (%)	26.14	26.18	30.47	30.25	36.65	36.8
CF (%)	7.32	12.47	7.7	12.55	7.65	12.25
Ash (%)	7.06	8.47	7.45	8.55	10.01	10.11
Cost/Kg (US\$)*	0.55	0.5	0.63	0.6	0.77	0.75

CP – Crude Protein, CF – Crude Fat,  
#not included, \*Sri Lankan rupees converted in to US\$ (12.09.2015)

**Table 1:** Ingredient percentage, cost and chemical composition of the experimental diets (mean, n=2).

Diets CP/CF	FMW (g)	SGR (% day <sup>-1</sup> )	FCR	PER	NY (Kg/2m <sup>3</sup> )	SR (%)
26/7	111.12 ± 6.0 <sup>a</sup>	2.41 ± 0.03 <sup>a</sup>	1.54 ± 0.04 <sup>a</sup>	2.48 ± 0.07 <sup>a</sup>	13.6 ± 3.14 <sup>a</sup>	90.0 ± 3.54 <sup>a</sup>
26/12	109.58 ± 7.23 <sup>a</sup>	2.40 ± 0.04 <sup>a</sup>	1.72 ± 0.09 <sup>a</sup>	2.23 ± 0.12 <sup>a</sup>	11.9 ± 6.25 <sup>a</sup>	89.16 ± 3.59 <sup>a</sup>
30/7	117.14 ± 2.93 <sup>a</sup>	2.45 ± 0.01 <sup>a</sup>	1.61 ± 0.07 <sup>a</sup>	2.04 ± 0.09 <sup>a</sup>	12.9 ± 6.88 <sup>a</sup>	85.5 ± 4.77 <sup>a</sup>
30/12	113.74 ± 10.0 <sup>a</sup>	2.42 ± 0.05 <sup>a</sup>	1.51 ± 0.04 <sup>a</sup>	2.18 ± 0.06 <sup>a</sup>	13.8 ± 4.78 <sup>a</sup>	89.00 ± 3.77 <sup>a</sup>
36/7	158.26 ± 3.56 <sup>b</sup>	2.65 ± 0.01 <sup>b</sup>	1.22 ± 0.04 <sup>b</sup>	2.23 ± 0.08 <sup>a</sup>	18.0 ± 6.45 <sup>b</sup>	89.16 ± 1.77 <sup>a</sup>
36/12	166.32 ± 6.90 <sup>b</sup>	2.68 ± 0.02 <sup>b</sup>	1.14 ± 0.03 <sup>b</sup>	2.63 ± 0.2 <sup>a</sup>	19.0 ± 6.45 <sup>b</sup>	86.33 ± 4.33 <sup>a</sup>

<sup>1</sup>Values in the same column having different letters are significantly different (P≤0.05).  
FMW – Final Mean Weight, SGR – Specific Growth Rate, FCR – Feed Conversion Ratio, PER – Protein Efficiency Ratio, NY – Net Yield, SR – Survival Rate  
CP- Crude Protein, CF- Crude Fat

**Table 2:** Growth performance indices of GIFT fed with six experimental diets for 150 days (mean ± SE, n = 4)<sup>1</sup>.

## Material and Methods

The study was carried out in cages established in Kattakaduwa perennial reservoir (80 ha, N 06°07' 26.2"E 80°52' 41.2"), at Hambantota District (Southern Province) in Sri Lanka. Twenty-four cages, each having 2 m<sup>3</sup> water capacities (size 1 m W × 2 m L × 1.5 m H), made of 2 mm high density polyethylene nets, were used. Twenty-four cages were randomly divided in to six groups to have four cages for each treatment. A total of 3600 GIFT fingerlings having average initial weights (2.94 ± 1.47 g) were stocked at a density of (75 fish/m<sup>3</sup>). The diets with three dietary Crude Protein (CP) levels (26%, 30% and 36%) and two dietary Crude Fat (CF) levels (7% and 12%) were used and the experiment was designed according to 3 × 2 factorial designs.

The six diets were prepared by locally available feed ingredients. Ingredient percentage, cost and proximate composition of the six diets are given in Table 1. The experimental fish in each cage fed at 10% biomass of fish at initial month and daily feeding percentage altered accordingly monthly mean weight gain using standards described by Lim and Webster [3].

Fish were fed for 150 days. Monthly samples from each cage (30%) were taken randomly. Individual body weights were recorded to the nearest 0.01 g by using an electronic balance (Digital Scale A&D).

Final Mean Weight (FMW), Net Yield (NY), Specific Growth Rate (SGR), Feed Conversion Ratio (FCR) and Protein Efficiency Ratio (PER) were calculated as described by Abdel-Tawwab [19] and Survival Rate (SR) also calculated at the end of the study.

$$NY = \text{Gross yield (g)} - \text{Initial total fish weight (g)}$$

$$SGR (\% \text{ day}^{-1}) = 100 \times (\text{Ln Final mean weight} - \text{Ln Initial mean weight}) / \text{Culture period (days)}$$

$$FCR = \text{Dry feed given (g)} / \text{Wet weight gain (g)}$$

$$PER = \text{Increment in body weight (g)} / \text{Protein intake (g)}$$

$$SR = 100 \times (\text{Number of fish harvested} / \text{Number of fish stocked})$$

At the end of the study three fish from each sex from each cage were collected and a total of 24 fish per treatment were pooled as a composite sample for proximate analysis. Six fish were randomly collected from this composite sample and subjected to proximate analysis using standard procedures [20].

Normally distributed data were subjected to Two way Analysis of Variance and Student Newman-Keuls test to find how the growth indices vary in fish fed with six different diets. Effect of three protein levels and two fat levels on growth were separately analysed by One way Analysis of Variance using SPSS 16 statistical software. Data are reported as mean ± standard error and the significance level was assessed at (P≤0.05).


## Results

The growth performance and feed utilization of GIFT fingerling under different levels of dietary protein and fat are given in the Table 2. Changing protein level from 26% to 36% resulted significant differences (P≤0.05) in growth performance indices; final mean weight, specific growth rate, net yield and feed conversion ratio (Table 2). The highest values for FMW, SGR, NY and better FCR were observed in fish fed with 36% CP diets (Table 3). There was no significant difference

Parameter	Mean values for each CP level			Mean values for each fat levels	
	26% CP	30% CP	36% CP	7% CF	12% CF
FMW	110.35 ± 4.36 <sup>a</sup>	115.44 ± 4.88 <sup>a</sup>	162.29 ± 3.9 <sup>b</sup>	128.84 ± 6.71 <sup>a</sup>	129.88 ± 8.88 <sup>a</sup>
SGR (% day <sup>-1</sup> )	2.40 ± 0.02 <sup>a</sup>	2.44 ± 0.02 <sup>a</sup>	2.66 ± 0.01 <sup>b</sup>	2.50 ± 0.03 <sup>a</sup>	2.50 ± 0.04 <sup>a</sup>
FCR	1.63 ± 0.05 <sup>a</sup>	1.58 ± 0.04 <sup>a</sup>	1.18 ± 0.03 <sup>b</sup>	1.46 ± 0.05 <sup>a</sup>	1.46 ± 0.07 <sup>a</sup>
PER	2.36 ± 0.08 <sup>a</sup>	2.11 ± 0.06 <sup>a</sup>	2.43 ± 0.15 <sup>a</sup>	2.25 ± 0.07 <sup>a</sup>	2.35 ± 0.11 <sup>a</sup>
NY (Kg)	12.8 ± 4.6 <sup>a</sup>	13.4 ± 4.21 <sup>a</sup>	18.6 ± 4.62 <sup>b</sup>	14.9 ± 7.44 <sup>a</sup>	14.90 ± 9.59 <sup>a</sup>

FMW – Final Mean Weight, SGR – Specific Growth Rate, FCR – Feed Conversion Ratio, PER – Protein Efficiency Ratio, NY – Net Yield, CP- Crude Protein, CF- Crude Fat  
<sup>a</sup>Values in the same row having different letters are significantly different (P≤0.05)

**Table 3:** Pooled value (mean ± SE, n = 8 for crude protein levels and n =12 for crude fat levels)<sup>1</sup> for FMW, SGR, FCR, PER and NY of GIFT during 150 days cage culture period.



Experimental diets CP/ CF	Fillet composition			
	CP	CF	Ash	DM
26/7	16.66 ± 0.13 <sup>a</sup>	2.36 ± 0.01 <sup>a</sup>	1.07 ± 0.04 <sup>ab</sup>	20.14 ± 0.2 <sup>a</sup>
26/12	16.16 ± 0.73 <sup>a</sup>	3.18 ± 0.47 <sup>b</sup>	1.11 ± 0.07 <sup>ab</sup>	20.51 ± 0.33 <sup>a</sup>
30/7	18.62 ± 0.22 <sup>bc</sup>	2.26 ± 0.13 <sup>a</sup>	0.99 ± 0.06 <sup>a</sup>	21.89 ± 0.15 <sup>b</sup>
30/12	20.15 ± 0.02 <sup>cd</sup>	2.42 ± 0.02 <sup>a</sup>	1.19 ± 0.10 <sup>ab</sup>	23.79 ± 0.14 <sup>c</sup>
36/7	17.89 ± 1.02 <sup>ab</sup>	3.44 ± 0.15 <sup>b</sup>	1.15 ± 0.10 <sup>ab</sup>	22.33 ± 0.87 <sup>b</sup>
36/12	20.56 ± 0.16 <sup>d</sup>	4.09 ± 0.10 <sup>c</sup>	1.34 ± 0.01 <sup>b</sup>	26.17 ± 0.20 <sup>d</sup>

CP – Crude Protein, CF – Crude Fat, DM – Dry Matter.  
<sup>1</sup>Values in the same column having different letters are significantly different (P≤0.05).

**Table 4:** Proximate composition of fish fillet (mean ± SE, n=4)<sup>1</sup> according to diets.

in final mean weight and other growth indices between protein levels, 26% to 30%.

No significant difference (p>0.05) in growth indices between fish fed with 7% and 12% CF having same protein level diets indicates that increasing fat levels have no marked contribution on growth performance indices of GIFT at any protein level (Tables 2 and 3). Fish fed with the diet 36/12 shows marked growth when they reached an average weight above 130 g which is not prominently seen in lower fat diet (36/7) (Figure 1).

Fillet of fish fed with 30% and 36% had higher protein content irrespective of the fat content than fish fed with 26% CP diet. No significant difference was observed in protein content of fish fillets of fish fed with 30% and 36% CP diets (Table 4). High fat deposition in fish fillets was obtained in 12% CF in 26% and 36% protein diets.

Protein efficiency ratio was not significantly different among treatments. Survival rates were 86.33% to 90% among six treatments.

## Discussion

Tilapia is known as herbivorous and omnivorous fish species feed on variety of natural food and artificial feeds [21]. Improving growth and feed efficiency with increased protein level is well known in fish [7]. Studies have shown that tilapia attain maximum growth for a range of 20% to 56% CP in diet El-Dahhar, Zeweil [6]. *Oreochromis niloticus* fry having weight of 0.8 g and 40 g need respectively 40% and 30% minimum amount of crude protein levels in diet for maximum growth [22]. The most cost effective dietary protein level lay between 25% to 28% [3]. These values are mostly taken by measuring the growth response of fish to quality graded protein (purified or semi-purified) [3], instead of commercially available cost effective feed ingredients. Present study demonstrates that increasing dietary protein level by using cheap (Table 1), easily available feed ingredients has positive effect on growth of GIFT in cage culture.

Most people prefer to consume larger tilapia while, higher

prizes claimed for large fish have made them not affordable. This has constrained them to eat smaller fish (<150 g). As such cost has overcome the consumer preference [23]. In the present study the GIFT fish attained a maximum size of 160 g within 150 days in cages is more favorable in terms of cost benefit to the consumers.

Dietary protein requirement of young Tilapia (*O. mossambicus*, *O. niloticus*, *O. aureus* and *Tilapia zilli*) often range from 34% to 36% in diet for maximum growth potential. Similarly, Bahnasawy [24] observed that mono-sex Nile tilapia reared in fertilized tanks reached significantly higher growth performance by different protein levels (17% CP to 35%) however no significant growth increase was observed between 30% and 35% CP diets. Present study also demonstrated that GIFT fed with 26 and 30% CP diet had no significant difference in growth performance revealing that dietary feed protein level within the optimal protein range makes no significant difference in growth of fish.

Abdel-Tawwab [19], Ahmad [25] reported that FCR decrease when protein level increased in diets. Similar effect was observed in the present study when CP level increased to 36% suggesting that requirement of high protein diet to maintain an efficient FCR. FCR observed between 26% and 30% CP diets was not significantly different. These results are in agreement with Bahnasawy [24], who revealed that there is no significant difference found for FCR when mono-sex tilapia when fed with 25, 30 and 35% CP.

In the present study GIFT has shown reduction in PER when CP level increase from 26% to 30%. Reduction of PER with increasing dietary CP level have been reported in many studies for different tilapia species [25-27]. This is mainly due to conversion of protein to energy when fish fed with high level of protein in diet [28]. Dietary energy supplementation through inclusion of high lipid levels proportionate to protein levels in diets may have promoted the efficient utilization of crude protein. Considering the PER and the cost of feed formulation 26% and 30% CP level diet can be considered as the most cost effective CP levels for the grow out phase of GIFT cage culture.

Diet containing 10-15% CF demonstrated good growth and feed utilization for hybrid tilapia [29]. They suggested that hybrid tilapia needs 5% CF for minimum growth and 12% CF for maximum growth in diet. Lim, Webster [3] has shown that tilapia does not tolerate as high CF in diet as carnivorous fish species and the growth can be depressed. As the growth performance and feed utilization are not significantly different in two fat levels in the present study, 7% lipid containing diet could be used for the growth of GIFT raised in cages.

High lipid diets have shown more beneficial effects for fish growth closer to the harvesting period when compared to low lipid level diets, when proper protein level is provided [30,31]. Increases of growth at the fourth month of the grow out phase in the fish fed with 36% CP and 12% CF (Figure 1) indicate that ability to utilize lipids more effectively, closer to maturity of GIFT.

Synthesis and deposition rate of protein and fat content in fish body could be related to muscle function [32,33] and the composition of diet. High fat level diets (12% CF) have not increased the fat content significantly among fish filets except in 36/12 CF diet and minimal contribution to the protein sparing of the fish (Table 4).

## Conclusion

GIFT in cages fed with elevated protein levels (36%) produce best growth performance. Fat levels (7 and 12%) has no significant difference effect on GIFT growth performance and feed utilization. Increased fat

levels (12%) have not contribute to protein sparing of fish too. As such, 36/7 diet is suitable for grow out phase and 36/12 for fattening stage of the GIFT in cages.

## Acknowledgment

Financial assistance by Transforming University of Ruhuna in to International Status - 2012 Grant (RU/DVC/RRO/143) is gratefully acknowledged.

## References

1. Fotedar R (2004) Effect of dietary protein and lipid source on the growth, survival, condition indices, and body composition of marron, *Cherax tenuimanus* (Smith). *Aquaculture* 230: 439-455.
2. Lavell RT (1989) Nutrition and feeding of fish. Springer.
3. Lim C, Webster CD (2006) Tilapia: Biology, culture, and nutrition. Taylor & Francis.
4. Parazo MM (1990) Effect of dietary protein and energy level on growth, protein utilization and carcass composition of rabbit-fish, *Siganus guttatus*. *Aquaculture* 86: 41-49.
5. Kim KD, Sang GL, Yong JK, Kim KW, Son MH, et al. (2012) Effects of dietary protein and lipid levels on growth and body composition of juvenile far eastern catfish *Silurus asotus*. *Asian-Australasian journal of animal sciences* 25: 369.
6. El-Dahhar A, Zeweil H, El-Tawil N (2000) Effect of protein and energy levels in commercial diets on growth performance of juvenile Nile Tilapia (*Oreochromis niloticus*). *Egyptian J. of Aquatic Biology and Fishries* 4: 267-285.
7. NRC (1993) Nutrient requirements of fish. National Research Council. Committee on Animal Nutrition.
8. Prather EE, Lovell RT (1973) Response of intensively fed channel catfish to diets containing various protein-energy ratios. *Game Fish Comm.*
9. Phillips AM (1972) Calorie and energy requirement, in fish nutrition. Academic Press New York, NY.
10. Cowey CB (1980) Protein and amino acid requirements of fin-fish in European Aquaculture Society Symposium Protein Metabolism and Nutrition. Hamburg: EAAP Publication.
11. Ringrose RC (1971) Calorie-to-energy ratio for brook trout (*Salvelinus fontinalis*). *Journal of the Fisheries Research Board of Canada* 28: 113-117.
12. El Sayed M, Garling L, Abd-El AS (1988) Protein and energy requirements of (*Tilapia zilli*) fingerlings. *Alexandria Journal of Veterinary Science*.
13. Cho C, Kaushik S (1990) Nutritional energetics in fish: energy and protein utilization in rainbow trout (*Salmo gairdneri*). *World review of nutrition and dietetics* 61: 132-172.
14. McGoogan BB, Gatlin DM (1999) Dietary manipulations affecting growth and nitrogenous waste production of red drum, *Sciaenops ocellatus* L. Effects of dietary protein and energy levels. *Aquaculture* 178: 333-348.
15. Page JW, Andrews JW (1973) Interactions of dietary levels of protein and energy on channel catfish (*Ictalurus punctatus*). *The Journal of Nutrition* 103: 1339-1346.
16. El-Dahhar A, Lovell R (1995) Effect of protein to energy ratio in purified diets on growth performance, feed utilization and body composition of Mozambique tilapia, *Oreochromis mossambicus* (Peters). *Aquaculture Research* 26: 451-457.
17. Jishun T (1998) Effect of different dietary protein and oil levels in diet on growth of tilapia: Comparison in of aquaculture grade (South American) and locally produced fish meal. *International Fishmeal & Oil Manufacturers Association*.
18. Dey MM, Eknath AE, Li S, Hussain MG, Tran MT, et al. (2000) Performance and nature of genetically improved farmed tilapia: a bioeconomic analysis. *Aquaculture Economics & Management* 4: 83-106.
19. Abdel-Tawwab M (2010) Effect of dietary protein level, initial body weight, and their interaction on the growth, feed utilization, and physiological alterations of Nile tilapia, *Oreochromis niloticus* (L.). *Aquaculture* 298: 267-274.
20. AOAC (1995) Official methods of analysis, (16th edn), 1995: Association of Official Analytical Chemists, Arlington, VA, USA.
21. Chowdhury K (2006) Effect of salinity on carrying capacity of adult Nile tilapia *Oreochromis niloticus* L. in recirculating systems. *Aquaculture Research* 37: 1627-1635.

22. Siddiqui AQ, Howlader MS, Adam AA (1988) Effects of dietary protein levels on growth, feed conversion and protein utilization in fry and young Nile tilapia, *Oreochromis niloticus*. Aquaculture 70: 63-73.
23. Murray FJ, Little DC (2000) Inland fisheries resources and the current status of aquaculture in Sri Lanka.
24. Bahnasawy MH (2009) Effect of dietary protein levels on growth performance and body composition of monosex Nile tilapia, *Oreochromis niloticus* L. reared in fertilized tanks. Pakistan Journal of Nutrition 8: 674-678.
25. Ahmad MH, Abdel-Tawwab M, Khattab YA (2004) Effect of dietary protein levels on growth performance and protein utilization in Nile tilapia (*Oreochromis niloticus*) with different initial body weights. The sixth international symposium on tilapia in aquaculture, Manila, Philippine.
26. Abdel-Tawwab M (2012) Effects of dietary protein levels and rearing density on growth performance and stress response of Nile tilapia, *Oreochromis niloticus* (L.). International Aquatic Research 4: 1-13.
27. Jauncey K (1982) The effects of varying dietary protein level on the growth, food conversion, protein utilization and body composition of juvenile tilapias (*Sarotherodon mossambicus*). Aquaculture 27: 43-54.
28. Kim KI, Kayes TB, Amundson CH (1991) Purified diet development and re-evaluation of the dietary protein requirement of fingerling rainbow trout (*Oncorhynchus mykiss*). Aquaculture 96: 57-67.
29. Chou BS, Shiau SY (1996) Optimal dietary lipid level for growth of juvenile hybrid tilapia, *Oreochromis niloticus* × *Oreochromis aureus*. Aquaculture 143: 185-195.
30. Kim LO, Lee SM (2005) Effects of the dietary protein and lipid levels on growth and body composition of bagrid catfish, *Pseudobagrus fulvidraco*. Aquaculture 243: 323-329.
31. Ahmad MH (2008) Response of African catfish, *Clarias gariepinus*, to different dietary protein and lipid levels in practical diets. Journal of the World Aquaculture Society 39: 541-548.
32. Fauconneau B (1984) The measurement of whole body protein synthesis in larval and juvenile carp (*Cyprinus carpio*). Comparative Biochemistry and Physiology Part B: Comparative Biochemistry 78: 845-850.
33. Abdel-Tawwab M, Khattab YAE, Ahmad MH, Adel MES (2006) Compensatory growth, feed utilization, whole-body composition, and hematological changes in starved juvenile Nile Tilapia, *Oreochromis niloticus* (L.). Journal of Applied Aquaculture 18: 17-36.

**Citation:** De Silva MPKSK, Senaarachchi WARK, Liyanage NPP (2016) Title-Combinatory Effects of Diets with Three Protein Levels and Two Fat Levels on Growth Performance and Fillet Composition of Cage Cultured Genetically Improved Farmed Tilapia (GIFT). J Aquac Res Development S2: 008. doi:[10.4172/2155-9546.S2-008](https://doi.org/10.4172/2155-9546.S2-008)

This article was originally published in a special issue, **Current and Emerging Diseases/Disorders of Fish in Aquaculture** handled by Editor(s). Prof. Patrick T.K. Woo, University of Guelph, Canada; Dr. Kenneth D. Cain, University of Idaho, USA.

### OMICS International: Publication Benefits & Features

#### Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

#### Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submit/>