

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.



ISSN: 1736-4337

Journal of Generalized Lie Theory and Applications

**The International Open Access
Journal of Generalized Lie Theory and Applications**

Editors-in-Chief

Eugen Paal

Tallinn University of Technology, Estonia

Alexander A Voronov

University of Minnesota, USA

Executive Editors

Wieslaw A Dudek

Wroclaw University of Technology, Poland

Alice Fialowski

Eotvos Lorand University, Hungary

Available online at: OMICS Publishing Group (www.omicsonline.org)

This article was originally published in a journal by OMICS Publishing Group, and the attached copy is provided by OMICS Publishing Group for the author's benefit and for the benefit of the author's institution, for commercial/research/educational use including without limitation use in instruction at your institution, sending it to specific colleagues that you know, and providing a copy to your institution's administrator.

All other uses, reproduction and distribution, including without limitation commercial reprints, selling or licensing copies or access, or posting on open internet sites, your personal or institution's website or repository, are requested to cite properly.

Digital Object Identifier: <http://dx.doi.org/10.4172/1736-4337.1000207>

To the Solution for Beal Conjecture

Victor P Baryshok*

Cathedra of Chemical Technology, Irkutsk State Technical University, Str. Lermontova 83, Irkutsk, 664074, Russia

Abstract

Found general expressions for infinite series of equations $A^x + B^y = C^z$, where A, B, C, x, y and z are positive integers and x, y and z are all greater than 2, (A), (B) and (C) have a Common Prime Factor. However, these expressions also hold true for fractional and negative values of x, y, and z, and cover the range of numbers from the infinitely small to the infinitely large.

Keywords: Generalized equations; $A^x + B^y = C^z$ for Beal Conjecture

Andrew Beal has formulated a Conjecture in number theory [1]:

$$\text{If } A^x + B^y = C^z \quad (1)$$

where A, B, C, x, y and z are positive integers and x, y and z are all greater than 2, then A, B and C must have a common prime factor. Is it appropriate to seek a systematic solution for the sum of two numbers in the assumption that A and B have even numbers. Minimum an even number of A (and B) is 2. Then the equation (1) with a Common Prime Factor 2 can be written as

$$2^x + 2^y = 2^z \quad (2a),$$

or for different A, B, C:

$$(2^n)^x + (2^m)^y = 2^z \quad (2b)$$

If $mx = nx$, then

$$A^x + B^y = 2 \cdot (2^n)^x; \quad C^z = 2^{nx+1} \quad (2c)$$

By the condition of Beal Conjecture are x, y and z are positive integers and all are greater than 2. In order to be agree with this condition for a different positive integers A, B and C, minimum value for $nx(=my)$ should be 23 and increase by 2^2 , i.e.

$$nx(=my) = 2^2(2+k), \quad (3)$$

where k is a positive integer from continuous integer row, including zero (k= 0, 1,2, 3...).

Then the expression (2b) satisfying Beal Conjecture, may be represented by expression (4):

$$(2^n)^x + (2^m)^y = 2^{nx+1}, \quad (4)$$

where $nx(=my) = 2^2(2+k)$; n is equal or unequal to m; n and m are equal or unequal 1; or (4a):

$$2^{3+k} + 2^{3+k} = 2^{4+k}, \quad (4a)$$

where k is any rational number.

Note that in the case k = 0, for differing A, B and C there is only one solution eq. 4, where $C \neq 2$ and $z \neq (nx+1)$:

$$k = 0, n = 1, x = 8, m = 2, y = 4, C = 8, z = 3 \quad (28 + 44 = 83).$$

Similarly, the known individual equations in which A, B and C have a Common Prime Factor [1]:

$$3^3 + 6^3 = 3^5 \Rightarrow 3^3 + (2 \cdot 3) = 3^5 \quad (5)$$

$$7^6 + 7^7 = 98^3 \Rightarrow 7^6 + 7^7 = (2 \cdot 7^2)^3 \quad (6)$$

$$33^5 + 66^5 = 33^6 \Rightarrow (3 \cdot 11)^5 + (6 \cdot 11)^5 = (3 \cdot 11)^6 \quad (7)$$

$$34^5 + 51^4 = 85^4 \Rightarrow (2 \cdot 17)^5 + (3 \cdot 17)^5 = (5 \cdot 17)^4 \quad (8)$$

$$19^4 + 38^3 = 57^3 \Rightarrow 19^4 + (2 \cdot 19)^3 = (3 \cdot 19)^3, \quad (9)$$

were reviewed and the common expressions of equations, satisfying the conjecture of Beal's infinite series, are obtained. The exception is a family of equations with Common Prime Factor 5, $y = 2$ (eq. 11).

$$3^{3(k+1)} + (2 \cdot 3^{k+1})^3 = 3^{3k+5}, \quad (10)$$

$$5^{2(k+1)} + (2 \cdot 5^{k+1})^2 = 5^{2k+3} \quad (11)$$

$$7^{3(k+2)} + 7^{3k+7} = (2 \cdot 7^{k+2})^3, \quad (12)$$

$$(3 \cdot 11^{6k+1})^5 + (6 \cdot 11^{6k+1}) = (3 \cdot 11^{5k+1})^6 \Rightarrow 33^{5(k+1)} + (2 \cdot 33^{k+1}) = 33^{5k+6}, \quad (13)$$

$$(2 \cdot 17^{4k+1})^5 + (3 \cdot 17^{5k+1})^4 = (5 \cdot 17^{5k+1})^4, \quad (14)$$

$$19^{3k+4} + (2 \cdot 19^{4k+1})^3 = (3 \cdot 19^{4k+1})^3, \quad (15)$$

In equations (10-15) k-value takes any rational number.

Other well-known equations [1], having a Common Prime Factor not represented in eqs. 5-9, are special cases of common expressions. For Example:

$$2^2 + 2^3 = 2^4 \quad (\text{eq. 4: } n, m=1, x=3, y=3 \text{ or eq. 4a: } k=0),$$

$$2^9 + 8^3 = 4^5 \Rightarrow 2^9 + (2^3)^3 = (2^2)^5 \quad (\text{eq. 4: } n=1, m=3, x=9, y=3 \text{ or eq. 4a: } k=6),$$

$$3^9 + 54^3 = 3^{11} \Rightarrow 3^9 + (2 \cdot 3^3)^3 = 3 \cdot 11 \quad (\text{eq. 10, } k=2),$$

$$27^4 + 162^3 = 9^7 \Rightarrow (3^3)^4 + (2 \cdot 3^3)^3 = 3^{14} \quad (\text{eq. 10, } k=3).$$

Using negative or fractional values k can get arithmetic addition of prime numbers. For example, if in eq. 11 take $k = -1$, the equation would be simplified to $1+4 = 5$. If $k = -2$ in equation 12, the result will be $1+7 = 8$. This may be of interest to number theory.

Thus, there are infinite rows of equations $A^x + B^y = C^z$, where A, B, C are positive integers and have a Common Prime Factor. At the

*Corresponding author: Victor P Baryshok, Cathedra of Chemical Technology, Irkutsk State Technical University, Str. Lermontova 83, Irkutsk, 664074, Russia, Tel: +7(3952)405-513; E-mail: baryvik@yandex.ru

Received November 21, 2013; Accepted November 24, 2013; Published November 28, 2013

Citation: Baryshok VP (2014) To the Solution for Beal Conjecture. J Generalized Lie Theory Appl 8: 207. doi: 10.4172/1736-4337.1000207

Copyright: © 2014 Baryshok VP. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

same time, x , y , and z are positive integers, and more 2, only private representations of generalized expressions, that occupy region of numbers to positive infinity.

Reference

1. R. Daniel Mauldin (1997) The Beal Conjecture, Notices of the Amer. Math. Soc., 44: 1436-1438.