

## Towards a More Sustainable Production of Fish as an Important Protein Source for Human Nutrition

Sabine Sampels\*

Faculty of Fisheries and Protection of Waters, University of South Bohemia in Ceske Budejovice, South Bohemian Research Center of Aquaculture and Biodiversity of Hydrocenoses, Institute of Aquaculture, Husova tř. 458/102, 370 05 České Budějovice, Czech Republic

Fish and seafood products, including aquatic plants are not only tasty constituents of meals and food products, but they also have a high nutritional valuable and contain beneficial amounts of protein, lipids as well as essential minerals and vitamins. Nowadays, aquaculture is the fastest growing sector in livestock production around the world. An annual increase of production during last 30 years is  $\pm 8\%$  [1]. In a recent review it has also been shown that fish and seafood have a major impact on global food supply [2]. Already today, the internationally traded amount of aquatic animal food products is higher than the total traded amount of beef, pork and poultry combined (26.85-27.45 compared to 20.38-21.99 million tons in 2009 respectively) [3,4]. In addition in several Asian and African countries the major animal protein source are aquatic animals [3,5].

Generally, aquatic animal foods have higher protein content than most terrestrial meats. In addition aquatic protein is highly digestible and rich in several essential amino acids that are limited in terrestrial meat proteins, as for example methionine (6.5% compared to 5.7% of total essential amino acids in fish compared to terrestrial meat respectively) and lysine (19.6% compared to 19.0% of total essential amino acids in fish compared to terrestrial meat respectively) [2]. Fish and seafood consumption has been connected to beneficial health effects, especially with the prevention of cancer, decreased risk of coronary heart and cardiovascular diseases, decreased inflammatory disease as arthritis. Historically the main effects of fish consumption have been attributed to the high content of long chain omega 3 (n-3) polyunsaturated fatty acids (PUFA). However it gets more and more clear that also other nutrients from fish have positive effects on human health. Beside the n-3 PUFA, fish and other seafood are a significant source of a well-balanced amino acid composition, taurine and choline, the Vitamins D and B12 as well as calcium, phosphorus, iodine and selenium. Depending on the general nutrition of a population, fish and seafood also might provide significant proportions of Vitamin A, iron and zinc. Another important aspect is that fish in general has a significantly lower feed conversion rate (FCR) than land living animals. For example beef cattle can have a FCR varying from 5-20 (National Research Council 2000) [6] while Masilko [7] reported for example a FCR for carp from 2.08-2.62 while a FCR from 1.6-1.8 was reported for tilapia [8] and salmonids usually have a FCR at around 1 [9]. This means more protein can be produced with a lower amount of feed with all included factors concerning feed production as energy and water use as well.

This combination makes fish an interesting object of research with a focus on an economical and easy as well as sustainable production in order to secure protein supply for human diet in the future. There has been some discussion however about the sustainability of aquaculture, where the main issues are the high use of fish meal and fish oil as well as the leakage of nutrients to the environment leading to an eutrophication of aquatic systems. There has been some research going on about substituting the more and more scarce and expensive fish-meal and fish oil from the fish feeds for carnivore fishes [10]. A second object towards a more sustainable production has been to produce feeds with a reduced FCR. For example it was shown that a simple pre-treatment of the used feed as for example pressing of cereals could reduce the FCR in carp as well as reduce the leakage of phosphorus to the pond water

[7]. In combination with other techniques and systems as for example aquaponics, where vegetables or flowers are grown in the nutrient rich flow out water from aquaculture recirculation systems [11]. The plants act as a bio filter and clean the water from excess nitrogen and other nutrients and provide additional food in form of vegetables or herbs. These systems also decrease the load of phosphorus, nitrogen and other nutrients from the waste water and have hence a positive effect on the environment [12]. Aquaponic systems can function already on a small base and could be hence a good solution for small or family businesses in rural areas. Therefore research should also focus on establishing simple, easy to operate and sustainable aquaculture/aquaponic systems, which can function with a minimum of technical support, and which have a low proportion of waste water. There is no doubt that aquaculture has still a big potential to grow and to be a major contributor to feed the world of tomorrow.

### Acknowledgments

This work was financially supported by the Ministry of Education, Youth and Sports of the Czech Republic - projects "CENAKVA" (No. CZ.1.05/2.1.00/01.0024), "CENAKVA II" (No. LO1205 under the NPU I program) and GAJU 074/2013/Z.

### References

1. FAO (2012) The State of World Fisheries and Aquaculture. Rome, Italy.
2. Tacon AGJ, Metian M (2013) Fish Matters: Importance of Aquatic Foods in Human Nutrition and Global Food Supply. *Reviews in Fisheries Science* 21: 22-38.
3. FAO/FISHSTAT (2012) FishstatJ a tool for fishery statistical analysis, release 2.0.0. Global capture fisheries production 1950-2010, Global aquaculture production 1950-2010, Global commodities production and trade 1978-2009.
4. FAO Fisheries Department, Data and Statistics Unit. Rome, FAO, FAS/USDA (2012). Livestock and poultry: World markets and trade. USDA, Foreign Agricultural Service.
5. FAO (2001) Food balance sheets-A Handbook. Rome, FAO Economic and Social Development Department: 99.
6. National Research Council (2000) Nutrient Requirements of Beef Cattle: Seventh Revised Edition: Update 2000. Washington, DC, The National Academies Press.
7. Masilko J (2014) Production efficiency of technological modified cereals in market carp farming. Faculty of Fisheries and Protection of waters. Vodňany, University of South Bohemia in Ceske Budejovice. PhD: 101.
8. LEAD (2006) Livestock's long shadow - environmental issues and options. Rome, Italy, FAO.

\*Corresponding author: Sabine Sampels, Faculty of Fisheries and Protection of Waters, University of South Bohemia in Ceske Budejovice, South Bohemian Research Center of Aquaculture and Biodiversity of Hydrocenoses, Institute of Aquaculture, Husova tř. 458/102, 370 05 České Budějovice, Czech Republic, Tel: +420 387 77 46; E-mail: [sampels@frov.jcu.cz](mailto:sampels@frov.jcu.cz)

Received October 01, 2014; Accepted October 04, 2014; Published October 14, 2014

Citation: Sampels S (2014) Towards a More Sustainable Production of Fish as an Important Protein Source for Human Nutrition. *J Fisheries Livest Prod* 2: 119. doi: [10.4172/2332-2608.1000119](http://dx.doi.org/10.4172/2332-2608.1000119)

Copyright: © 2014 Sampels S. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

9. Nordgarden UF, Oppedal F, Taranger GL, Hemre GI, Hansen T (2003) Seasonally changing metabolism in Atlantic salmon (*Salmo salar* L.) I – Growth and feed conversion ratio. *Aquaculture Nutrition* 9: 287-293.
10. Naylor RL, Hardy RW, Bureau DP, Chiu A, Elliott M, et al. (2009) Feeding aquaculture in an era of finite resources. *Proc Natl Acad Sci USA* 106: 15103-15110.
11. Trang NTD, Brix H (2014) Use of planted biofilters in integrated recirculating aquaculture-hydroponics systems in the Mekong Delta, Vietnam. *Aquaculture Research* 45: 460-469.
12. Turcios AE, Papenbrock J (2014) Sustainable Treatment of Aquaculture Effluents-What Can We Learn from the Past for the Future? *Sustainability* 6: 836-856.

**Citation:** Sampels S (2014) Towards a More Sustainable Production of Fish as an Important Protein Source for Human Nutrition. J Fisheries Livest Prod 2: 119. doi: [10.4172/2332-2608.1000119](https://doi.org/10.4172/2332-2608.1000119)

### Submit your next manuscript and get advantages of OMICS Group submissions

#### Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

#### Special features:

- 350 Open Access Journals
- 30,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission/>

