

## Traumatic Brain Injury

Ninh Doan<sup>\*</sup>, Mohit Patel, Hayley Doan, Karl Janich, Ha S Nguyen and Saman Shabani

Department of Neurosurgery, Medical College of Wisconsin, Milwaukee, Wisconsin, USA

<sup>\*</sup>Corresponding author: Ninh Doan, Department of Neurosurgery, Medical College of Wisconsin, Milwaukee, Wisconsin, 53226, USA, Tel: 1414-955-8296; E-mail: [ndoan@mcw.edu](mailto:ndoan@mcw.edu)

Received date: October 31, 2016; Accepted date: November 08, 2016; Published date: November 14, 2016

Copyright: © 2015 Doan N, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### Introduction

Traumatic brain injury (TBI) remains a significant problem in both the pediatric and adult populations. Approximately 500,000-700,000 pediatric TBI incidents and 2.5 million adults are suffered from TBI every year [1-4]. TBI is ranged from mild to severe and its management depends of the severity of the injury. The injuries from TBI can be classified into primary and secondary injury. Primary injury, which is considered irreversible but preventable, is the injury took place at the time of trauma such as the direct damage to the parenchyma of the brain, producing axonal shear and rupture. On the other hand, the secondary injury defines as the injury occurring following the initial TBI, ranging from inflammation that can increase intracranial pressure (ICP), which in turn causing further damages. Therefore, the goal of most medical management after TBI is to decrease the insult caused by secondary injury with the aim of maximizing the patient's recovery from the primary insult [5]. Similar treatments are also employed in pediatric patients with TBI. Preventing of secondary injury is even more important pediatric population given their brains still have significant potential for plasticity [6].

Concussion is one of the most common pediatric injuries. Its symptoms vary depending on the given situation and can include cognitive challenges, sleep disturbances, headaches, and ataxia. Headache is one of the most disabling symptoms that patients often complain about after a concussion [7]. Children often need a longer rehabilitation time post-concussion compared to adults [8,9].

In the case of trauma patients with a suspicion of TBI, usually computed tomography (CT) is the imaging of choice given its high sensitivity for blood and the short duration of scanning. However, in certain mild TBI cases imaging may not be needed. The classically used "Canadian CT Head Rule" is a useful guide to follow to help deciding the need for a CT of the head in the setting of mild TBI [10].

The coagulation panel that includes Normalized Ratio or PT/INR, and activated Partial Thromboplastin Time or aPTT) should be sent on admission. A prompt correction of a patient's INR to 1.3 or lower has been shown to decrease hematoma expansion and improves outcomes [11]. While platelet transfusion is recommended for thrombocytopenia (platelet count < 100,000) for patients with intracranial hemorrhage, others have found that its use to counter the effect of antiplatelet agents, such as aspirin or clopidogrel, may not improve outcome [12]. Novel new agents for reversal of newer anticoagulants, such as direct thrombin inhibitors or Factor Xa inhibitors are promising, such as idarucizumab for dabigatran reversal [13] or andexanet alfa for reversal of anti-Factor Xa agents [14].

Monitoring of ICP is recommended for all TBI patients without clinically followable neurological exams [15] or whose GCS < 8 with an

abnormal CT scan of the head [16]. Most institutions use the external ventricular drain (EVD) or the fiber optic intraparenchymal probe to record ICP. Due to the loss of cerebral blood flow autoregulation at high ICP, which can lead to hypoperfusion of the brain, ICP must be kept below 20 mmHg, commonly through the employment of sedation and hypertonic therapy [17].

Decompressive craniectomy is often employed as a last resort to treat medically intractable ICP [18-21]. It has been showed in the DECRA trial that improvement ICP control and shorter stay in ICU in patients treated with the bilateral frontotemporoparietal craniectomy, but this also led to more unfavorable outcomes based on the extended glasgow outcome scale (GOS), which was believed to be secondary to the improvement in the survival of vegetative patients [22]. In consistent with this, a recent randomized study called RESCUEicp demonstrated that decompressive craniectomy in patients with TBI resulted in a lower mortality rate at 6 months compared to medical treatment alone [23,24].

### References

- Centers for Disease Control and Prevention (2015) Report to Congress on Traumatic Brain Injury in the United States: Epidemiology and Rehabilitation. National Center for Injury Prevention and Control, Division of Unintentional Injury Prevention, Atlanta, GA.
- Dewan MC, Mummareddy N, Wellons JC, Bonfield CM (2016) Epidemiology of Global Pediatric Traumatic Brain Injury: Qualitative Review. *World Neurosurg* 91: 497-509.
- Li L, Liu J (2013) The effect of pediatric traumatic brain injury on behavioral outcomes: a systematic review. *Dev Med Child Neurol* 55: 37-45.
- Wilde EA, Hunter, Bigler ED (2012) Pediatric traumatic brain injury: neuroimaging and neurorehabilitation outcome. *NeuroRehabilitation* 31: 245-260.
- Karl JH, Patel M, Shabani S, Montoure A, Doan N (2016) Management of Adult Traumatic Brain Injury: A Review. *J Trauma Treat* 5:320.
- Doan N, Patel M, Nguyen HS, Montoure A, Shabani S, et al. (2016) A rare remarkable recovery in a pediatric patient with the bi-hemispheric, transventricular trajectory craniocerebral gunshot wound. *J Surg Case Rep* 5: 1-3.
- Blume HK (2015) Headaches after Concussion in Pediatrics: a Review. *Curr Pain Headache Rep* 19: 42.
- Simma B, Lütschg J, Callahan JM (2013) Mild head injury in pediatrics: algorithms for management in the ED and in young athletes. *Am J Emerg Med* 31: 1133-1138.
- Barlow KM, Crawford S, Stevenson A, Sandhu SS, Belanger F, et al. (2010) Epidemiology of postconcussion syndrome in pediatric mild traumatic brain injury. *Pediatrics* 126: e374-381.
- Stiell IG, Wells GA, Vandemheen K, Clement C, Lesiuk H, et al. (2001) The Canadian CT Head Rule for patients with minor head injury. *Lancet* 357: 1391-1396.
- Huttner HB, Schellinger PD, Hartmann M, Köhrmann M, Juettler E, et al. (2006) Hematoma growth and outcome in treated neurocritical care

- patients with intracerebral hemorrhage related to oral anticoagulant therapy: comparison of acute treatment strategies using vitamin K, fresh frozen plasma, and prothrombin complex concentrates. *Stroke* 37: 1465-1470.
12. Hemphill JC, Greenberg SM, Anderson CS, Becker K, Bendok BR, et al. (2015) Guidelines for the Management of Spontaneous Intracerebral Hemorrhage: A Guideline for Healthcare Professionals from the American Heart Association/American Stroke Association. *Stroke* 46: 2032-2060.
  13. Pollack CV, Reilly PA, Eikelboom J, Glund S, Verhamme P, et al. (2015) Idarucizumab for Dabigatran Reversal. *N Engl J Med* 373: 511-520.
  14. Siegal DM, Curnutte JT, Connolly SJ, Lu G, Conley PB, et al. (2015) Andexanet Alfa for the Reversal of Factor Xa Inhibitor Activity. *N Engl J Med* 373: 2413-2424.
  15. Chesnut RM, Temkin N, Carney N, Dikmen S, Rondina C, et al. (2012) A trial of intracranial-pressure monitoring in traumatic brain injury. *N Engl J Med* 367: 2471-2481.
  16. Bratton SL, Chestnut RM, Ghajar J, McConnell Hammond FF, Harris OA, et al. (2007) Guidelines for the management of severe traumatic brain injury. VI. Indications for intracranial pressure monitoring. *J Neurotrauma* Suppl 1: S37-44.
  17. Czosnyka M, Smielewski P, Piechnik S, Steiner LA, Pickard JD (2001) Cerebral autoregulation following head injury. *J Neurosurg* 95: 756-763.
  18. Bullock MR, Chesnut R, Ghajar J, Gordon D, Hartl R, et al. (2006) Surgical management of acute subdural hematomas. *Neurosurgery* 58: S16-24.
  19. Bullock MR, Chesnut R, Ghajar J, Gordon D, Hartl R, et al. (2006) Surgical management of acute epidural hematomas. *Neurosurgery* 58: S7-15.
  20. Bullock MR, Chesnut R, Ghajar J, David G, Roger H, et al. (2006) Surgical management of traumatic parenchymal lesions. *Neurosurgery* 58: S25-46.
  21. Bullock MR, Chesnut R, Ghajar J, Gordon D, Hartl R, et al. (2006) Surgical management of posterior fossa mass lesions. *Neurosurgery* 58: S47-55.
  22. Chi JH (2011) Craniectomy for traumatic brain injury: results from the DECRA trial. *Neurosurgery* 68: N19-20.
  23. Hutchinson PJ, Koliass AG, Timofeev IS, Corteen EA, Czosnyka M, et al. (2016) Trial of Decompressive Craniectomy for Traumatic Intracranial Hypertension. *N Engl J Med* 375: 1119-1130.
  24. Appelboom G, Zoller SD, Piazza MA, Szpalski C, Bruce SS, et al. (2011) Traumatic brain injury in pediatric patients: evidence for the effectiveness of decompressive surgery. *Neurosurg Focus* 31: E5.