

Two Cases of Chronic Myelomonocytic Leukemia Combined with Monoclonal Gammopathy of Undetermined Significance and a Literature Review

Fengling M¹, Lijia Z¹, Wei Z¹, Xiaohui G¹, Lina Z¹ and Jianyong L^{2*}

¹Department of Hematology, The First People's Hospital, Yangzhou, Jiangsu, China

²The First Affiliated Hospital of Nanjing Medical University, Jiangsu Province Hospital, China

Abstract

Objective: To describe myelodysplastic syndrome (MDS)/ myeloproliferative neoplasm (MPN) combined with monoclonal gammopathy of undetermined significance (MGUS) in order to investigate the potential association between these 2 diseases.

Methods: Two cases of confirmed chronic myelomonocytic leukemia (CMML) combined with MGUS were reported here. In addition, reports related to cases with combined MDS or MPN with MGUS were reviewed.

Results: The first case was a 77-year-old man whose routine blood tests showed abnormal hemogram results. The diagnosis was CMML combined with IgM monoclonal gammopathy, and the disease course was 4 years. The CMML gradually progressed, and the patient presented with anemia, thrombocytopenia, autoimmune hemolysis, and an increase in the number of immature cells in the bone marrow. Although the MGUS caused fluctuations in the concentrations of IgM, no IgM-associated organ damage was observed. Eventually, this patient died of lung infection. The second case was a 78-year-old man who sought treatment because of fever and a cough. An increase in the number of monocytes was discovered in the peripheral blood. Bone marrow smear results suggested obvious active granulocytes and an increase in the percentages of promyelocytes, myelocytes, and metamyelocytes. Unhealthy granulocytes and immature monocytes could also be observed, and the percentage of monocytes was increased. In addition, serum IgG levels were increased, and immunofixation electrophoresis results showed IgG-k type M proteins. The diagnosis was CMML combined with IgG monoclonal gammopathy. These diseases were stable, and follow-up was conducted for 1 year after diagnosis.

Conclusion: The cases in this study combined with those that were reviewed from the relevant literature indicated that the presence of the above 2 tumors in 1 patient might not be a coincidence. The development of the 2 diseases in case 1 was different, and it was speculated that they might have had different clonal origins. Whether CMML may serve as a risk factor for MGUS as well as the function of clonal plasma cells in the occurrence and development of MDS or MDS/MPN still requires further studies on a large number of clinical cases.

Keywords: Bone marrow; Myelomonocytic; Leukemia; Monoclonal; Gammopathy

Introduction

Chronic myelomonocytic leukemia (CMML) constitutes a group of clonal diseases that originate from committed hematopoietic myeloid stem cells or from pluripotent stem cells. Monoclonal gammopathy of undetermined significance (MGUS) is a common pre-malignant plasma cell disease. These diseases are both common neoplastic diseases in elderly people. Although these 2 neoplastic diseases originate from different types of committed hematopoietic stem cells, case reports of patients with these 2 diseases are not rare. Do these 2 diseases present together coincidentally or are they correlated? The current report describes 2 cases of CMML combined with MGUS and presents a review of the relevant literature.

Case Report

Case 1

The patient in this case was a 77-year-old man. He was admitted to the hospital on July 15th, 2011 because of an "increase in white blood cells for more than 1 year and gum bleeding for 3 days". The patient first sought treatment at a local health care institution in October 2010 because of a "cold". Routine blood tests revealed white blood cell (WBC): $11.4 \times 10^9/L$, hemoglobin (Hb): 132 g/L and platelet (PLT): $71 \times 10^9/L$. The patient did not seem to pay attention. After admission,

physical examination demonstrated that the patient was lucid and was in good spirits. No signs of anemia were present, the skin featured scattered petechiae and ecchymosis, there were no palpable superficial lymph nodes, and no abnormalities were found in the heart, lungs, or abdomen. The full abdominal CT showed mild fatty liver, a cyst on the right kidney, and some pelvic fluid. Routine blood tests revealed WBC: $34.7 \times 10^9/L$ with a differential count of 51.9% neutrophils, 10.8% lymphocytes, and 36.6% monocytes, Hb: 108 g/L, PLT: $26 \times 10^9/L$. Blood smear showed metamyelocytes: 0.01%, rod neutrophils: 0.05, lobed neutrophils: 0.60, lymphocytes: 0.18, and monocytes: 0.16. Moreover, small clusters of platelets, but no eosinophils or basophils, were observed. Complete immunologic and serologic tests showed IgA: 1.9 g/L, IgG: 7.59 g/L, IgM: 48.44 g/L, IgE: 15 IU/ml, and the erythrocyte

***Corresponding author:** Jianyong L, The First Affiliated Hospital of Nanjing Medical University, Jiangsu Province Hospital, China, Tel: 862583714511; E-mail: gaoqx@lzu.edu.cn

Received May 17, 2016; **Accepted** September 24, 2016; **Published** September 30, 2016

Citation: Fengling M, Lijia Z, Wei Z, Xiaohui G, Lina Z, et al. (2016) Two Cases of Chronic Myelomonocytic Leukemia Combined with Monoclonal Gammopathy of Undetermined Significance and a Literature Review. Med Rep Case Stud 1: 122. doi: 10.4172/2572-5130.1000122

Copyright: © 2016 Fengling M, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

sedimentation rate (ESR) was 90 mm/hr. Blood biochemistry showed globulin: 48.1 g/L, total bilirubin concentration: 24.3 $\mu\text{mol/L}$, indirect bilirubin: 20.9 $\mu\text{mol/L}$, creatinine: 135 $\mu\text{mol/L}$, uric acid: 452 $\mu\text{mol/L}$, and the electrolyte concentrations were normal. Serum protein electrophoresis revealed the presence of immunoglobulin M protein. The bone marrow smear showed clear active proliferation of nucleated cells and an increased granulocyte/erythrocyte ratio. The percentage of cells of the granulocyte lineage was 62%, of which myeloblasts, promyelocytes, myelocytes, metamyelocytes, rod neutrophils, lobed neutrophils, and eosinophils accounted for 0%, 1.2%, 12.4%, 13%, 21%, 12.4%, and 1.6%, respectively. Monocytes accounted for 12.4%, and out of these, immature monocytes accounted for 6.4%, while mature monocytes accounted for 6%. A total of 16 megakaryocytes appeared on the whole smear, and the distribution of platelets was scattered. The percentage of lymphocytes was 13.4%. No abnormal cell morphology was observed. Iron staining showed that extracellular iron staining was positive (+). The intracellular iron positivity rate was 21%, the neutrophil alkaline phosphatase (NAP) positivity rate was 5%, and the NAP score was 5 points (the reference score in our laboratory was 7-63 points). The bone marrow immunology typing showed that 0.7% of the immature cell population was of the myeloid lineage; in addition, the 77% of cells of the granulocyte lineage had maturation defects. A bone marrow biopsy showed active bone marrow hyperplasia, active granulocyte lineage proliferation, scattered primitive immature cells, and an increased percentage of mature granulocytes. In addition, increased cells of the plasma cell lineage were observed on histologic sections, and some regions contained aggregated clusters. Quantitative polymerase chain reaction (qPCR) results for the BCR-ABL fusion gene were negative, tests for the JAK2V617F gene mutation were negative, and the karyotype of the chromosomes was normal. The diagnosis was 1. CMML combined with MGUS, 2. chronic renal insufficiency, and 3. hyperuricemia. To alleviate his symptoms, the patient received treatment that included hemostasis, anti-inflammation, and renal protection; after the symptoms improved, he was released from the hospital. After that, the patient was admitted several more times because of recurrent dizziness and fatigue, coughing, asthma, and combined gum bleeding. The results of a second exam showed that the patient's renal function was stable. The numbers of white blood cells and monocytes were increased, and the patient was anemic. The results of routine blood tests from February 2013 showed WBC: $10.57 \times 10^9/\text{L}$ with a differential count of 28.9% neutrophils, 20.3% lymphocytes, and 47.8% monocytes, Hb: 69 g/L, and PLT: $31 \times 10^9/\text{L}$. Blood biochemistry showed globulin: 54.9 g/L, total bilirubin level: 52.1 $\mu\text{mol/L}$, and indirect bilirubin level: 49.3 $\mu\text{mol/L}$. Coombs test was positive. The level of IgM was 68.51 g/L. The results from the bone marrow puncture smear showed active proliferation. In addition, the percentage of immature monocytes increased to 28%; in this group, the cells each had a larger cell body with an irregular shape, the cytoplasm was abundant, the nuclei were irregular, the nuclear chromatin was fine and granular, and most cells did not have a nucleolus. Mature monocytes accounted for 10.5% of the cells. Proliferation of cells of the granulocyte lineage was clear. Myeloblasts, promyelocytes, myelocytes, metamyelocytes, rod neutrophils, and lobed neutrophils accounted for 10%, 5%, 2.5%, 1.5%, 7.5%, and 22.5%, respectively. Abnormalities such as nuclear atypia and a decrease in the number of cytoplasmic granules could also be observed. Proliferation of cells of the erythrocyte lineage was decreased; the number of lymphocytes was also decreased and accounted for 7.5% of the cells. Thirty-nine megakaryocytes were present on the whole section, and the number of platelets was decreased. A thoracic CT revealed inflammation in

the superior lobe of the left lung and in the inferior lobe of the right lung as well as emphysema. After the patient received oral hydroxyurea and prednisone as well as treatment for his symptoms (anti-infection, kidney protection, activating blood circulation to dissipate blood stasis, blood perfusion, and improvement of myocardial blood supply), the patient's condition improved. The patient ultimately died in December 2014 of lung infection combined with cardiopulmonary insufficiency.

Case 2

The patient was a 78-year-old man. He was admitted to the hospital on February 9, 2014 because of "coughing for 10 days and fever for 2 days". Ten days before admission, the patient caught a cold and experienced a cough. He was treated with "azithromycin and levofloxacin". The effects of these drugs were poor, and the patient then experienced fatigue and a fever of 39.4°C. Physical examination showed that the patient was lucid and in poor spirits. There were no obvious signs of anemia, both lungs exhibited coarse breath sounds and did not exhibit dry or wet rales; the heart rate was 90 beats/min, and atrial fibrillation without a murmur was detected. After admission, a thoracic X-ray showed bronchitis. Electrocardiography showed rapid atrial fibrillation, complete right bundle branch block, and partial lead ST-T changes. Routine blood test revealed WBC: $7.93 \times 10^9/\text{L}$ with a differential count of 58.9% neutrophils, 10.2% lymphocytes, and 29.6% monocytes, red blood cells (RBCs): $3.2 \times 10^{12}/\text{L}$, Hb: 108 g/L, and PLT: $137 \times 10^9/\text{L}$. Blood biochemistry results showed that the albumin, globulin, creatinine, and uric acid levels were 34.6 g/L, 44.5 g/L, 122 $\mu\text{mol/L}$, and 492 $\mu\text{mol/L}$, respectively. The ESR was 101 mm/h. Blood smear showed that the percentage of mature monocytes was significantly increased and accounted for 36% of the white blood cells. The ferritin concentration was 1203 ng/ml, and tests for the detection of serum folate and vitamin B12 revealed normal levels. Bone marrow smear showed active bone marrow hyperplasia, proliferation of cells of the granulocyte lineage, and an increase in the percentages of promyelocytes, myelocytes, metamyelocytes, and monocytes. Additionally, unhealthy granulocytes could be observed; 0.4% of the total cells were immature monocytes, while 17.2% were mature monocytes. Complete serology showed IgA: 0.23 g/L, IgG: 27.8 g/L, and IgM: 0.26 g/L. Immunofixation electrophoresis showed the presence of IgG- κ type M proteins; M proteins accounted for 28.2%, the level of free light chain κ : 9.31 mg/L (range: 3.3-19.4 mg/L), λ : 11.7 mg/L (range: 5.7-26.3 mg/L), and κ/λ : 0.79 (range: 0.26-1.65). The patient had a history of "hypertension" for more than 20 years and a history of "coronary heart disease and atrial fibrillation". He had received a pacemaker implantation in the heart and had a history of gallstone surgery. A review of the patient's several routine blood tests from 2012 and 2013 showed that the patient had a normal white blood cell count and an increased percentage of monocytes, which fluctuated between 15-35%. The final diagnosis included the following: 1. CMML combined with MGUS, 2. bronchitis, and 3. hyperuricemia. Plasma cells develop from B lymphocytes (B cells), a type of white blood cell that is made in the bone marrow. Normally, when bacteria or viruses enter the body, some of the B cells will change into plasma cells. The plasma cells make antibodies to fight bacteria and viruses, to stop infection and disease. Plasma cell neoplasms are diseases in which abnormal plasma cells or myeloma cells form tumors in the bones or soft tissues of the body. The plasma cells also make an antibody protein, called M protein that is not needed by the body and does not help fight infection. These antibody proteins build up in the bone marrow and can cause the blood to thicken or can damage the kidneys.

The patient also received anti-infection treatment, which was

ineffective after 5 days, as the patient still exhibited a high fever, poor appetite, and fatigue. The patient was then treated with an intravenous injection of methylprednisolone, and his body temperature returned to normal. However, the patient again exhibited signs of a fever after 4 days. After he received cyclophosphamide combined with methylprednisolone, the symptoms of fever and fatigue gradually disappeared. Treatment with these drugs was stopped after 2 months. The clinical follow-up was performed until recently; the disease was stable, and the patient did not complain of any discomfort. Routine blood tests, complete immunology tests, and immunofixation electrophoresis were performed regularly, and the results did not show any significant changes.

Cyclophosphamide is usually used in conjunction with steroids (prednisone). It frequently helps decrease the doses of steroids that are necessary to control vasculitis. In this regard, cyclophosphamide is sometimes referred to as a “steroid-sparing agent”. Cyclophosphamide may cause infertility in both men and women who are treated with the medication. As with many of cyclophosphamide’s side-effects, the risk of infertility appears to be related to the length of time a patient receives the drug (and probably the cumulative dose, as well). In a long-term study of the effectiveness of cyclophosphamide in treating lupus, the risk of infertility among women of childbearing potential was 57%.

Discussion

MDS is composed of a group of clonal diseases that have the following clinical presentation: ineffective hematopoiesis in the bone marrow and a high-risk of transformation into acute myeloid leukemia. Before 2008, CMML had already been classified as a subtype of MDS. In recent years, the World Health Organization (WHO) decided that the French-American-British (FAB) classification system did not adequately reflect the clinical prognosis and biological behaviors of patients; therefore, CMML was classified into an independent, larger disease group (MDS/MPN). In fact, according to the new classification system, CMML includes previous MDS-CMML (more emphasis on dysmyelopoiesis) and MPN-CMML (more emphasis on bone marrow proliferative changes) [1]. MGUS is a common pre-malignant plasma cell disease. MGUS may be divided into 3 different subtypes, as follows: non-IgM type MGUS, IgM type MGUS, and light chain type MGUS. Non-IgM type MGUS is further divided into IgG, IgA, double clone, and the rare IgD and IgE types [2]. All types typically evolve into multiple myeloma (MM), AL amyloidosis, or other diseases at a percentage of 1% every year.

Although the diseases mentioned above originate from different types of committed hematopoietic stem cells, case reports on the presence of these 2 diseases in a single patient are not rare. In 1986, the Spanish researcher Costa [3] first reported 1 case of CMML combined with lymphoma. Since then, researchers from different countries also successively reported cases of CMML combined with lymphoma [4,5]. A prospective study by Economopoulos [6] showed that 5 out of 61 patients with CMML had combined MGUS; the percentage was 8.2% and was much higher than the 1-3% in the general population. However, Cesana [7] studied disease transformation of 1231 patients with either MGUS or smoldering MM and showed that only 0.4% of cases had combined MPN. This leads to the question of whether MGUS combined with MDS or MDS/MPN are coincidental or whether these 2 diseases are closely associated. Mailankody et al. [8] retrospectively analyzed secondary hematological malignancies in 8740 cases of MM and 5652 cases of MGUS that occurred in Sweden between 1986 and 2005. In cases of MGUS, the risk of development of MPN/polycythemia

vera (PV) was increased 5-fold. Compared with MGUS, the risk of development of AML/MDS in patients with MM was even higher, and patients with the IgG/IgA type were more prone to the development of acute myeloid leukemia (AML)/MDS than were patients with IgM type gammopathy. However, the incidence of MPN/PV was not increased significantly in patients with MM. These results were not consistent with those of a study with a small sample size conducted by Duhrsen et al. [9], which showed that the number of cases of plasma cell tumors combined with MPN/PV was significantly increased. The authors admitted that the high risk of incidence obtained in the study might have been due to a certain detection bias. For example, the clinical data of patients with MGUS were obtained from the Swedish Inpatient Registry; therefore, they could not account for some patients who sought treatment for other diseases and who were diagnosed with MGUS after they underwent MGUS-related examinations. In addition, routine blood tests during the clinical follow-up of patients with MGUS might help in the discovery of MDS at an early stage. To exclude bias associated with targeted examinations for other diseases, the Mayo Clinic [10] performed a survey on local healthy residents above the age of 50 years. The results showed that the development of MDS in patients with MGUS increased 2.4-fold compared with the control group; moreover, no patients with MGUS developed acute lymphoid leukemia (ALL). These results were consistent with the results of the Swedish report. Because the number of patients with AML in the cohort study was small, the results could not confirm whether MGUS was associated with AML. Compared with the Swedish study, which showed that the development of MDS in patients with MGUS was 8.1 times that in the controls, the number was significantly decreased. The researchers considered that the reason for this was that the subjects who were enrolled in the Swedish study were mostly patients who had clinical symptoms and intentionally went to the hospital for a serum protein electrophoresis examination; therefore, some patients who did not have clinical symptoms were missed. The findings of the Mayo Clinic study, which differed from those of the Swedish study, did not reveal that patients with IgA/IgM MGUS were prone to the development of MDS or AML. Although conclusions from these 2 large studies had some differences, they both found that the incidence of MDS in patients with plasma cell diseases was significantly increased compared with that in the general healthy population.

Do the 2 types of hematological tumors in the same individual arise from the same stem cells or do they have different cell origins? The observation by Shvidel et al. [11] was that the disease progression of CMML was different from the progression of lymphoma. Therefore, they considered that these tumor types did not arise from a common abnormal stem cell population. Zagaria [12] reported 1 case of MGUS in a patient who developed anemia and thrombocytosis after 2 years. The patient was diagnosed with MDS combined with a 5q deletion. From the first diagnosis of MGUS to the development of MDS, the presence of a MYD88 mutation was discovered in the bone marrow. To clarify the origins of these 2 clones, allele-specific PCR (AS-PCR) combined with fluorescence in situ hybridization (FISH) was used to detect the molecular cytogenetic changes in hematopoietic stem cells. The results showed that CD34+ hematopoietic stem cells all had the 5q deletion; however, no mutations in the MYD88 gene were observed. At the cytogenetic and molecular levels, the results showed that these 2 tumors were independent of each other. Anelli [13] also reported that patients with polycythemia vera with the JAK2 V617F mutation developed IgG type lymphoplasmacytic lymphoma (LPL) after several years. The detection of the MYD88 L265P mutation and the JAK2 V617F mutation confirmed that these 2 diseases were two clonal diseases that co-existed

in the same person. However, some studies also reported that leukemia cells in patients with CMML expressed both monocyte and NK cell markers together; therefore, it was speculated that these tumor cells might have originated from common progenitor cells of monocytes and NK cells [14]. During treatment for a nose bleed following a cold, it was discovered that the first patient (case 1) in the current study had CMML combined with IgM MGUS. The disease course lasted more than 4 years, and the CMML gradually progressed; the clinical presentation included anemia, thrombocytopenia, autoimmune hemolysis, and an increase in immature cells in the bone marrow. Although MGUS was associated with fluctuations in IgM values, no IgM-associated organ damage was observed. For example, lymphadenopathy, hyperviscosity, hypercalcemia, and kidney and bone damage did not significantly worsen during disease transformation compared with when the patient was first diagnosed. These results also suggested that these tumors had different clonal origins; if 2 types of tumor clones were present, CMML might have been the dominant clone. Because MGUS and MDS or MDS/MPN primarily occur in elderly people who typically do not exhibit clinical symptoms, MGUS is usually discovered incidentally when patients seek treatment for other diseases. The diagnosis of MDS or MPN/MDS is usually confirmed because of the presence of clinical symptoms induced by abnormal blood cells. Therefore, when a patient presents with these 2 diseases simultaneously, it is difficult to know which disease developed first. Malhotra et al. [15] retrospectively analyzed 90 patients with MPN who were treated at their medical center within the past 5 years. Immunofixation electrophoresis results from the blood and urine of a total of 32 patients were examined. Out of these, 15 were diagnosed with MGUS/MM; however, the majority of patients was diagnosed with MPN at the same time or before the MGUS/MM was diagnosed, and only 3 patients were diagnosed first with MGUS/MM followed by MPN. None of these 3 cases featured an increase in the number of red blood cells or platelets, which indicates that the diagnosis was delayed relative to the actual course of disease development. The 2 elderly patients described in this article were discovered to have both CMML and MGUS during treatment for other diseases; therefore, no treatment-related factors were involved. Because immunoglobulin detection was not performed initially, this test could not confirm which disease developed first.

Studies have shown that monocytes and nurse-like cells derived from monocytes are important components of the tumor microenvironment that play important roles in the maintenance and protection of tumor cell growth in the blood [16]. Whether CMML may serve as a risk factor of MGUS and the function of clonal plasma cells in the occurrence and development of MDS or MDS/MPN still requires further studies. Yoshida et al. [17] discovered 14 cases of MDS combined with MGUS from their medical center within the past decade; this article is currently the clinical report with the largest number of cases of MDS combined with MGUS. These cases accounted for 10.2% of MDS patients within the same period, which was significantly higher than the number of people of the same age in the control population with anemia. When the patients were diagnosed with MDS, the majority of them were also diagnosed with MGUS, which suggests that MGUS was not associated with treatment and that MGUS could be present in patients with any MDS subtype according to the WHO classification. Among the 12 cases, 11 cases had low-risk chromosome karyotypes; combined with other risk factors, the clinical outcomes of the MDS combined with MGUS group were not significantly different from those in the MDS group, which suggests that MGUS did not influence the prognosis of the patients with MDS. Case 1 in this article had a 4-year-long disease course. The patient's disease did not transform into acute leukemia,

and he eventually died from a lung infection. Compared with the median survival period of 20-40 months of patients with CMML without MGUS, the survival period was not decreased; therefore, this observation supported the idea that MGUS might not have an adverse effect on the prognosis of patients with CMML.

The elderly are susceptible to both MGUS and MDS/MPN. After a review of the relevant literature, a limited number of studies showed that the incidence of secondary MDS/MPN in patients with MGUS was higher than that in the general population. The association between these 2 diseases and the mutual influence on each of their prognoses still require further studies of a large number of clinical cases. In addition, these results suggested that in future work, routine blood tests should be the focus for patients with MGUS. Finally, an analysis of serum M protein should not be ignored in patients with MDS/MPN.

References

1. Hebeda KM, Fend F (2009) Changed concepts and definitions of myeloproliferative neoplasms (MPN), myelodysplastic syndromes (MDS) and myelodysplastic/myeloproliferative neoplasms (MDS/MPN) in the updated 2008 WHO classification. *J Hematop* 2: 205-210.
2. Korde N, Kristinsson SY, Landgren O (2011) Monoclonal gammopathy of undetermined significance (MGUS) and smoldering multiple myeloma (SMM): novel biological insights and development of early treatment strategies. *Blood* 117: 5573-5581.
3. Costa B, Llorente A, Vidal F (1986) Coexistence of chronic myelomonocytic leukemia, lymphoplasmacytoid lymphoma and lymph node leishmaniasis. *Med Clin (Barc)* 87: 73-75.
4. Cervero M, Obeso G, Marco J, Fernández-Ballesteros A, Martín-Joven A (1994) Pleomorphic T – lymphoma associated with chronic myelomonocytic leukemia. *Sangre (Barc)* 39: 61.
5. Gaulier A, Jary-Bourguignat L, Serna R, Pulik M, Davi F, et al. (2000) Occurrence of angioimmunoblastic T cell lymphoma in a patient with chronic myelomonocytic leukemia features. *Leuk Lymphoma* 40: 197-204.
6. Economopoulos T, Economidou J, Papageorgiou E, Dervenoulas J, Christodoulides C, et al. (1989) Monoclonal gammopathy in chronic myeloproliferative disorders. *Blut* 58: 7-9.
7. Cesana C, Klersy C, Barbarano L, Nosari AM, Crugnola M, et al. (2002) Prognostic factors for malignant transformation in monoclonal gammopathy of undetermined significance and smoldering multiple myeloma. *J Clin Oncol* 20: 1625-1634.
8. Mailankody S, Pfeiffer RM, Kristinsson SY, Korde N, Bjorkholm M, et al. (2011) Risk of acute myeloid leukemia and myelodysplastic syndromes following multiple myeloma and its precursor disease (MGUS). *Blood* 118: 4086-4092.
9. Duhren U, Uppenkamp M, Meusers P, König E, Brittinger G (1988) Frequent association of idiopathic myelofibrosis with plasma cell dyscrasias. *Blut* 56: 97-102.
10. Roeker LE, Larson DR, Kyle RA, Kumar S, Dispenzieri A (2013) Risk of acute leukemia and myelodysplastic syndromes in patients with monoclonal gammopathy of undetermined significance (MGUS): a population-based study of 17315 patients. *Leukemia* 27: 1391-1393.
11. Shvidel L, Sigler E (1995) Chronic myelomonocytic leukemia preceding high-grade immunoblastic lymphoma. *Eur J Haematol* 54: 270.
12. Zagaria A, Coccaro N, Tota G, Anelli L, Minervini A, et al. (2015) Myelodysplastic syndrome with 5q deletion following IgM monoclonal gammopathy, showing gene mutation MYD88 L265P. *Blood Cells Mol Dis* 54: 51-52.
13. Anelli L, Zagaria A, Minervini A, Casieri P, Coccaro N, et al. (2014) IgG-lymphoplasmacytic lymphoma following polycythemia vera: JAK2 V617F and MYD88 L265P mutations separated in the same house. *Ann Hematol* 93: 1605-1607.
14. Kojima H, Bai A, Mukai HY, Hori M, Komeno T, et al. (2000) Chronic myelomonocytic leukemia derived from a possible common progenitor of monocytes and natural killer cells. *Leuk Lymphoma* 37: 617-621.
15. Malhotra J, Kremyanskaya M, Schorr E, Hoffman R, Mascarenhas J (2014) Coexistence of myeloproliferative neoplasm and plasma-cell dyscrasia. *Clin Lymphoma Myeloma Leuk* 14: 31-36.

16. Filip AA, Cisel B, Koczkodaj D, Wąsik-Szczepanek E, Piersiak T, et al. (2013) Circulating microenvironment of CLL: are nurse-like cells related to tumor-associated macrophages. *Blood Cells Mol Dis* 50: 263-270.
17. Yoshida Y, Oguma S, Ohno H, Nakabo Y, Katsurada T, et al. (2014) Co-occurrence of monoclonal gammopathy and myelodysplasia: a retrospective study of fourteen cases. *Int J Hematol* 99: 721-725.