

Type One Diabetic Patients Knowledge and Attitude on Insulin Self Administration in Mekele Tigray, Ethiopia

Hadgu Gerensea¹, Admasu Moges², Birikity Shumiyee², Frweiyne Abrha², Medina Yesuf², Teweldde Birihan², Tsiyon Birhanu² and Zomanesh Getahun²

¹School of Nursing, Axum University, Axum, Ethiopia

²Axum University, Referral Hospital, Ethiopia

*Corresponding author: Hadgu Gerensea, School of Nursing, Axum University, Axum, Ethiopia, Tel: +251-914 -38- 8918; E-mail: hadguellen1@gmail.com

Rec date: August 11, 2015; Acc date: February 05, 2016; Pub date: February 12, 2016

Copyright: © 2016 Gerensea. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Introduction: Insulin is a valuable drug for those who are in need of it. Insulin will have tremendous impact, when used properly. Intensive insulin therapy was essential in the maintenance of strict glycemic control among insulin requiring patients with diabetes. However this presents challenges in the face of the complexities associated with insulin use and also taking in the consideration the potential danger associated with inappropriate use. Insufficient knowledge and attitude of insulin self administration can result in preventable complication, adverse patient outcome, poor adherence to therapy and invariably poor glycemic control.

Objectives: To assess knowledge and attitude on insulin self administration among type 1 DM patients in Mekele hospital 2015.

Methodology: Institution based descriptive cross sectional study was conducted. The study subject were type 1 DM patients in Mekele hospital (attending chronic follow up). Data was collected by interview using pre-tested structured questionnaires. The data was entered and analyzed using SPSS version 16. All type one DM patients were included in the study.

Result: Among 141 respondents, more than half 78 (55.3%) of them had average knowledge and majority 96 (68.0%) of the participants had favorable attitude.

Conclusion and recommendation: The finding of this showed that type 1 diabetic patient had average knowledge and favorable attitude towards self administration of insulin injection. It is suggested that more should be done on areas like medias, health professionals on health education towards insulin self administration for type 1 DM patients.

Keywords: Knowledge; Attitude; Insulin self administration

Introduction

Diabetes is a public health problem which is a chronic disease and is now as an epidemic in both developed and developing countries. DM is recognized as one of the leading cause of death and disability worldwide. The physical, social and economic factors involved in the management of diabetes are a continuous strain for the health sector as well as to government agencies while there has been a great improvement in the knowledge, epidemiology and management of this condition in the developed world, there has been little or no improvement in sub-Saharan Africa including Ethiopia [1].

Insulin is one of the harmful medications if used inappropriately. For this reason, it is identified as one of the five "high alert" medications that have the greatest risk of causing injury to patients because of medication errors [2,3].

Intensive insulin therapy is essential in the maintenance of strict glycemic control among insulin requiring patients with diabetes. However this presents a challenge in the face of the complexities associated with insulin use and also taking into consideration the potential dangers associated with inappropriate use. Insufficient

knowledge of insulin self injection can result in preventable complication, adverse patient outcome, full adherence to therapy and invariably poor glycemic control [4]. Many insulin requiring diabetes patients possess poor knowledge of insulin use and are conversant with some important terms such as ketoacidosis, insulin reaction hypoglycemia [5].

It is a growing concern worldwide; while there has been a great improvement in the knowledge, epidemiology and management of this condition in the developed world, there has been little or no improvement in sub-Saharan Africa including Ethiopia [1].

Knowledge about insulin self administration is a fully requested for individuals and communities to take action for control of diabetes. The treatment for type 1 diabetes (IDDM) is injectable insulin therapy which requires coordination and understanding of both the individuals with diabetes and those responsible for diabetic care. The diabetes patients who are on insulin need to be knowledgeable regarding the disease and insulin therapy and also they must have a competency and positive attitude towards self administration of insulin injection to overcome the barriers of insulin injection and to have good glycemic control. Patients need knowledge to improve the competency level of self administration of insulin which in turn will help in better glycemic

control and avoid complications. Glycemic control and prevention of complication will reduce the cost of treatment and diabetes related mortality [6].

The long term prognosis of type 1DM is said to improve with a complex demanding and often intensive regimen and maintenance of lower blood glucose level is said to reduce the risk of long term complication by as much as 60% [7].

Many studies conducted elsewhere in the world on insulin therapy have showed that the knowledge and awareness about the disease can have positive influence on knowledge and attitude of patients regarding Insulin Self-administered that could lead to better management of diabetes and eventually good quality of life. However, knowledge and attitude gap exists in type 1 diabetes mellitus management that does not allow patients and healthcare professionals to implement life style changes could reduce the morbidity and mortality associated with diabetes.

Diabetes is one of the major causes of premature illness and death worldwide. More than 80% of diabetes death occurs in low and middle- income countries. In our study area, Mekele hospital health professionals alone can't provide high quality care.

Diabetes being a chronic illness requires sound knowledge and attitude of self insulin administration by patients, so that they can contribute meaningfully in the management of their lives. However there was no enough study conducted on knowledge and attitude of insulin self administration among type 1 DM patients.

The study has been conducted to assess the knowledge and attitude of insulin self administration with type 1 DM patients in Mekele hospitals. The findings of this study will help in identifying population knowledge gap and their attitude towards insulin self administration which might help in designing effective educational program for safe insulin self administration and also will be used as a base line in the future for further study.

Objectives

General objectives

To assess knowledge and attitude on insulin self administration among type one diabetic patients in Mekele hospital, Tigray, Ethiopia 2015.

Specific objectives

- To determine knowledge on insulin self administration among type 1 diabetic patients.
- To explore attitude of type 1 diabetic patients on insulin self administration.

Methodology

Study area

The study was conducted in Mekele hospital, which is a teaching and regional hospital located in Mekele town, North Ethiopia, at 783 km away from Addis Ababa, the capital city of Ethiopia. Mekele is one of the biggest cities in Tigray national regional state with a population more than 150341. It is a member of governmental health institution;

Study design and period

Institution based cross sectional study design was used to assess knowledge and attitude on self- administration of insulin from January to August 4, 2015.

Population

Source population: The source populations for this study were all type 1 diabetic patients who have follow up in Mekele hospital.

Study population: The study populations for this study were all type 1 diabetes patients who came to Mekele hospital for follow up during the study period.

Inclusion and exclusion criteria

Inclusion criteria: All type 1 diabetic patient attending at Mekele hospital chronic follow up clinic were included in this study

Exclusion criteria: Type 1 diabetes patients who were severely ill, age less than 18 year and psychiatric patients were excluded from the study.

Sampling size determination

All type one DM who has follow up on Mekele hospital was included in the study. A total of 141 were taken from 155 types one DM with 90% response rate.

Study variable

Dependant variable:

- Knowledge of insulin self administration
- Attitude of insulin self administration

Independent variable: Age, Sex, Marital status, Education, Religion, Health profile related variables, Duration of treatment, DM association member ship and family history of diabetics were independent variables.

Data collection method

Interview with structured closed ended questionnaire was performed by revising questionnaires developed from similar study, which was conducted in Mangalore and adapted to the objectives of the present study to collect quantitative data. The questionnaire was designed in English and translated to Tigrigna (local language) for better understanding of the study participants and again translated back to English by other person to check validity.

Data quality assurance

The prepared questionnaire was pre-tested on 15 type 1 DM patients in shire, suhul hospital, then some modification were made before applied on the study subject.

Data processing and analysis

Data processing, analysis and interpretation have been done using SPSS. The frequency and percentage of all variables was determined.

Operational definition

Good knowledge-a subject who answered 9-13 (above 66%) correct responses from knowledge questions.

Average knowledge-a subject who answered 5-8 (33.3-66.6%) response from knowledge question.

Poor knowledge-a study participant who answered 0-4 (below 33.3%) corrects response from knowledge question

Attitude-these are feelings and beliefs expressed by patients towards self administration and are evident from score obtained by their response in validated rating scale according to their attitude response the study participants were categorized as 6 and above (>49%) favorable attitude and below 6 (50%) unfavorable attitude.

Insulin self administration –the injection of insulin expected to be administered at home without assistance. This includes preparation of article, withdrawal of injection, and administration after care.

Ethical consideration

Ethical clearance was obtained from Aksum University College of health sciences department of nursing and permission obtained from Mekele hospital administrative office. Informed verbal consent was obtained from each study subjects after explanations of the importance of the study. They had all the right not to be involved in the study or not to answer any question. We were also informed that all of their response was confidential.

Result

Socio demographic characteristics of study subjects

Total of 141 study participants were interviewed using structured questionnaires giving response rate of 90%.

From the total sample size population, almost half 68 (48.1%) of the respondents were between the age group of 18-31 years followed by 31-40 years and 41-50 years which were 34 (24.11%) and 22 (15.6%) respectively. More than half 72 (51%) of them were male. The majority 99 (70.2%) of the participants were orthodox followers. sixty four (45.3%) Subjects were married and only 8 (5.6%) were widowed. Regarding the educational status of study subjects, majority of them 67 (47.5%) was primary school (Table 1).

Variables		Frequency (%)
Age in year	18-30	68 (48.1)
	31-40	34 (24.11)
	41-50	24 (15.5)
	51-60	12 (8.5)
	>60	5 (3.5)
Sex	Male	72 (51)
	Female	69 (49)
	Orthodox	99 (70.2)
	Muslim	30 (21.27)
Religion	Protestant	12 (8.5)

	Other	-
	Married	64 (45.39)
	Single	52.6 (37.58)
Marital Status	Divorced	17 (12)
	Widowed	7 (4.9)
	Illiterate	15 (10.6)
Educational Status	Primary school	67 (47.5)
	Secondary school	46 (32.6)
	Higher education and above	13 (9.2)

Table 1: Socio demographic characteristics of type one DM in Mekele hospital, 2015.

Health profile of study subjects

Majority 104 (73.8%) of the study participant had no family history of diabetes. Most of 121 (85.8%) participants were not a member of diabetic association. Concerning duration of drug treatment, 49 (34.7%) of study participants were less than one year and 30 (21.2%) were between 3-5 years duration (Table 2).

Variables		Frequency (%)
Family history DM	Yes	35 (24.5)
	No	104 (73.5)
DM association membership	Yes	20 (14.2)
	No	121(85.8)
Duration of insulin therapy	<1yr	47 (32.9)
	1-3yr	33 (23.4)
	3-5yr	30 (21.2)
	>5yr	31 (22.3)

Table 2: Health profile of type one DM in Mekele Hospital, 2015.

Knowledge regarding insulin self administration

More than half 89 (63%) of the study subjects knew the definition of diabetes is high blood glucose and majority of the participants knew about insulin. Nearly two third 97 (68.8%) of study subject knew insulin vial is stored in refrigerator or cold place .greater than half 75 (53.1%) of respondents got the information about insulin from health workers and only 5 (3.1%) got from mass media.

The majority 117 (83%) of the study participant just had taken insulin soon after or just before taking food. Nearly three fourth (72.5%) of the study subjects knew the advantage of site rotation. 55 (39%) of respondent knew more than three sites for insulin injection.

Only 23 (16.3%) of respondent knew the angle of insulin injection is 45°. Majority 97 (68.9%) of participant didn't know the distance to rotate insulin injection on the same site is one thumb.

30 (21.2%) Study subjects understood the ways to reduce pain during insulin injection. 55 (39%) respondents answered the correct responses on the reason of massaging the site after injection (Table 3).

Only 15 (10.6%) study participants had knowledge on the complication of insulin therapy and nearly three fourth 108 (76.5%) knew the benefit of insulin self administration.

Item	Correct answer Frequency (%)	Wrong answer Frequency (%)
Know about diabetes mellitus	120 (85.1)	21 (14.91)
DM means high blood sugar	89 (63.1)	52 (36.87)
Know about insulin	118 (83.6)	23 (17.8)
Insulin vial is stored in the refrigerator or Cold place	98 (68.7)	43 (31.2)
Insulin injection is taken soon after Or just before taking food	117 (82.9)	24 (17)
The sites for insulin injection are abdomen, Thigh, Glutei and deltoid	56 (39)	85 (60.9)
The angle to administer insulin is 45°	23 (16.3)	118 (83.3)
The distance to rotate on the same site is One thumb	44 (31.2)	97 (68.7)
Ways to reduce pain during insulin injection are Inters the skin, do not manipulate the needle once Inserted, avoiding re using of the same site	30 (21)	110 (78.8)
The complications of insulin therapy are low Blood Sugar, insulin allergy, insulin resistance and wasting of subcutaneous tissue and wasting of subcutaneous tissue	15 (10.6)	126 (89.3)
The use of rotation of the injection site is to Reduce pain, prevent wasting of subcutaneous Tissues	98 (69.5)	43 (30.4)
Massage after injection is used to Reduce the rapid absorption of insulin	75 (76.5)	33 (23.4)
The benefit of insulin self administration Are, time saving, inexpensive and easy to take On self while traveling	108 (76.5)	33 (23.4)

Table 3: Data on knowledge regarding general information on DM and insulin self administration.

The mean knowledge score of study subjects was 7.07 with a maximum possible score of 13. Only 18 (13.07%) of participants had good knowledge, 78 (55%) participants had average knowledge and 45 (31.8%) had poor knowledge regarding diabetes and insulin self administration.

the cause for other health problems. Majority 66 (46.9%) study subjects thought insulin self administration correctly decrease glucose in the blood only 22 (15.9%) of respondents disagree with insulin self administration correctly decrease glucose in the blood (Table 4).

Attitude regarding insulin self administration

31 (22%) Study participants were agreeing with insulin causes other health problems and 86 (60.7%) study subjects believed insulin is not

Greater than half 91 (64.6%) of study subjects believed that insulin self administration is not tiresome and half 71 (50.3%) of respondent thought insulin self administration brought stigma.

Item	Agree Frequency (%)	Not Sure Frequency (%)	Disagree Frequency (%)
Insulin cause other health problem	31 (22)	25 (17.7)	85 (60.2)
Insulin self-administration correctly Decrease glucose in the blood	66 (46.8)	52 (36.8)	23 (15.6)
Insulin self administration is a tiresome	44 (32.1)	14 (9.9)	91 (64.5)
Insulin self administration bring stigma	71 (50.3)	16 (11.3)	54 (38.2)
Insulin self administration is beneficiary	99 (70.2)	30 (21.2)	12 (8.5)

Table 4: Data on attitude of type 1 diabetic patients regarding on insulin self administration.

Out of 141 type 1 diabetic patients included in this study, 95 (67.4%) had favorable attitude and 46 (32.5%) had un favorable attitude .the

mean attitude score of the participants was 3.07 with maximum score of 15. This result showed that most type 1 participant had favorable attitude on insulin self administration.

Discussion

The study was carried out to assess the knowledge and attitude of type 1 diabetic patients on insulin self administration in Mekele hospital. The study showed that the mean knowledge score of study subjects were 7.27. Study participants had average knowledge regarding insulin self administration. This study had demonstrated average level of knowledge regarding insulin self administration. The prevalence of knowledge was lower compared to studies done in India and Bangalore which were 68% and 86.7% [6,8]. This difference from the previous studies might be due to high illiteracy rate of study participants, lack of self insulin injection education facilities and low access of media and NGOs in awareness creation about insulin self administration.

Of the study participants thought that diabetes mellitus mean high blood sugar. This was better than a similar study done in Bahir Dar, Fellegehiwot hospital which was 33.4% [9]. This difference may be related with patient health profile difference.

Majority of type 1 diabetic patients (82%) awarded that the action of insulin is to lower glucose in the blood. This study was supported by similar study conducted in Bangalore (80%) [8].

Another important finding of this study was a limited knowledge about the complication of insulin therapy (10%), which was much lower compared to studies done in Bangalore (86.67%) and Nigeria (66.4%) [8,10] this difference might be due to high illiteracy rate, environmental factor and socio economic status of the study participant of the present study. This level of inadequate knowledge regarding complication of insulin therapy may lead to decrease awareness of patients for complication and these are high economic burden of the country in the management of complications which comes due to inadequate awareness of the complications and lack of adequate organized health education about diabetes.

Participant aware the sites of insulin injection are abdomen, thigh, glutei and deltoid. This finding was different from a study conducted in Bangalore, which was 53.3% [8]. This could be lack of organized health education and negligence of health care workers on insulin self administration. Inadequate knowledge regarding site of insulin injection may lead to develop complication of insulin therapy. These findings high light the need of emphasizing insulin injection during diabetes education.

Among respondent, 68.6% knew an open vial could be preserved in refrigerator or cold place. This study had slight different from study conducted in Indian and Bangalore 72.4% and 78.33% respectively [6,8]. This difference from the previous studies could be low economics status of study subjects and health patient negligence to educate about insulin vial preservation.

The mean attitude score of the study subjects was, participants had favorable attitude and study participants had favorable attitude and study participants had unfavorable attitude. It was lower compared to study done in Bangalore, which was 81.7% had favorable attitude. 18.3% had most favorable attitude and none of them had unfavorable attitude [8]. The difference from Bangalore finding might be lack of trained man power to create adequate awareness toward insulin therapy. The present finding was better than another study conducted

in India which was only 32% had favorable attitude and others had unfavorable attitude towards insulin self administration [6]. This difference might be due to environmental and socio-cultural factors of participants.

Limitation of the study

This study was also limited to diabetic patients who were on subcutaneous insulin therapy and only site was included in Mekele hospital (chronic follow up) it assessed only their knowledge and attitude practice was not included. We did not observe what actually do while they insert insulin.

Conclusion

The finding of this study showed that type 1 diabetes patients had average knowledge and less favorable attitude among type 1 diabetes toward insulin self administration but there is still there is a gap on knowledge and attitude of type 1 diabetes patients on insulin therapy.

Recommendation

To Medias; most of the patients get the information from health workers, there is still knowledge gap so that it is better to deliver updated information from the media.

To health science professionals: at the time of follow up strict education has to give for each patient in order to increase their knowledge, attitude and practice.

References

1. Majaliwa ES, Elusiyani BE, Adesiyun OO, Laigong P, Adeniran AK, et al. (2008) Type 1 diabetes mellitus in the African population: epidemiology and management challenges. *Acta Biomed* 79: 255-259.
2. Derr RL, Sivanandy MS, Bronchi-hall L, Rodriguez A (2007) Insulin related knowledge among health care professional in internal medicine. *Diabetes spectr* 20: 177-185.
3. The Joint commission on accreditation of health care organization (JCAHO) JCAHO suggests ways to prevent medical errors. *Health risk man* 22: 19-20.
4. Garg MK (2002) Current perspective in insulin therapy in the management of diabetes mellitus. *J Indian Med Assoc* 100: 194-195, 202.
5. Majra JP, Acharya D (2009) Awareness regarding self care among diabetics in Rural India. *Middle East journal of family medicine* 7: 25-26.
6. Surendranath A, Nagaraju B, Padmavathi GV, Anand SC, Fayaz P, et al. (2012) A Study to assess the knowledge and attitude of insulin self administration among patients with diabetes mellitus in India. *Asian journal of pharmaceutical and clinical research* 5: 1.
7. [No authors listed] (1994) Effect of intensive diabetes treatment on the development and progression of long-term complications in adolescents with insulin-dependent diabetes mellitus: Diabetes Control and Complications Trial. *Diabetes Control and Complications Trial Research Group. J Pediatr* 125: 177-188.
8. Namita S (2005) A descriptive study to assess the knowledge and attitude regarding self administration of insulin injection among DM patients in Kempa chaluwamba general hospital at Malleswaram, Bangalore.
9. Sr. Tresa P (2005) Effectiveness of individual planned teaching on self insulin Administration for patients with diabetes mellitus in A selected hospital at mangalore. Rajiv Gandhi University of health sciences, Karnataka.

10. Jasper US, Opara MC, Pyiki EB, Akinrolie O (2014) Knowledge of insulin use and its determination among Nigerian insulin requiring diabetes patients. Journal of diabetes and metabolic disorder 13: 10.