

Unraveling the Progression of Ischemic Core Genome-Wide by Bioinformatics Analysis of Permanent Middle Cerebral Artery Occlusion (PMCAO) Mouse Model Brain Regions Genes

Motohide Hori^{1,2}, Tomoya Nakamachi^{2,3}, Junko Shibato^{2,4}, Randeep Rakwal^{2,4,5*}, Seiji Shioda^{2,4} and Satoshi Numazawa¹

¹Division of Toxicology, Department of Pharmacology, Toxicology and Therapeutics, School of Pharmacy, Showa University, 1-5-8 Hatanodai, Shinagawa, Tokyo, Japan
²Department of Anatomy I, School of Medicine, Showa University, 1-5-8 Hatanodai, Shinagawa, Tokyo, Japan
³Laboratory of Regulatory Biology, Graduate School of Science and Engineering, University of Toyama, Toyama, Toyama, Japan
⁴Global Research Center for Innovative Life Science, Peptide Drug Innovation, School of Pharmacy and Pharmaceutical Sciences, Hoshi University, 4-41 Ebara 2-chome, Shinagawa, Tokyo, Japan
⁵Faculty of Health and Sport Sciences and Tsukuba International Academy for Sport Studies (TIAS), University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki, Japan

Keywords: Brain ischemia; Ischemic core and penumbra; DNA microarray; IPA analysis, PACAP38

Introduction

The neuropeptide, pituitary adenylate-cyclase activating polypeptide (PACAP) [1-3], is the major focus of research as a neuroprotective factor in the group of Prof. S. Shioda [4-16]. A major emphasis of the research is on the neuroprotective effects of PACAP38 on the brain, particularly in brain ischemia [17]. Therein, a permanent middle cerebral artery occlusion (hereafter referred to as PMCAO) mouse model has been established and used for unraveling of the genome-wide gene expression profiles by high-throughput omics approach, namely DNA microarray technology. DNA microarray analysis of the whole brain/ischemic hemisphere and specific brain regions of the ischemic core hereafter, the IC and penumbra hereafter, the P with or without PACAP38 treatment have been performed [11-16]. It is to be noted that the reason for utilizing intraluminal filament technique-based PMCAO model over the transient MCAO is to avoid reperfusion injury, in our research model (PMCAO). However, it should be emphasized that different research groups are using different stroke models such as the MCAO, resulting in greater insight into how PACAP treatment influences the brain ischemia. Overall, these genomic data on differential gene expression in the brain of mouse PMCAO model resulted in us i) obtaining the transcriptome profiles of ischemic brain hemispheres along with the diverse categories of gene families being modulated under the ischemic condition [11-13], ii) unraveling specific gene expressions and localization of molecular factors in the IC and P effected specifically by the PACAP38 treatment [14,15], and iii) providing the explanation and validation of the dye-swap, two-color DNA microarray approach [16]. The vast inventories of differential gene expressions generated from these numerous analyses have not only revealed the importance of both whole hemisphere and region-specific analyses in genome-wide identification of target molecular factors that might play a role in the neuroprotective function of PACAP38, but also provided a valuable resource for further study by the scientific community.

Short Communication

Our most recent data analyzing the specificity of the PACAP38 treatment in delineating the molecular expressions in the IC and the P at 6 h post-treatment (early) and 24 h (late) progression of the ischemia revealed that PACAP38 indeed has a positive influence in helping recover the ischemic insult [15]. However, we did not clarify the progression of ischemia itself, i.e. in the absence of any treatment (PACAP38). As also kindly suggested by an anonymous reviewer in our previous research paper [15] it would be important to investigate the progression of ischemia in order to effectively understand the neuroprotective effects of PACAP38, the overall goal of the research. Therefore, in this communication, we primarily present the DNA

microarray analysis data of the IC and P regions at 6h in the ischemic brain compared to the healthy IC and P regions, as outlined in the experimental design (Figure 1). In other words, our research aimed to unravel differential gene expression profiles in the IC and P regions without PACAP38 treatment in order to know which genes are specifically functioning during progression of the ischemic core. To do so, we carried out a whole genome DNA microarray analysis (Agilent mouse whole genome 4x44K DNA chip; G4131F) of the healthy versus ischemic IC and P regions. Briefly, three mice each in PMCAO groups for IC and P regions in the ischemic brain over corresponding controls were used that exhibited neurological grades G1 and G2, for the subsequent downstream analysis. Ischemic core and P regions and corresponding healthy core and P regions were carefully removed with a sterile scalpel, and placed in sterile 2 ml Eppendorf tubes. Samples were




Figure 1: Strategy for analyses of the specific brain regions, ischemic core (IC) and the penumbra (P), in the permanent middle cerebral artery occlusion (PMCAO) mouse model and DNA microarray approach for identifying the molecular factors potentially involved in PACAP38-mediated neuroprotection. Two analyses approaches are presented here for progression of infarct, first, gene profiling of the 6 h IC and P regions (minus PACAP38 treatment), and second, use of the Ingenuity Pathway Analysis (IPA) tool for identification of the annotated genes and networks therein.

*Corresponding author: Randeep Rakwal, Faculty of Health and Sport Sciences and Tsukuba International Academy for Sport Studies (TIAS), University of Tsukuba, 1-1-1 Tennodai, Tsukuba 305-8574, Ibaraki, Japan, Tel: +81-029-853-2681; E-mail: plantproteomics@gmail.com

Received September 09, 2015; Accepted October 16, 2015; Published October 19, 2015

Citation: Hori M, Nakamachi T, Shibato J, Rakwal R, Shioda S, et al. (2015) Unraveling the Progression of Ischemic Core Genome-Wide by Bioinformatics Analysis of Permanent Middle Cerebral Artery Occlusion (PMCAO) Mouse Model Brain Regions Genes. *Transcriptomics* 3: 112. doi:10.4172/2329-8936.1000112

Copyright: © 2015 Hori M, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

then quickly immersed in liquid nitrogen and stored in -80°C prior to further analysis. Animal care and experimental procedures were used as approved by the Institutional Animal Care and Use Committee of Showa University (School of Medicine), Tokyo, Japan, and the PMCAO model mice, (PACAP38 treatment), dissection of brain, sampling and storage, and total RNA extraction followed by DNA microarray analysis was performed using our established dye-swap approach as described previously [11-16]. The outputs of DNA microarray analysis are freely available to the public under the series number GSE 67421 [18] at the NCBI GEO public functional genomics data repository [19].

Additional bioinformatics analysis was carried out using the

Ingenuity Pathway Analysis (IPA; Ingenuity® Systems, www.ingenuity.com; Content version: 23814503, Release Date: 2015-03-22, Qiagen) bioinformatics tool that provided us with the latest annotations of genes. The biological function and network analysis were also generated through the use of IPA. The data set from microarray (6 h, IC and P), which is the differentially expressed ($\geq/\leq 1.5/0.75$ -fold compared to saline control) genes, and their corresponding fold change values were uploaded as an Excel spread sheet into the IPA tool. To create gene networks, genes were overlaid onto a global molecular network developed from information contained in the ingenuity knowledge base. The functional analysis identified the biological functions that were most significant to the data set (p -value < 0.05) according to Right-tailed Fisher's exact test. Further, we also used the PACAP38 treatment data (GSE 62884) for the IC and P regions to generate a parallel list of annotated genes. In this PMCAO-PACAP38 model, an intracerebroventricularly PACAP38 (1 pmol) injection over a control saline (0.9% sodium chloride, NaCl) treatment was used.

Results presented in Figure 2A show the number of differentially expressed (up- and down-regulated) annotated genes in the IC (456) and P (667) compared to the whole brain (hemisphere, 881). Similarly, in Figure 2B, the number of differentially expressed annotated genes after PACAP38 treatment for the IC (582) and P (501) compared to the whole brain (hemisphere, 800) is presented. These genes are presented as a list of color-coded up-/down-regulated genes along with their Entrez gene names, Agilent probe number and fold-change, for clarity in Supplementary Tables 1 and 2 (Figure 2A) 3 and 4 (Figure




Figure 2: Venn diagram showing the numbers of differentially expressed annotated genes in the PMCAO ischemic hemisphere (881), as compared to the IC (456), and P (667) regions (A), and also in the PACAP38 influenced PMCAO ischemic hemisphere (800), as compared to the IC (582), and P (501) regions (B) at 6 h. Genes were annotated using the IPA bioinformatics tool. IC, ischemic core; P, penumbra.




Figure 3: The top 3 networks for the IC (upper panel) and P (lower panel) regions by the IPA bioinformatics tool.

Ischemic Core (IC)		Penumbra (P)	
Fold Change up-regulated		Fold Change up-regulated	
Molecules	Exp. Value (+)	Molecules	Exp. Value (+)
<i>Ngp</i>	6.341	<i>CAMK2D</i>	8.03
<i>CCR2</i>	4.449	<i>SHOX2</i>	7.571
<i>ANXA10</i>	4.357	<i>COL10A1</i>	5.096
<i>TMEM190</i>	3.973	<i>SAMD3</i>	4.693
<i>AVPR1A</i>	3.858	<i>CCKAR</i>	4.29
<i>FAM90A1</i>	3.616	<i>OPN4</i>	3.736
<i>Cxcl3</i>	3.583	<i>KRT23</i>	3.522
<i>LTF</i>	3.217	<i>TNFRSF8</i>	3.35
<i>TREM3</i>	3.202	<i>SNCA</i>	3.132
<i>NMU</i>	3.157	<i>SIX3</i>	3.041
Fold Change down-regulated		Fold Change down-regulated	
Molecules	Exp. Value (-)	Molecules	Exp. Value (-)
<i>TBX22</i>	3.043	<i>TUB</i>	4.147
<i>ADAMTS12</i>	2.782	<i>TMEM199</i>	3.648
<i>N4BP2</i>	2.596	<i>SLC13A1</i>	3.116
<i>FGF16</i>	2.565	<i>CYP11A1</i>	2.78
<i>RAB38</i>	2.552	<i>ISL2</i>	2.647
<i>MB</i>	2.425	<i>SOX17</i>	2.613
<i>ART4</i>	2.368	<i>LY6D</i>	2.598
<i>SAMD11</i>	2.359	<i>OR2G6</i>	2.59
<i>ITIH5</i>	2.238	<i>S100a7a</i>	2.584
<i>RAPGEF6</i>	2.306	<i>PTCH2</i>	2.581

Table 1. The top molecules identified in the IPA analysis for the ischemic core (IC) progression. The penumbra (P) top molecules are also presented for comparison.

2B) (see Supplementary Information Tables). At a glance these data reveal differences in the type of genes expressed in the IC and P regions during progression of infarct, including that after PACAP38 treatment. Subsequently, these gene lists were used to generate the networks for the molecules/pathways being influenced in the IC and P regions (for without PACAP38 treatment), examples of which are shown in Figure 3. The top up- and down-regulated molecules or the IC and P regions are also presented in Table 1 and these genes might be linked to the progression of the ischemic core.

Here we have newly communicated the importance of using a specific analysis of the gene expression in IC and P brain regions compared to healthy control regions to know the gene profile status in the progression of IC using IPA bioinformatics tool providing the latest annotated gene lists and generated gene networks. This study will form the basis for a detailed bioinformatics analysis functionally identifying the genes (Figure 2A, Figure 3, and Table 1) involved specifically in the progression of ischemia, which is not well understood as yet. Furthermore, by revisiting the PACAP38 influenced annotated gene lists in light of the progression of ischemia we may be able to further understand the diversity and specific function of molecular factors, pathways and networks that may underlie the potential mechanism behind PACAP38 neuroprotective action in the ischemic brain. For this, functional genomics studies will be the only way forward to understand how and why PACAP38 is neuroprotective in the brain for any meaningful development of a stroke therapy using PACAP38 or its analogs.

Acknowledgments

MH gratefully acknowledges the members of Professor Seiji Shioda's laboratory (at the Department of Anatomy I) for their support and encouragement during this study. This work was supported by Grants-in Aid for Scientific Research (KAKENHI: 23249079, 24592681, 24592680), and by the MEXT-Support Program for the Strategic Research Foundation at Showa University (2012-16) to SS. RR acknowledges the support of Grants-in Aid for Scientific Research, KAKENHI (C): 26462745 for conducting this research.

References

- Kimura C, Ohkubo S, Ogi K, Hosoya M, Itoh Y, et al. (1990) A novel peptide which stimulates adenylate cyclase: molecular cloning and characterization of the ovine and human cDNAs. *Biochem Biophys Res Commun* 166: 81-89.
- Arimura A (1998) Perspectives on pituitary adenylate cyclase activating polypeptide (PACAP) in the neuroendocrine, endocrine, and nervous systems. *Jpn J Physiol* 48: 301-331.
- Shioda S, Ozawa H, Dohi K, Mizushima H, Matsumoto K, et al. (1998) PACAP protects hippocampal neurons against apoptosis: involvement of JNK/SAPK signaling pathway. *Ann N Y Acad Sci* 865: 111-117.
- Ohtaki H, Funahashi H, Dohi K, Oguro T, Horai R, et al. (2003) Suppression of oxidative neuronal damage after transient middle cerebral artery occlusion in mice lacking interleukin-1. *Neurosci Res* 45: 313-324.
- Ohtaki H, Nakamachi T, Dohi K, Aizawa Y, Takaki A, et al. (2006) Pituitary adenylate cyclase-activating polypeptide (PACAP) decreases ischemic neuronal cell death in association with IL-6. *Proc Natl Acad Sci U S A* 103: 7488-7493.
- Ohtaki H, Nakamachi T, Dohi K, Shioda S (2008) Role of PACAP in ischemic neural death. *J Mol Neurosci* 36: 16-25.
- Ohtaki H, Takeda T, Dohi K, Yofu S, Nakamachi T, et al. (2007) Increased mitochondrial DNA oxidative damage after transient middle cerebral artery occlusion in mice. *Neurosci Res* 58: 349-355.
- Ohtaki H, Ylostalo JH, Foraker JE, Robinson AP, Reger RL, et al. (2008) Stem/progenitor cells from bone marrow decrease neuronal death in global ischemia by modulation of inflammatory/immune responses. *Proc Natl Acad Sci U S A* 105: 14638-14643.
- Nakamachi T, Endo S, Ohtaki H, Yin L, Kenji D, et al. (2005) Orexin-1 receptor expression after global ischemia in mice. *Regul Pept* 126: 49-54.
- Nakamachi T, Ohtaki H, Yofu S, Dohi K, Watanabe J, et al. (2010) Endogenous pituitary adenylate cyclase activating polypeptide is involved in suppression of edema in the ischemic brain. *Acta Neurochir Suppl* 106: 43-46.
- Hori M, Nakamachi T, Rakwal R, Shibato J, Nakamura K, et al. (2012) Unraveling the ischemic brain transcriptome in a permanent middle cerebral artery occlusion model by DNA microarray analysis. *Disease Models & Mechanisms* 5: 270-283.

12. Hori M, Nakamachi T, Rakwal R, Shibato J, Ogawa T, et al. (2012) Transcriptomics and proteomics analyses of the PACAP38 influenced ischemic brain in permanent middle cerebral artery occlusion model mice. *J Neuroinflammation* 9: 256.
13. Hori M, Nakamachi T, Rakwal R, Shibato J, Ogawa T, et al. (2013) Correction: Transcriptomics and proteomics analyses of the PACAP38 influenced ischemic brain in permanent middle cerebral artery occlusion model mice. *J Neuroinflammation* 10: 18.
14. Hori M, Nakamachi T, Shibato J, Rakwal R, Tsuchida M, et al. (2014) PACAP38 differentially effects genes and CRMP2 protein expression in ischemic core and penumbra regions of permanent middle cerebral artery occlusion model mice brain. *Int J Mol Sci* 15: 17014-17034.
15. Hori M, Nakamachi T, Shibato J, Rakwal R, Shioda S, Numazawa S (2015) Unraveling the specific ischemic core and penumbra transcriptome in the permanent middle cerebral artery occlusion mouse model brain treated with the neuropeptide PACAP38. *Microarrays* 4: 2-24.
16. Hori M, Shibato J, Nakamachi T, Rakwal R, Ogawa T, et al. (2015) Two-color Dye-swap DNA Microarray approach toward confident gene expression profiling in PMCAO mouse model for ischemia-related and PACAP38-influenced genes. *Genom Data* 3: 148-154.
17. Gusev EI, Skvortsova VI (2003) *Brain Ischemia*. Kluwer Academic/Plenum Publishers: New York.
18. <https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE67421>.
19. <http://www.ncbi.nlm.nih.gov/geo/info/linking.html>.

Citation: Hori M, Nakamachi T, Shibato J, Rakwal R, Shioda S, et al. (2015) Unraveling the Progression of Ischemic Core Genome-Wide by Bioinformatics Analysis of Permanent Middle Cerebral Artery Occlusion (PMCAO) Mouse Model Brain Regions Genes. *Transcriptomics* 3: 112. doi:[10.4172/2329-8936.1000112](https://doi.org/10.4172/2329-8936.1000112)

OMICS International: Open Access Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission/>