

## Utility of Upper Gastrointestinal Endoscopy for Management of Patients with Roundup® Poisoning

Kenji Iwai<sup>1</sup>, Masato Miyauchi<sup>2</sup>, Daisuke Komazawa<sup>1</sup>, Ryoko Murao<sup>1</sup>, Hiroyuki Yokota<sup>2</sup>, and Atushi Koyama<sup>1</sup>

<sup>1</sup>Department of Emergency and Critical Care Medicine, Iwaki Kyoritu General Hospital, Fukushima, Japan

<sup>2</sup>Department of Emergency and Critical Care Medicine, Nippon Medical School, Tokyo, Japan

\*Corresponding author: Masato Miyauchi, Department of Emergency and Critical Care Medicine, Nippon Medical School, Tokyo, Japan, Tel: +81-3-3822-2131; E-mail: [m-masa@nms.ac.jp](mailto:m-masa@nms.ac.jp)

Received date: Nov 03, 2014, Accepted date: Dec 05, 2014, Published date: Dec 08, 2014

Copyright: © 2014, Miyauchi M, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### Abstract

**Introduction:** Roundup® is a herbicide widely used in Japan in gardening and agriculture. When ingested, Roundup is highly toxic, but gastrointestinal decontamination, including gastric lavage, is not routinely performed after ingestion. Endoscopy may be useful in managing individuals with liquid herbicide poisoning, by identifying gastric residual contents, assessing mucosal damage and retrieving herbicide directly by aspiration.

**Case report:** A 73 year old, 40 kg female with a history of depression was transported to our emergency room by ambulance 1 h after attempting suicide by ingesting 100 ml Roundup, which contains 48% glyphosate-potassium, and 52% surfactant and water. This volume was below a fatal dose (<5000 mg/kg), but may have caused organ dysfunction and mucosal damage. After confirming respiratory and circulatory stability and after obtaining informed consent from the patient, endoscopy (XQ 260; Olympus, Tokyo, Japan) was performed in the emergency room to retrieve residual herbicide. About 80 ml of herbicide in the stomach were aspirated endoscopically with only mild erosion observed in the mucosa of the stomach. The patient was able to resume oral intake 2 days after endoscopy and was discharged without any complications on day 5.

**Conclusion:** Endoscopy may be useful in cases of liquid poisoning including, Roundup, both to determine the amount of residual toxin and to remove it from the stomach.

**Keywords:** Endoscopy, Liquid poisoning, Gastric lavage, Round up®


### Introduction

Roundup® is an herbicide widely used in Japan for gardening and agriculture. Many preparations of this herbicide are available, containing various amounts of glyphosate (N-[phosphonomethyl] glycine), surfactant, and water. The toxicity of herbicide derives from the glyphosate and surfactant [1,2], with mortality rates following its intentional ingestion reported to range from 3.2% to 29% [3,4]. Patients poisoned with these preparations generally receive symptomatic and supportive management [5]. Although gastrointestinal decontamination, including gastric lavage, is not performed routinely [6,7], the retrieval of residual toxic contents from the stomach may benefit patients. Direct observation of gastric residual contents by upper gastrointestinal endoscopy may reveal the need for gastric decontamination [8]. Endoscopy in patients poisoned with liquid herbicide may be useful for assessing gastric residual contents and mucosal damage, as well as for direct retrieval of the toxic agent by aspiration.

### Case Report

A 73 year old, 40 kg female with a history of depression was transported to our emergency room by ambulance 1 h after attempting suicide by ingestion of 100 ml of the herbicide Roundup®, containing 48% glyphosate-potassium, and 52% surfactant and water. At

admission, her blood pressure was 177/85 mmHg, her heart rate was 60 beats/min, her respiratory rate was 20 breaths/min and her temperature was 36.4°C. Initial blood gas analysis showed a pH of 7.398, arterial carbon dioxide tension (PaCO<sub>2</sub>) of 44.9 mmHg, arterial oxygen tension (PaO<sub>2</sub>) of 72.1 mmHg, actual bicarbonate level of 30.1 mmol/l (normal range, 22 to 32 mmol/l), and base excess of +4.8 mmol/l (-2 to 2 mmol/l). Laboratory data were normal. She had a Glasgow Coma Scale score of 15 (E4V5M6). The patient had ingested 1200 mg/kg of glyphosate, less than a fatal dose (<5000 mg/kg); however, the surfactant used in this product was not published. The herbicide may have caused organ dysfunction and mucosal damage. Endoscopy was proposed to retrieve residual herbicide by aspiration. The purpose and contraindications for endoscopy were fully explained to the patient, who provided informed consent. After confirming respiratory and circulatory stabilization, endoscopy (XQ 260; Olympus, Tokyo, Japan) was performed in the emergency room. Sedation was not used. The patient was placed in the left lateral decubitus position, and the endoscope was slowly advanced. As much residue in the stomach was aspirated as possible (about 80 ml), with only mild erosion observed in the mucosa of the stomach (Figure 1). After removing the endoscope, a laxative and activated carbon were administered to the patient, and she was admitted to the intensive care unit. Because peritonitis and abdominal tenderness were absent, the patient was able to resume oral intake on day 2. She was discharged without any complications on day 5.



**Figure 1:** Endoscopic images in our patient. A: residual toxic substance. B and C: erosion of gastric mucosa by toxic substances and aspiration by endoscopy.

## Discussion

Roundup®, a trade name for a herbicide containing various formulations of glyphosate salts, is widely used worldwide for agricultural purposes. The herbicide ingested by our patient contained 48% potassium glyphosate salts. Glyphosate has an oral LD50 >5000 mg/kg in animals. Our patient had ingested 1200 mg/kg of glyphosate. Although below a fatal dose, the volume [9] and concentrations [10] put her at risk factor for severe toxic effects, including gastric erosion, renal dysfunction, hepatic impairment and shock [11]. Furthermore the surfactant used in this product could not be determined.

Although gastric lavage for gastrointestinal decontamination is not performed routinely in these patients [12], the need to retrieve any residual toxic substances in the stomach is unclear. A three-step method of evaluating the need for gastrointestinal decontamination has been proposed [13], with treatment decisions based on assessing the risk of poisoning and the immediate and long-term benefits and risks of the decontamination procedure. Endoscopy was previously used in the diagnosis and treatment by gastrointestinal decontamination of a patient with gastric bezoars [14]. Because the herbicide ingested by our patient is a liquid, it could be retrieved by aspiration using endoscopy. Endoscopy thus confirmed the gastric contents, with endoscopic aspiration facilitating gastric lavage and preventing further passage of toxic substances into the small bowel. Furthermore endoscopy could estimate the erosion of the stomach, not only improving patient outcomes but avoiding hospital admission. Endoscopy in poisoning patients should only be used after airway protection. Although there is little clinical evidence on the effectiveness of gastric lavage for managing patients with liquid poisoning, we found that endoscopy confirmed the presence of residual substances, and also facilitated the clearance of residue from the stomach.

## Conclusion

Endoscopy may be useful in patients with liquid poisoning, including Roundup, both for determining the amount of residual substance and for removing it from the stomach.

## References

1. Adam A, Marzuki A, Abdul Rahman H, Abdul Aziz M (1997) The oral and intratracheal toxicities of ROUNDUP and its components to rats. *Vet Hum Toxicol* 39: 147-151.
2. Martinez TT, Long WC, Hiller R (1990) Comparison of the toxicology of the herbicide roundup by oral and pulmonary routes of exposure. *Proc West Pharmacol Soc* 33: 193-197.
3. Roberts DM, Buckley NA, Mohamed F, Eddleston M, Goldstein DA, et al. (2010) A prospective observational study of the clinical toxicology of glyphosate-containing herbicides in adults with acute self-poisoning. *Clin Toxicol (Phila)* 48: 129-136.
4. Lee CH, Shih CP, Hsu KH, Hung DZ, Lin CC (2008) The early prognostic factors of glyphosate-surfactant intoxication. *Am J Emerg Med* 26: 275-281.
5. Talbot AR, Shiaw MH, Huang JS, Yang SF, Goo TS, et al. (1991) Acute poisoning with a glyphosate-surfactant herbicide ('Roundup'): a review of 93 cases. *Hum Exp Toxicol* 10: 1-8.
6. Vale JA (1997) Position statement: gastric lavage. *American Academy of Clinical Toxicology; European Association of Poisons Centres and Clinical Toxicologists. J Toxicol Clin Toxicol* 35:711-719.
7. Vale JA, Kulig K; American Academy of Clinical Toxicology; European Association of Poisons Centres and Clinical Toxicologists (2004) Position paper: gastric lavage. *J Toxicol Clin Toxicol* 42: 933-943.
8. Miyauchi M, Hayashida M, Hirata K, Hirata K, Yokota H (2013) Gastric lavage guided by ultrathin transnasal esophagogastroduodenoscopy in a life-threatening case of tobacco extract poisoning: a case report. *J Nippon Med Sch* 80: 307-311.
9. Chen YJ, Wu ML, Deng JF, Yang CC (2009) The epidemiology of glyphosate-surfactant herbicide poisoning in Taiwan, 1986-2007: a poison center study. *Clin Toxicol (Phila)* 47: 670-677.
10. Moon JM, Chun BJ (2010) Predicting acute complicated glyphosate intoxication in the emergency department. See comment in *PubMed Commons below Clin Toxicol (Phila)* 48: 718-724.
11. Bradberry SM, Proudfoot AT, Vale JA (2004) Glyphosate poisoning. *Toxicol Rev* 23: 159-167.
12. Benson BE, Hoppu K, Troutman WG, Bedry R, Erdman A, et al. (2013) Position paper update: gastric lavage for gastrointestinal decontamination. *Clin Toxicol (Phila)* 51: 140-146.
13. Bailey B (2008) To decontaminate or not to decontaminate? The balance between potential risks and foreseeable benefits. *Clin Ped Emerg Med* 9: 17-23.
14. Wells CD, Luckritz TC, Rady MY, Zornik JM, Leighton JA, et al. (2006) Bezoar formation requiring endoscopic removal after intentional overdose of extended-release nifedipine. *Pharmacotherapy* 26:1802-1805.