

Utilizing a Dietary Supplement to Modify Small Intestinal Secretory Function

Layla Al-Nakkash*

Midwestern University, Glendale, AZ, USA

*Corresponding author: Layla Al-Nakkash, Midwestern University, Glendale, AZ, USA, Tel: 623-572-3719; Fax: 623-572-3673; E-mail: lalnak@midwestern.edu


Received date: November 24, 2016; Accepted date: November 28, 2016; Published date: November 30, 2016

Copyright: © 2016 Al-Nakkash L. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Editorial

During the last decade the hypothesis that consuming genistein may modify intestinal secretory function has been investigated. This is of importance in clinically relevant illnesses whereby intestinal abnormalities result in modifications of secretory function, for example in Cystic Fibrosis (CF) and diabetes. Fortunately, both of these diseases have murine models to better test the effects of dietary manipulations.

Genistein, a naturally occurring isoflavononic phytoestrogen found in soy [1], is readily absorbed across the intestines and reaches micro molar concentrations in the serum of individuals consuming a soy milk diet [2]. Within the small intestine, chloride secretion occurs at the crypts, a process that requires several transporters working in concert (Figure 1). The $\text{Na}^+/\text{K}^+/\text{2Cl}^-$ (NKCC1) co-transporter provides the entry point for Cl^- into the epithelial crypt cells, a driving force for Cl^- exit across the apical membrane is maintained by recycling of K^+ across the basolateral membrane, the Na^+/K^+ -ATPase maintains Na^+ and K^+ concentration gradients across the membrane, and the major route for Cl^- exit (secretion) across the apical membrane into the lumen is via the CF trans membrane conductance regulatory protein, CFTR, Cl^- channel [3,4]. Loss of any individual transporter (or the regulation of) will result in intestinal secretory dysfunction.



Genistein is a known modifier of several of those key proteins/transporters involved in successful intestinal secretory transport; NKCC1 [5], K^+ channels [6] and CFTR [7]. Indeed, initial studies revealed several beneficial effects of acute application of genistein to freshly isolated intestinal tissue. Acute application of genistein (75 μm)

to jejunal tissue from male Swiss Webster mice has been shown to elicit increased trans epithelial ion secretion (a measure of chloride secretion), via inhibition of a PDE3-mediated pathway [8,9], whereas acute application of genistein (50 μm) to duodenal tissue from C57BL6J mice has been shown to stimulate HCO_3^- secretion via an estrogen receptor/PI3K pathway [10]. Those studies, and others, led the way to the development of dietary studies. Consumption of a genistein-containing diet (600 mg genistein/kg diet) for 4-weeks has been shown to increase trans epithelial chloride secretion across jejunum in lean (C57BL6J) female mice [11], while concomitantly increasing serum genistein levels to low micro molar range (comparable to humans consuming soy milk). In broiler chickens, improvements in intestinal morphometry have been measured (villus length, villus width, crypt depth, villus length/crypt depth ratio) following dietary genistein (5 mg genistein/kg feed) [12], indicating such food additives can promote gut health and function.

Murine models exhibiting intestinal secretory dysfunction such as those with CF and diabetic obesity provide useful settings with which to test genistein's effects. Mice homozygous for the DF508 mutation (the most common clinical mutation) are characterized with severe intestinal disease requiring continual treatment with laxative for survival [13]. This murine model mimics the intestinal obstruction (meconium ileus) seen in some CF patients [14]. Examination of the effects of dietary supplementation with genistein (600 mg genistein/kg diet) in the absence of laxative, in DF508 CF mice between 21-65 days of age, resulted in a significant increase in survival rate from 68% to 84% [15], suggesting potential therapeutic benefits. Favorable effects of genistein consumption on jejunal function in the diabetic obese ob/ob mouse have recently been evaluated. This diabetic obese mouse model exhibits a slowed gastrointestinal transit, thus mimicking one of the clinical symptoms noted in diabetics [16,17]. After consuming genistein (600 mg genistein/kg diet) for a period of 4-weeks, ob/ob mice had rescued basal Trans epithelial chloride secretion (i.e., levels returned to those measured in lean controls). Interestingly, genistein's beneficial effect on jejunum secretory function was mediated by sex-dependent mechanisms: Increasing KCa sensitive transepithelial secretion in females, and increasing both Na^+/K^+ -ATPase activity along with NKCC1 expression in males [18], suggesting potential novel pharmacotherapeutic objectives in the treatment of intestinal dysfunction in diabetes.

The aforementioned studies aimed at improving intestinal secretory transport are relevant given that loss of secretory function can lead to increased bacterial translocation. There is no doubt that the mechanism(s) of action of dietary genistein are complex and its targets are countless and complicated, with sex-differences making matters more interesting. Thus, it is incumbent upon the scientific community to be especially rigorous and thorough in the assessment of the usefulness of such dietary manipulations. That said, understanding the

benefits of dietary supplementation with genistein in experimentally clinically relevant models of intestinal secretory dysfunction will certainly pave the way towards potential future applicability in clinical settings.

References

1. Murphy PA (1982) Phytoestrogen content of processed soybean products. *Food Technol* 36: 60-64.
2. Xu X, Wang HJ, Murphy PA, Cook L, Hendrich S (1994) Daidzein is a more bioavailable soy milk isoflavone than is genistein in adult women. *J Nutr* 124: 825-832.
3. Anderson MP, Welsh MJ (1991) Calcium and cAMP activate different chloride channels in the apical membrane of normal and cystic fibrosis epithelia. *Proc Natl Acad Sci* 88: 6003-6007.
4. Clarke LL, Grubb BR, Yankaskas JR, Cotton CU, McKenzie A, et al. (1994) Relationship of a non-cystic fibrosis transmembrane conductance regulator-mediated chloride conductance to organ-level disease in Cfr(-/-) mice. *Proc Natl Acad Sci* 91: 479-483.
5. Niisato N, Ito Y, Marunaka Y (1999) Activation of Cl⁻ channel and Na⁺/K⁺/2Cl⁻ cotransporter in renal epithelial A6 cells by flavonoids: genistein, daidzein, and apigenin. *Biochem Biophys Res Commun* 254: 368-371.
6. Diener M, Hug F (1996) Modulation of Cl⁻ secretion in rat distal colon by genistein, a protein tyrosine kinase inhibitor. *Eur J Pharmacol* 299: 161-170.
7. Goddard CA, Evans MJ, Colledge WH (2000) Genistein activates CFTR-mediated Cl⁻ secretion in the murine trachea and colon. *Am J Physiol Cell Physiol* 279: C383-392.
8. Chao PC, Hamilton KL (2009) Genistein stimulates electrogenic Cl⁻ secretion via phosphodiesterase modulation in the mouse jejunum. *Am J Physiol Cell Physiol* 297: C688-C698.
9. Tuo B, Wen G, Song P, Xu J, Liu X, et al. (2011) Genistein stimulates duodenal HCO₃⁻ secretion through PI3K pathway in mice. *Eur J Pharmacol* 651: 159-167.
10. Al-Nakkash L, Clarke LL, Rottinghaus GE, Chen YJ, Cooper K, et al. (2006) Dietary genistein stimulates anion secretion across female murine intestine. *J Nutr* 136: 2785-2790.
11. Kamboh AA, Zhu WY (2014) Individual and combined effects of genistein and hesperidin on immunity and intestinal morphology in lipopolysaccharide-challenged broiler chickens. *Poult Sci* 93: 2175-2183.
12. Clarke LL, Gawenis LR, Franklin CL, Harline MC (1996) Increased survival of CFTR knockout mice with an oral osmotic laxative. *Lab Anim Sci* 46: 612-618.
13. Kelly T, Buxbaum J (2015) Gastrointestinal Manifestations of Cystic Fibrosis. *Dig Dis Sci* 60: 1903-1913.
14. Al-Nakkash L, Drumm ML (2004) Dietary genistein increases survival in a CF mouse model. *Pediatr Pulmonol* 27: 250.
15. El-Salhy M (2002) Gastrointestinal transit in an animal model of human diabetes type 2: relationship to gut neuroendocrine peptide contents. *Ups J Med Sci* 107: 101-110.
16. El-Salhy M, Spangeus A (2002) gastric emptying in animal models of human diabetes: correlation to blood glucose level and gut neuroendocrine peptide content. *Ups J Med Sci* 107: 89-99.
17. Catmull S, Masood F, Schacht S, Dolan R, Stegman D (2016) Dietary genistein rescues reduced basal chloride secretion in diabetic jejunum via sex-dependent mechanisms. *Cell Physiol Biochem* 40: 335-346.
18. Asfaha S, MacNaughton WK, Appleyard CB, Chadee K, Wallace JL (2001) Persistent epithelial dysfunction and bacterial translocation after resolution of intestinal inflammation. *Am J Physiol* 281: G635-G644.