

Validated HPLC Method for the Determination of JWU1497 in Rat Plasma and Its Application to a Comparative Pharmacokinetic Study of the Free Base and Hydrophosphate Salt Forms of JWU1497

Tae Kon Kim* and Oui Hyun Jung

College of Science & Engineering, Jungwon University, 85 Munmu-ro, Geosan-gun, Chungbuk, South Korea

Abstract

In this study, a sensitive and reliable method for the quantitation of JWU1497 in rat plasma was developed and validated using high performance liquid chromatography (HPLC). The pharmacokinetics of 2 forms of JWU1497, namely the base form and the hydrophosphate salt form, were investigated in rats. The 2 forms were orally administered to rats and the plasma concentrations of JWU1497 were determined using HPLC. The JWU1497 base and hydrophosphate salt forms showed similar pharmacokinetic profiles in terms of their maximum plasma concentration (C_{max}) and area under the concentration-time curve (AUC). The time to peak concentration (T_{max}) of the base form was slightly greater than that of the salt form, but this difference was not statistically significant. These results suggest that the JWU1497 base and hydrophosphate forms are pharmacokinetically equivalent in rats, and thus the base form could be used in various JWU1497 formulations as a substitute for the existing JWU1497 hydrophosphate form.

Keywords: JWU1497; HPLC; Rat; Pharmacokinetics**Introduction**

Male erectile dysfunction, the present inability to achieve or maintain an erection for satisfactory sexual performance, is a common and important medical problem [1]. Recently, phosphodiesterase type 5 (PDE-5) inhibitors are used to improve erectile dysfunction by binding cyclic guanosine monophosphate and maintaining sufficient cellular levels in the smooth muscles [2,3]. However, PDE-5 inhibitors have common adverse reactions such as headache, flushing, nasal congestion, and dyspepsia [4]. Thus, JWU1497 (Figure 1), a new PDE-5 inhibitor, was developed to alleviate drawbacks of above common side effects of PDE-5 inhibitors. JWU1497 appears to be safe and effective in the treatment of male erectile dysfunction. The log partition coefficient (octanol / water) of JWU1497 was approximately 3.59. The solubilities of JWU1497 in methanol, acetonitrile, and distilled water were 150, 195, and 1.12 mg/ml, respectively, at $20 \pm 5^\circ\text{C}$. The IC_{50} value of JWU1497 (0.235 nM) for the inhibition of PDE-5 was smaller than other PDE-5 inhibitors such as sildenafil and tadalafil (an internal report). JWU1497 of which the active pharmaceutical ingredient (API) is JWU1497 hydrophosphate salts is about to enter the clinical study.


Generally, drugs in salt forms are preferable as they have higher dissolution rate, which facilitates absorption from the gastrointestinal tract [5]. However, the salt forms of drugs can have limitations when they are used in some formulations, such as gels, patches, films, or

chewable tablets, due to their greater volume compared with free base forms. Furthermore, salt forms may cause corrosion to equipment used in the manufacturing process. With the aim of developing a new JWU1497 formulation, free base form of JWU1497 was selected and used as the API instead of the existing salt form. The free base form negated the disadvantages of the salt form mentioned above but was also expected to improve the taste of the product, which is one of the important properties of drugs [6,7].

In this study, the pharmacological equivalence of 2 forms of JWU1497, the free base and hydrophosphate salt forms, was compared in the context of the development of JWU1497 new formulation that use JWU1497 free base as the API. We determined the JWU1497 concentration in the plasma after the administration of JWU1497 free base and hydrophosphate salt forms to rats by using HPLC and compared their pharmacokinetic properties.

Materials and Methods**Materials**

JWU1497 free base and hydrophosphate salt (JWU1497-HPO₄) were provided by Jungwon university (Chungbuk, Korea) with a chemical purity of more than 99%. Sildenafil, an internal standard for the HPLC analysis of JWU1497 were purchased from Sigma-Aldrich Corporation (St. Louis, MO). Acetonitrile and methanol were products



*Corresponding author: Tae Kon Kim, College of Science & Engineering, Jungwon University, 85 Munmu-ro, Geosan-gun, Chungbuk 367-700, South Korea, Tel: +82-43-830-8619; Fax: +82-43-830-8579; E-mail: tkkim@jwu.ac.kr

Received July 12, 2015; Accepted September 02, 2015; Published September 09, 2015

Citation: Kim TK, Jung OH (2015) Validated HPLC Method for the Determination of JWU1497 in Rat Plasma and Its Application to a Comparative Pharmacokinetic Study of the Free Base and Hydrophosphate Salt Forms of JWU1497. J Bioequiv Availab 7: 267-269. doi:10.4172/jbb.1000251

Copyright: © 2015 Kim TK, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

from Burdick & Jackson (Muskegon, MI, USA). Other chemicals were of reagent grade or HPLC grade.

Preparation of JWU1497 dosing solutions

JWU1497 free base and JWU1497-HPO₄ were dissolved in distilled water with 0.5% sodium carboxymethylcellulose to a final concentration of 3 mg/mL as free base.

Animal experiments

Male Sprague–Dawley rats, 6-8 week old and weighing 220-300 g, were purchased from the Samtako Bio Korea (Osan, South Korea). Rats were maintained in a Clean room at a temperature of between 23 ± 2°C with 12-h light (07:00-19:00) and dark (19:00-07:00) cycles, and a relative humidity of 55% ± 5%. Rats were housed in metabolic cages (Tecniplast, Varese, Italy) under filtered pathogen-free air and with food (Sam Yang Company, Pyeongtaek, South Korea) and water available ad libitum. The rats were fasted overnight before drug administration and for 4 hr after dosing. The rats were placed in a restrainer and were orally administered a dose of 30 mg/kg with a catheter. Blood was collected in a heparinized tube at the pre-dose stage, and at 0.25, 0.5, 1, 1.5, 2, 4, 6, 8, and 12 h after p.o. administration. Plasma was harvested after centrifugation at 3,000 rpm and 4°C for 10 min and stored frozen at -70°C until it was analyzed.

Preparation of calibration standards and quality control samples

Stock solutions of JWU1497 (1 mg/mL) were prepared in methanol. Appropriate dilutions of the stock solutions of JWU1497 were made with methanol (0.003, 0.005, 0.01, 0.02, 0.05, 0.1, 0.2, 0.5, or 1 mg/mL). Standard solutions of JWU1497 in rat plasma were prepared by spiking with an appropriate volume (10 µL/mL of plasma) of the diluted stock solutions, giving final concentrations of 0.03, 0.05, 0.1, 0.2, 0.5, 1, 2, 5, or 10 µg/mL for plasma. The IS working solution was prepared by dissolving sildenafil in acetonitrile to give a final concentration of 10 µg/mL.

Preparation of plasma samples

A 50 µL aliquot of sample was deproteinized with a 75 µL of acetonitrile containing 10 µg/mL sildenafil (an IS). After vortex-mixing and centrifugation at 3,000 rpm for 10 min, the supernatant was transferred into a vial and a 20 µL aliquot was injected directly onto the HPLC column.

HPLC analysis

The HPLC system consisted of a Gilson-234 autosampler (Gilson, Middleton, WI, USA), a Gilson 307 pump (Gilson), a Capcell PAK (C₁₈) column (250 mm × 4.6 mm, i.d.; particle size, 5 µm; Shiseido, Tokyo, Japan), a model UV-118 UV/VIS detector (Gilson), and a model Gilson unipoint system software (Gilson). The mobile phase, 0.02 M ammonium acetate buffer:acetonitrile (45:55, v/v), was run at a flow rate of 1.0 mL/min, and the column eluent was monitored using an ultraviolet detector at 254 nm at room temperature. The retention times of sildenafil (an internal standard) and JWU1497 were approximately 5.7 and 8.3 min, respectively.

Analytical method validation

The analytical method was validated with regards to its specificity, linearity, intra- and interday precision and accuracy, matrix effect, and stability according to the US Food and Drug Administration's "Guidance for Industry, Bioanalytical Method Validation, 2001 [8]".

Pharmacokinetic and statistical analyses

The total area under the plasma concentration-time curve to the last time (AUC_{last}), the maximum plasma concentration (C_{max}), the time to reach C_{max} (T_{max}), and the half-life (T_{1/2}) were estimated using non-compartmental calculations carried out within WinNonlin™ 5.2 (Pharsight, Sunnyvale, CA, USA). All data are expressed as the mean ± standard deviation (SD). The statistical significance of the differences between the 2 groups was analyzed using Student's *t*-tests carried out within SPSS (IBM, Yorktown Heights, NY, USA). A *p* value of <0.05 was considered statistically significant.

Results and Discussion

Development and validation of the HPLC method

The HPLC method for the determination of JWU1497 in rat plasma was developed and validated with regard to specificity, linearity, accuracy, and sensitivity. No interferences from endogenous substances were observed in the blank rat plasma samples. The retention times of sildenafil and JWU1497 were 5.7 and 8.3 min, respectively. The analytical method used was linear over the range of 0.03-10 µg/mL, with correlation coefficients (*r* values) greater than 0.9997. The lower limit of quantitation was 0.03 µg/mL with relative standard deviation (RSD) values less than 20% and relative errors within ± 20%. Intra- and inter-day accuracies (as relative error values) ranged between 1.0% and 11.5% and intra- and inter-day precisions (as RSDs) were 3.0-10.1% for all QC samples, with the result that they all met the criteria for bioanalysis method validation (Table 1). The matrix effect, recovery, and process efficiency values for JWU1497 and sildenafil in rat plasma are provided in (Table 2). The recovery was, on average, more than 90% for both compounds. JWU1497 was found to be stable under various

Nominal conc. (ng/mL)	Measured conc. (ng/mL)	Coefficient of variation (%)	Relative error (%)
Intra-day (n=6)			
30	30 ± 2	8.1	0.0
100	106 ± 6	5.2	6.0
1000	1109 ± 34	3.0	10.9
7500	7827 ± 315	4.0	4.4
Inter-day (n=18, 6 runs per day)			
30	32 ± 3	10.1	6.7
100	112 ± 8	7.4	12.0
1000	1081 ± 55	5.1	8.1
7500	7428 ± 438	5.9	-1.0

Data represent mean ± SD.

Coefficient of variation (%) = (SD/mean) × 100

Relative error (%) = ((Measured conc. - Nominal conc.) / Nominal conc.) × 100

Table 1: Intra- and inter-day precision and accuracy for JWU1497 in rat plasma QC samples.

Concentration (ng/mL)	Matrix effect (%) (B/A×100)	Recovery (%) (C/B×100)	Process efficiency (%) (C/A×100)
JWU1497			
100	72.5 ± 12.9	96.0 ± 2.9	69.3 ± 11.2
1000	71.6 ± 4.3	93.9 ± 6.3	67.3 ± 6.0
7500	82.6 ± 3.8	86.5 ± 2.1	71.4 ± 3.5
Sildenafil			
10000	73.8 ± 1.9	91.6 ± 2.6	67.6 ± 2.1
A, Peak area of analytes in mobile phase			
B, Peak area of analytes spiked after extraction			
C, Peak area of analytes spiked before extraction			

Table 2: Matrix effect, recovery, and process efficiency data for JWU1497 and sildenafil in rat plasma.

conditions, whether in the plasma or in the stock solution, and the detailed stability data are presented in (Table 3). In summary, the HPLC method developed in the current study was found to be suitable for the quantification of JWU1497 in plasma with acceptable specificity, linearity, accuracy, precision, recovery, and stability. On the basis of this HPLC method, JWU1497 concentrations in rat plasma were determined.

Comparative pharmacokinetics of JWU1497 free base and hydrophosphate salt forms in rat plasma

Plasma samples were collected after the oral administration of the free base and hydrophosphate salt forms of JWU1497 and the concentrations of the API, JWU1497, were determined using the validated HPLC method. Figure 2 shows the mean plasma concentration-time curves for JWU1497 after the oral administration of the 2 JWU1497 formulations in rats; the pharmacokinetic parameters are presented in Table 4. The maximum plasma concentrations of JWU1497 were achieved 1.1 and 0.6 hr after oral administration for the free base and hydrophosphate forms, respectively. The C_{max} values were 6.31 ± 2.90 and 6.25 ± 2.85 $\mu\text{g/mL}$, and the AUC_{last} values were 22.25 ± 9.79 and 21.85 ± 8.95 $\mu\text{g}\cdot\text{h/mL}$ for the free base and salts forms, respectively. The C_{max} and AUC_{last} values for these 2 groups were comparable. The free base form of JWU1497 appeared to have been absorbed more slowly from the




Figure 2: Mean plasma concentration-time curves for JWU1497 after oral administration of JWU1497 free base and hydrophosphate salt forms at a dose of 30 mg/kg as free base in rats. Each point represents the mean \pm standard deviation ($n = 5$).

gastrointestinal tract than the hydrophosphate salt form, but there was no statistically significant difference between the pharmacokinetic profiles of the 2 groups.

Conclusions

The HPLC method was developed and validated for the determination of JWU1497 in rat plasma and developed method was successfully applied to a comparative pharmacokinetic study of the free base and salt forms of JWU1497. The pharmacokinetic profile of the free base form was comparable to that of hydrophosphate salt form in rats. This suggests that these 2 forms could be used interchangeably to produce a variety of pharmaceutical preparations.

Acknowledgement

This work was supported by the J2H Grants.

References

- Hatzimouratidis K, Hatzichristou DG (2008) Looking to the future for erectile dysfunction therapies. *Drugs* 68: 231-250.
- Lincoln TM (2004) Cyclic GMP and phosphodiesterase 5 inhibitor therapies: what's on the horizon? *Mol Pharmacol* 66: 11-13.
- Aversa A, Bruziches R, Pili M, Spera G (2006) Phosphodiesterase 5 inhibitors in the treatment of erectile dysfunction. *Curr Pharm Des* 12: 3467-3484.
- Broderick GA (2003) Oral pharmacotherapy and the contemporary evaluation and management of erectile dysfunction. *Rev Urol* 5 Suppl 7: S9-9S20.
- Engel GL, Farid NA, Faul MM, Richardson LA, Winneroski LL (2000) Salt form selection and characterization of LY333531 mesylate monohydrate. *Int J Pharm* 198: 239-247.
- Dixit RP, Puthli SP (2009) Oral strip technology: overview and future potential. *J Control Release* 139: 94-107.
- Goel H, Rai P, Rana V, Tiwary AK (2008) Orally disintegrating systems: innovations in formulation and technology. *Recent Pat Drug Deliv Formul* 2: 258-274.
- <http://www.fda.gov/downloads/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/ucm070107.pdf>

Nominal conc. (ng/mL)	Duration	Measured conc. (ng/mL)	Relative error (%)
Short-term stability (at room temperature, RT)			
100	4 h	103 \pm 7	3.0
1000		1031 \pm 35	3.1
7500		7626 \pm 324	1.7
Long-term stability (at -80°C)			
100	7 days	110 \pm 10	10.0
1000		1076 \pm 46	7.6
7500		7703 \pm 418	2.7
Freeze and thaw stability			
100	3 cycles	105 \pm 8	5.0
1000		1070 \pm 41	7.0
7500		8010 \pm 381	6.8
Auto-sampler stability (at 4°C)			
100	24 h	104 \pm 8	4.0
1000		1054 \pm 51	5.4
7500		7866 \pm 639	4.9
Stock solution			
500	2 h at RT	510 \pm 5	2.0
	11 days at 4°C	490 \pm 4	-2.0

Table 3: Stability of JWU1497 in rat plasma and stock solutions ($n=3$).

Parameters	JWU1497 base ($n=5$)	JWU1497-HPO ₄ ($n=5$)
AUC_{last} ($\mu\text{g}\cdot\text{hr/mL}$)	22.25 \pm 9.79	21.85 \pm 8.95
C_{max} ($\mu\text{g/mL}$)	6.31 \pm 2.90	6.25 \pm 2.85
T_{max} (hr)	1.10 \pm 0.22	0.60 \pm 0.22
$T_{1/2}$ (hr)	1.76 \pm 0.69	1.52 \pm 0.78

Data represent mean \pm SD ($n=5$).

AUC: Area under the curve to the collected time point ($\mu\text{g}\cdot\text{hr/mL}$).

C_{max} : Peak plasma concentration ($\mu\text{g/mL}$)

T_{max} : Time to reach peak plasma concentration (hr)

$T_{1/2}$: Elimination half life (hr)

Table 4: Pharmacokinetic parameters of JWU1497 after a single oral administration of JWU1497 base and hydrophosphate salts (JWU1497-HPO₄) forms at a dose of 30 mg/kg (as a base form) to male rats.

Citation: Kim TK, Jung OH (2015) Validated HPLC Method for the Determination of JWU1497 in Rat Plasma and Its Application to a Comparative Pharmacokinetic Study of the Free Base and Hydrophosphate Salt Forms of JWU1497. *J Bioequiv Availab* 7: 267-269. doi:10.4172/jbb.1000251