

# Water-mediated Desymmetrization Reactions

Satomi Niwayama\*

Graduate School of Engineering, Muroran Institute of Technology, 27-1, Mizumoto-cho, Muroran, Hokkaido, Japan

The development of environmentally friendly organic reactions has been of central importance in synthetic organic chemistry. As water is among the most environmentally friendly solvents, water-mediated organic reactions thus represent a typical “green chemistry.” Among various synthetic conversions, desymmetrization of symmetric compounds is one of the most cost-effective reactions, because the starting symmetric compounds are typically obtained on a large scale from inexpensive sources, or are commercially available inexpensively. Therefore, water-mediated desymmetrization of symmetric organic compounds is of remarkable synthetic values.

However, distinguishing the identical functional groups in symmetric compounds is considered challenging. Organic compounds are also generally considered not to significantly dissolve in water due to their hydrophobic nature. Therefore, the studies on water-mediated desymmetrization reactions have been rather limited. This editorial highlights some of the recent progress in the research of water-mediated desymmetrization reactions.

Enzyme reactions are a relatively common approach to accomplishing desymmetrization in aqueous media [1-8]. Conditions for enzyme reactions are typically mild and environmentally friendly, and therefore this approach is considered green chemistry. However, the mechanisms of enzyme reactions are mostly unknown, in particular, in the area of desymmetrization, and therefore one must screen various enzymes and substrates in order to reach desirable outcomes.

Among the most well-known non-enzymatic desymmetrization reactions are ring opening reactions of strained cyclic compounds such as epoxides and aziridines [9-14]. The driving force of these reactions is the relief of bond angle strain. Some asymmetric versions are also reported in this area.

Without such a driving force, desymmetrization is more difficult. Perhaps among the most pioneering non-enzymatic water-mediated desymmetrization are selective monohydrolysis of symmetric diesters reported by Niwayama [15-23]. This reaction distinguishes the two identical ester groups with high selectivity (>99%) and with high yields (>99%) in water, a selectivity which was previously impossible in the organic media. The reaction mechanism remains unknown, but potential conformational bias due to attractive interactions between two closely located carboalkoxy groups in these diesters may have contributed for this selectivity, because especially high selectivities were observed in the symmetric diesters with “*cis*” stereochemistry, facilitating the hydrolysis of a carboalkoxy group from one particular direction. The existence of the  $n \rightarrow \pi^*$  interaction is suggested by theoretical studies for such an interaction [16]. An additional proposed mechanistic insight includes formation of micelle-like aggregates in which the hydrophobic carboalkoxy groups are pointed inside and hydrophilic COO<sup>-</sup> groups are pointed outside, prohibiting further hydrolysis by the aqueous NaOH solution [23].

Chong et al. also reported selective monobromination of symmetric diols in the aqueous HBr/toluene two-phase media with high selectivities, producing high (>90%) yields of monobromides, despite the fact that the expected yields are a maximum of 50% as in the above monohydrolysis [24]. They propose that the reversed micelle-

like structure by the bromo alcohol in which the polar hydroxy groups are pointed inside, which is expected to keep away from the exposure from the aqueous HBr.

More recently, other non-enzymatic desymmetrization reactions in aqueous media such as selective monobenzoylation of 1,2- and 1,3-diols [25], selective monocycloisomerization of bispropargylic carboxylic acids [26], desymmetrization of 1-ferrocenedicarboxaldehyde [27], and aldol reactions toward cyclohexanone [28] have been reported. The key intermediate for the mechanism of the first reaction is proposed to involve a cyclic stannyl structure [25]. The transition state involving the enamines formed by the proline-based chiral amines is proposed for the last reaction [28]. Yields vary for all these reactions, but these reactions were performed in aqueous media with the use of water as a main solvent or a co-solvent. However, considering the obscure nature of the desymmetrization and the difficulty of the elucidation of the mechanisms, the reported studies and cited references in this editorial are expected to stimulate organic chemists and help pave the way for further development of additional green reactions.

## References

1. Yu MS, Lantos I, Peng ZQ, Yu J, Cacchio T (2000) Asymmetric Synthesis of (-)-Paroxetine Using PLE Hydrolysis. *Tetrahedron Lett* 41: 5647-5651.
2. Sano S, Ushirogouchi H, Morimoto K, Tamai S, Nagao Y (1996) A New Enantiodivergent Procedure Utilizing the Chemoselective Dieckmann-type Cyclization of Chiral Mono-thiol Diesters. *Chem Commun* 27: 1775-1776.
3. Schoffers E, Golebiowski A, Johnson CR (1996) Enantioselective Synthesis through Enzymatic Asymmetrization. *Tetrahedron* 56: 3769-3826.
4. Guanti G, Narisano E, Riva R (1998) Lipase Mediated Preparation of Differently Protected Homochiral 2-Aryl-2-fluoro-1,3-propanediols. *Tetrahedron Asymmetry* 9: 1859-1862.
5. Guanti G, Banfi L, Brusco S, Narisano E (1994) Enzymatic Preparation of Homochiral 2-(N-carbobenzyloxypiperid-4-yl)-1,3-propanediol Monoacetate. A Facile Entry to Both Enantiomers of 3-Hydroxymethylquinuclidine *Tetrahedron Asymmetry* 5: 537-540.
6. Guanti G, Banfi L, Narisano E (1990) Enzymes in Organic Synthesis: Remarkable Influence of  $\pi$  System on the Enantioselectivity in PPL Catalyzed Monohydrolysis of 2-Substituted 1,3-Diacetoxypropanes. *Tetrahedron Asymmetry* 1: 721-724.
7. Guanti G, Narisano E, Podgorski T, Thea S, Williams A (1990) Enzyme Catalyzed Monohydrolysis of 2-Aryl-1,3-propanediol Diacetates. A Study of Structural Effects of the Aryl Moiety on the Enantioselectivity *Tetrahedron* 46: 7081-7092.
8. Ohno M, Otsuka M (1989) Chiral Synthons by Ester Hydrolysis Catalyzed by Pig Liver Esterase. *Org React* 37: 1-55.

\*Corresponding author: Satomi Niwayama, Graduate School of Engineering, Muroran Institute of Technology, 27-1, Mizumoto-cho, Muroran, Hokkaido, 050-8585, Japan, Tel: 806-742-3118; E-mail: [niwayama@mmm.muroran-it.ac.jp](mailto:niwayama@mmm.muroran-it.ac.jp)

Received September 01, 2014; Accepted September 03, 2014; Published September 05, 2014

Citation: Niwayama S (2014) Water-mediated Desymmetrization Reactions. *Organic Chem Curr Res* 1:e138. doi: [10.4172/2161-0401.1000e138](https://doi.org/10.4172/2161-0401.1000e138)

Copyright: © 2014 Niwayama S. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

9. White DE, Tadross PM, Lu Z, Jacobsen EN (2014) A Broadly Applicable and Practical Oligomeric (salen) Co Catalyst for Enantioselective Epoxide Ring-opening Reactions. *Tetrahedron* 70: 4165-4180.
10. Li J, Liao Y, Zhang Y, Liu XH, Lin LL, et al. (2014) Chiral Magnesium (II)-catalyzed Asymmetric Ring-opening of meso-Aziridines with Primary Alcohols. *Chem Commun* 50: 6672-6674.
11. Ogawa C, Kokubo M, Kobayashi S (2010) Asymmetric Ring-Opening Reactions of meso-Epoxides Using Metal-Chiral Bipyridine Complexes. *Journal of Synthetic Organic Chemistry Japan* 68: 718-728.
12. Kokubo M, Naito T, Kobayashi S (2010) Chiral Zinc (II) and Copper (II)-Catalyzed Asymmetric Ring-opening Reactions of meso-Epoxides with Aniline and Indole Derivatives. *Tetrahedron* 66: 1111-1118.
13. Bonollo S, Fringuelli F, Pizzo F, Vaccaro L (2008) Zn (II)-catalyzed Desymmetrization of meso-Epoxides by Aromatic Amines in Water. *Synlett* 10: 1574-1578.
14. Ogawa C, Wang N, Kobayashi S (2007) Scandium-catalyzed Ring-opening Desymmetrization of meso-Epoxides. *Heterocycles* 36: 34-35.
15. Niwayama S (2000) Highly Efficient Selective Monohydrolysis of Symmetric Diesters. *J Org Chem* 65: 5834-5836.
16. Cho H, Alexander RB, Niwayama S (2012) Conformational Analysis of Symmetric Diesters. *Curr Org Chem* 16: 1151-1158.
17. Niwayama S, Wang H, Hiraga Y, Clayton JC (2007) Influence of co-solvents in the highly efficient selective monohydrolysis of a symmetric diester. *Tetrahedron Lett* 48: 8508-8510.
18. Niwayama S, Rimkus A (2005) Effects of Counter Cations in Selective Monohydrolyses of Symmetric Diesters. *Bull Chem Soc Jpn* 78: 498-500.
19. Niwayama S, Cho H, Lin C (2008) Highly Efficient Selective Monohydrolysis of Dialkyl Malonates and Their Derivatives. *Tetrahedron Lett* 49: 4434-4436.
20. Niwayama S, Cho H (2009) Practical Large Scale Synthesis of Half-esters of Malonic Acid. *Chem Pharm Bull* 57: 508-510.
21. Niwayama S, Cho H, Zabet-Moghaddam M, Whittlesey BR (2010) Remote Exo/Endo Selectivity in Selective Monohydrolysis of Dialkyl Bicyclo[2.2.1]heptane-2,3-dicarboxylate Derivatives. *J Org Chem* 75: 3775-3780.
22. Niwayama S, Hiraga Y (2003) New Exo/Endo Selectivity Observed in Monohydrolysis of Dialkyl Bicyclo[2.2.1]hept-5-ene-2,3-dicarboxylates. *Tetrahedron Lett* 44: 8567-8570.
23. Niwayama S (2008) Highly Efficient and Practical Selective Monohydrolysis of Symmetric Diesters: Recent Progress and Scope. *J Synth Org Chem Jpn* 66: 983-994.
24. Chong JM, Heuft MA, Rabbat P (2000) Solvent Effects on the Monobromination of  $\alpha, \omega$ -Diols. A Convenient Preparation of  $\omega$ -Bromoalkanols. *J Org Chem* 65: 5837-5838.
25. Muramatsu, W, William JM, Onomura O (2012) Selective Monobenzylation of 1,2- and 1,3-Diols Catalyzed by  $\text{Me}_2\text{SnCl}_2$  in Water (Organic Solvent Free) under Mild Conditions. *J Org Chem* 77: 754-759.
26. Tomás-Mendivil E, Toullec PY, Díez J, Conejero S, Michelet V, et al. (2012) Cycloisomerization versus Hydration reactions in Aqueous Media: A Au(III)-NHC Catalyst That Makes the Difference. *Org Lett* 14: 2520-2523.
27. Benniston AC, Sirbu D, Turta C, Probert MR, Clegg W (2014) A Simple Method for Desymmetrizing 1,1'-Ferrocenedicarboxaldehyde. *Tetrahedron Lett* 55: 3777-3780.
28. Luo S, Xu H, Li J, Zhang L, Mi X, et al. (2007) Facile evolution of asymmetric organocatalysts in water assisted by surfactant Brønsted acids. *Tetrahedron* 63: 11307-11314.

Citation: Niwayama S (2014) Water-mediated Desymmetrization Reactions. Organic Chem Curr Res 1:e138. doi: 10.4172/2161-0401.1000e138

### Submit your next manuscript and get advantages of OMICS Group submissions

#### Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

#### Special features:

- 300 Open Access Journals
- 30,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, DOAJ, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscripts as E- mail: [editor.organicchemistry@omicsonline.org](mailto:editor.organicchemistry@omicsonline.org)