

Work-Related Stress and Stress-Coping Strategies among Patients' Companions at a Tertiary Care Hospital in Saudi Arabia

Fahad D Alosaimi^{1,2}, Ayedh H Alghamdi², Bandar S Aladwani², Sana N Kazim³ and Auroabah S Almufleh³

¹SABIC Psychological Health Research & Applications Chair, College of Medicine, King Saud University, Riyadh, Saudi Arabia

²Department of Psychiatry, King Saud University, Riyadh, Saudi Arabia

³Department of Psychiatry, Saudi Commission for Health Specialties, Riyadh, Saudi Arabia

Abstract

Providing care to hospitalized patients may be associated with stress, reduced quality of life, and even psychiatric disorders. This case-control study aimed to compare the level of stress (using the 14-question perceived stress scale (PSS-14), potential risk factors, and stress-coping strategies (using the 28-item brief coping scale (BCS-28) between two convenient samples of patients' companions (56) and administrative employees (98) working in a tertiary care hospital in Saudi Arabia. The average PSS-14 was slightly higher in companions than administrative employees (27.4 ± 9.9 vs. 25.1 ± 10.1 , $p=0.179$). After stratifying by gender, the difference in males (but not females) was marginally significant in the unadjusted comparison and significant in the adjusted comparison. Companions had similar scores of adaptive stress-coping strategies but higher scores of maladaptive stress-coping strategies compared with administrative employees. This was especially apparent in the denial and self-distraction strategies. PSS-14 in all participants had a moderate significant positive correlation with maladaptive stress-coping strategies. Companions exhibited a gender-specific slight increase in the level of stress and adopted more maladaptive stress-coping strategies compared with administrative employees. There were no group differences in blood lipids, serum glucose, or cortisol levels.

Keywords: Companions; Administrative; Stress; Coping; Gender; Saudi Arabia

Introduction

The concept of "constant observation" refers to continuous one-to-one monitoring that is used to assure the safety and well-being of patients [1]. In the West, this is usually achieved by allocated nurses, hired or volunteer non-relative personnel, or patient family members [2]. In Saudi Arabia, a companion refers to the person (usually a family member or a friend) who provides "constant observation" for a patient during his/her stay at the hospital [3,4]. In the literature, several terminologies are used interchangeably with companion, such as patient sitter and patient attendant [5]. The companion policy at King Khalid University Hospital (KKUH) defines a companion as a healthy adult (more than 18 years of age) who is not overtly pregnant or taking care of infants. The policy allows one companion for children, senior and disabled patients, but does not allow companions for patients in critical care areas or for isolated patients due to the risk of infection and medical-related issues.

Several studies showed that family caregiving to patients with different disabling diseases are associated with high levels of distress, reduced quality of life, and even psychiatric disorders among the caregivers [6-8]. On the other hand, a limited number of studies examined the psychological impact of caregiving on the companions of patients with chronic and aging diseases admitted to hospitals [9-11]. Although this type of data is critical for any future attempt to improve the effectiveness of the current companion policy [3,12], data examining stress and stress-coping strategies among patient companions in Saudi Arabia are largely lacking. The objective of the current study was to compare the level of stress, potential risk factors, and stress-coping strategies among companions in a tertiary care hospital in Saudi Arabia. Similar to previous studies, administrative employees working in the same hospital were used as a control group [13].

Methods

Population

The current study was conducted among companions and

administrative employees at King Khalid University Hospital (KKUH) located in Riyadh, Saudi Arabia. KKUH is an 800-bed University-affiliated government hospital that provides free primary to tertiary care services to residents of the Northern Riyadh area. The KKUH provides inpatient services for more than 50,000 patients per year. Companions who were non-healthcare sitters/caregivers of admitted patients were invited to join the study during their stay in the hospital. The KKUH hired approximately 3,000 administrative employees at the time of the study. Administrative employees with no clinical responsibilities or direct patient contact, such as accountants and receptionists, were invited to join the study.

Study design

A case-control study was carried out between December 2013 and June 2014. The study obtained all of the required ethical approvals from the Institutional Review Board at the Faculty of Medicine at King Saud University, Riyadh, Saudi Arabia.

Sample size and recruitment

Previous studies that used the 14-question perceived stress scale (PSS-14) showed that the average score was 25 ± 8 points [14]. Having two administrative employees for each companion, it was estimated

***Corresponding author:** Fahad D Alosaimi, SABIC Psychological Health Research & Applications Chair, College of Medicine, King Saud University, P.O. Box 7805, Riyadh 11472, Kingdom of Saudi Arabia, Tel: +96614671226/+966503441370; Fax: +966-14672571; E-mail: dr.fahad.alosaimi@gmail.com

Received April 17, 2016; Accepted June 21, 2016; Published June 27, 2016

Citation: Alosaimi FD, Alghamdi AH, Aladwani BS, Kazim SN, Almufleh AS (2016) Work-Related Stress and Stress-Coping Strategies among Patients' Companions at a Tertiary Care Hospital in Saudi Arabia. J Psychiatry 19: 374 doi:10.4172/2378-5756.1000374

Copyright: © 2016 Alosaimi FD, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

that 48 companions and 96 administrative employees are required to detect a 4 point difference (23-27) between the groups, with 80% power and a level of significance of 95%. The study participants were a convenient sample of companions who were personally invited to join the study. An age-matched control group of administrative employees who were working at KCUH during the study period were invited to join the study, either personally or by email.

Data collection tool

The self-administered questionnaire that was developed included socio-demographic characteristics, clinical history, occupational characteristics, and recently faced stressors. The perception of stress over the past month was assessed using PSS-14 [15,16]. Each question was answered on a 5-point Likert-type scale (never, almost never, sometimes, fairly often, or very often) and was given a score (ranging from 0 to 4). The PSS-14 score was calculated by summing the scores of individual questions, with higher scores representing higher levels of stress (possible range between 0 and 56 points). The scores of seven questions (4, 5, 6, 7, 9, 10, and 13) were reversed as they were positively stated. Stress-coping strategies employed in the past month were assessed using the 28-item brief coping scale (BCS-28) [17]. Each question was answered on a 4-point Likert-type scale (not at all, a little bit, a medium amount, or a lot). Each question was given a score ranging from 1 to 4 and each strategy was given a score ranging from 2 to 8. The scale assesses 14 stress-coping strategies using 28 questions (2 questions per strategy). Stress-coping strategies were grouped into adaptive or maladaptive strategies, as described previously [18]. Standing height (in cm) and weight (in kg) while the participant is wearing light clothes with bared feet were measured and used in calculating BMI ($\text{height}/\text{weight}^2$). Waist circumference (in cm) at the level of umbilicus was measured while the participant is standing and comfortably breathing. Blood pressure was recorded by using a standard mercury sphygmomanometer after the subject seats for at least 5 minutes. Two readings will be taken with 5 minutes interval and their mean will be recorded. For biochemical investigations, a venous blood sample was taken in the morning after an eight hours fasting for the determination of blood lipids (Low density lipoprotein (LDL) cholesterol, high density lipoprotein (HDL) cholesterol, total cholesterol, and triglycerides), fasting blood glucose, and cortisol.

Validation of data collection tool

Because the majority of companions and administrative employees do not speak English, the study questionnaire and both scales were first translated into Arabic by a linguistic specialist, fluent in both English and Arabic. Then, another specialist, fluent in both English and Arabic, carried out back translation into English. During this time, the back-translation and the original scale were compared and any differences were discussed and resolved. The content of the study questionnaire was validated by experts in psychiatry, epidemiology and ethics to ensure the relevance and applicability of different questions. The study questionnaire and both scales were then piloted on 15 individuals before the study began. The wording and suggested answers were modified for some questions based on the feedback from the pilot sample. Both PSS-14 [19,20] and BCS-28 [21] have been shown to be valid tools for perceived stress and stress-coping strategies, respectively. In the current study, the overall PSS-14 as well as the adaptive and non-adaptive BCS-28 had acceptable internal consistency among their items, as indicated by overall Cronbach Alpha values of 0.83, 0.77, and 0.64, respectively.

Statistical analysis

Data were presented as frequencies and percentages for categorical

data and mean and standard deviation (SD) for continuous data. Significant differences between companions and administrative employees in PSS-14, BCS-28, socio-demographic characteristics, clinical characteristics, and occupational characteristics were tested using Student's t-test for continuous data and chi-square test or Fisher's exact test (as appropriate) for categorical data. The adjusted marginal means of PSS by study groups and gender were calculated using a general linear model. The correlations between PSS-14 and BCS-28 were examined using Spearman's correlation. All p-values were two-tailed. A p-value <0.05 was considered significant. SPSS software (release 20.0, SPSS Inc., Chicago, U.S.) was used for the statistical analyses.

Results

A total of 154 participants (56 companions and 98 administrative employees) completed the study questionnaire and both the PSS-14 and coping scales. As shown in Table 1, the average age was 28.5 ± 4.5 years with almost two-thirds (64%) of the participants below the age of 30. Females represented 57% of the participants, with more females among the companions than the administrative employees (74% vs. 47%). All participants were Saudi, and approximately half (51%) of them were currently married. Obesity parameters including body mass index (BMI) (31.3 ± 6.7 vs. 26.6 ± 5.2), waist circumference (86.6 ± 14.8 vs. 79.6 ± 13.5), and the prevalence of obesity (56.1% vs. 24.5%) were much higher in the companions compared with the administrative employees. There were more current smokers among the administrative employees (26%) compared with the companions (11%). Approximately 34% of the participants had chronic medical illnesses, which primarily included obesity, diabetes, hypertension, and hyperlipidemia. Approximately 22% of the participants had a past history of psychiatric illness; 19% received psychiatric help and 11% had a family history of psychiatric illness. On examination, 29% of the participants had a psychiatric illness, which primarily included major depressive episodes, panic disorder, and generalized anxiety disorder. The average SBP was 115.1 ± 13.2 mmHg and the average DBP was 75.6 ± 8.7 mmHg. Compared with the administrative employees, the companions were older ($p=0.009$), more likely to be female ($p=0.002$), and had higher BMI ($p<0.001$), waist circumference ($p=0.003$), prevalence of obesity (<0.001), and history of hypertension ($p=0.024$); however, the companions were less likely to smoke ($p=0.024$) and had lower SBP ($p=0.007$). There were no group differences in blood lipids, serum glucose, or cortisol levels.

As shown in Table 2, approximately 55% of the participants had a monthly income between SR 5,000 and SR 9,999. While the income was lower in the companions compared with the administrative employees, both were equally satisfied (57%) with their income. The majority of the participants were sleeping less than 7 hours per day (63%) and working less than 8 hours per day (59%). More than half (55%) of the participants considered their job stressful and the majority (54%) had (always/sometimes) thoughts of changing their job or specialty. The majority (87%) of the participants reported facing stressors during the last month. These included work-related stressors (34%) followed by financial (33%), family (31%), and other stressors. Only 33% of the participants had training in stress management. Approximately 10% of the participants had (always/sometimes) suicidal thoughts and 4% had thoughts of harming themselves. Compared to the administrative employees, the companions had lower income ($p=0.048$), worked fewer hours ($p<0.001$), faced less work-related stressors ($p<0.001$), had more thoughts about changing their job/specialty ($p=0.001$), and had more suicidal thoughts (0.006).

	Companions (N=56)	Administrative employees (N=98)	Total (N=154)	p-value
Age (years)				
Mean ± SD	29.3 ± 4.4	28.1 ± 4.6	28.5 ± 4.5	0.098
<27	19 (33.9%)	37 (38.5%)	56 (36.8%)	0.009
27-29	9 (16.1%)	32 (33.3%)	41 (27.0%)	
>29	28 (50.0%)	27 (28.1%)	55 (36.2%)	
Gender				
Male	15 (26.3%)	54 (52.9%)	69 (43.4%)	0.002
Female	42 (73.7%)	48 (47.1%)	90 (56.6%)	
BMI				
Mean ± SD	31.3 ± 6.7	26.6 ± 5.2	28.4 ± 6.2	<0.001
Normal	9 (15.8%)	44 (43.1%)	53 (33.3%)	<0.001
Overweight	16 (28.1%)	33 (32.4%)	49 (30.8%)	
Obese	32 (56.1%)	25 (24.5%)	57 (35.8%)	
Waist circumference (cm)	86.6 ± 14.8	79.6 ± 13.5	82.1 ± 14.3	0.003
Marital Status				
Married	30 (53.6%)	48 (49.0%)	78 (50.6%)	0.341
single	22 (39.3%)	47 (48.0%)	69 (44.8%)	
Divorced	4 (7.1%)	3 (3.1%)	7 (4.5%)	
Medical and behavioral history				
Comorbidity	22 (39.3%)	31 (31.6%)	53 (34.4%)	0.38
Diabetes	6 (10.7%)	3 (3.1%)	9 (5.8%)	0.073
Hypertension	5 (8.9%)	1 (1.0%)	6 (3.9%)	0.024
Hyperlipidemia	1 (1.8%)	5 (5.1%)	6 (3.9%)	0.417
Obesity	4 (7.1%)	12 (12.2%)	16 (10.4%)	0.318
Others	9 (16.1%)	14 (14.3%)	23 (14.9%)	0.765
Current smoking	6 (10.7%)	25 (26.0%)	31 (20.4%)	0.024
History of psychiatric illness				
Personal history of psychiatric illness	11 (19.6%)	22 (22.7%)	33 (21.6%)	0.66
Received psychiatric help	10 (17.9%)	19 (19.6%)	29 (19.0%)	0.792
Family history of psychiatric illness	7 (12.7%)	10 (10.5%)	17 (11.3%)	0.682
Diagnosis of psychiatric illness				
No	37 (67.3%)	70 (72.9%)	107 (70.9%)	0.463
Yes	18 (32.7%)	26 (27.1%)	44 (29.1%)	
Major depressive episode	13 (23.6%)	14 (14.6%)	27 (17.9%)	0.162
Panic disorder	2 (3.6%)	4 (4.2%)	6 (4.0%)	1
Generalized anxiety disorder	0 (0.0%)	2 (2.1%)	2 (1.3%)	0.534
Others	3 (5.5%)	6 (6.2%)	9 (6.0%)	1
Blood pressure measurement				
Average SBP (mmHg)	111.3 ± 12.2	117.3 ± 13.4	115.1 ± 13.2	0.007
Average DBP (mmHg)	75.5 ± 8.0	75.7 ± 9.1	75.6 ± 8.7	0.905
Blood examination				
LDL cholesterol (nmol/L)	2.39 ± 0.71	2.49 ± 0.82	2.45 ± 0.78	0.477
HDL cholesterol (nmol/L)	1.23 ± 0.32	1.25 ± 0.36	1.24 ± 0.34	0.766
Triglycerides (nmol/L)	1.13 ± 0.55	1.06 ± 0.79	1.09 ± 0.71	0.543
Total cholesterol (nmol/L)	4.76 ± 0.84	4.79 ± 0.86	4.78 ± 0.85	0.791
Fasting blood glucose (nmol/L)	5.05 ± 1.78	4.80 ± 2.96	4.89 ± 2.59	0.572
Cortisol (nmol/L)	411.4 ± 169.4	392.7 ± 151.7	399.5 ± 158.1	0.483

Table 1: Demographic and clinical characteristics by study groups.

Table 3 shows the actual answers to the PSS-14 questions from both companions and administrative employees. Some responses were similar in both groups. For example, more than 60% of the participants in both groups were fairly/very often thinking about things that they needed to accomplish. On the other hand, several questions were answered with opposing responses in each of the groups. For example, 64% of the companions, compared with 46% of the administrative employees, were fairly/very often feeling nervous and stressed.

As shown in Figure 1, the average PSS-14 was slightly higher in companions than administrative employees (27.4 ± 9.9 vs. 25.1 ± 10.1); however, the difference did not reach statistical significance ($p=0.179$). After adjusting for the significant potential risk factors (shown in the univariate analysis above, which includes demographic, clinical, and work-related characteristics), the difference in PSS-14 widened and the p-value became marginally significant (29.7 vs. 25.5 , $p=0.074$). Furthermore, after stratifying by gender, the difference in males (but

	Companions (N=56)	Administrative employees (N=98)	Total (N=154)	p-value
Monthly income (SR)				
< 5,000	15 (27.8%)	10 (10.2%)	25 (16.4%)	0.048
5,000 – 9,999	25 (46.3%)	59 (60.2%)	84 (55.3%)	
10,000 – 14,999	10 (18.5%)	21 (21.4%)	31 (20.4%)	
≥15,000	4 (7.4%)	8 (8.2%)	12 (7.9%)	
Satisfied with the income				
Agree/strongly agree	30 (54.5%)	56 (58.3%)	86 (57.0%)	0.87
Not sure	7 (12.7%)	10 (10.4%)	17 (11.3%)	
Disagree/strongly disagree	18 (32.7%)	30 (31.2%)	48 (31.8%)	
Average sleeping per day				
<7 hours	33 (60.0%)	63 (64.3%)	96 (62.7%)	0.131
7-8 hours	18 (32.7%)	34 (34.7%)	52 (34.0%)	
> 8 hours	4 (7.3%)	1 (1.0%)	5 (3.3%)	
Working hours per day				
≤8 hours	46 (83.6%)	44 (44.9%)	90 (58.8%)	<0.001
>8 hours	9 (16.4%)	54 (55.1%)	63 (41.2%)	
Working weekends				
No	29 (55.8%)	69 (70.4%)	98 (65.3%)	0.073
Yes	23 (44.2%)	29 (29.6%)	52 (34.7%)	
Consider job stressful				
Agree/strongly agree	28 (50.0%)	56 (57.1%)	84 (54.5%)	0.657
Not sure	9 (16.1%)	12 (12.2%)	21 (13.6%)	
Disagree/strongly disagree	19 (33.9%)	30 (30.6%)	49 (31.8%)	
Facing stressors during last month				
No	7 (12.5%)	13 (13.3%)	20 (13.0%)	1
Yes	49 (87.5%)	85 (86.7%)	134 (87.0%)	
Work	9 (16.1%)	43 (43.9%)	52 (33.8%)	<0.001
Financial	17 (30.4%)	33 (33.7%)	50 (32.5%)	0.672
Family	21 (37.5%)	27 (27.6%)	48 (31.2%)	0.2
Marital	7 (12.5%)	13 (13.3%)	20 (13.0%)	0.892
Academic	4 (7.1%)	13 (13.3%)	17 (11.0%)	0.243
Death of loved person	5 (8.9%)	11 (11.2%)	16 (10.4%)	0.653
Household sick person	8 (14.3%)	5 (5.1%)	13 (8.4%)	0.069
Trained in stress management				
No	39 (69.6%)	63 (65.6%)	102 (67.1%)	0.611
Yes	17 (30.4%)	33 (34.4%)	50 (32.9%)	
Thoughts/wishes*				
To change job/specialty	40 (71.4%)	43 (43.9%)	83 (53.9%)	0.001
To die	11 (19.6%)	5 (5.1%)	16 (10.4%)	0.006
To harm own-self	3 (5.5%)	3 (3.1%)	6 (3.9%)	0.668

*Always or sometimes

Table 2: Work-related characteristics, stressors, and ideations by study groups.

not females) was marginally significant ($p=0.055$) in the unadjusted comparison and was significant ($p=0.036$) in the adjusted comparison.

As shown in Table 4, the companions had similar scores of adaptive stress-coping strategies but higher scores of maladaptive stress-

coping strategies compared with the administrative employees. This was especially apparent in the higher denial and self-distraction scores in the companions ($p=0.006$ and $p=0.007$, respectively). As shown in Figure 2, PSS-14 in all of the participants had a moderate significant positive correlation with maladaptive stress-coping strategies ($\rho=0.35$,

S. No	In the last month, how often have you been	Companions (N=56)					Administrative employees (N=98)				
		Never	Almost never	Sometimes	Fairly often	Very often	Never	Almost never	Sometimes	Fairly often	Very often
1	Upset because of something that happened unexpectedly	8 (14.5%)	6 (10.9%)	17 (30.9%)	15 (27.3%)	9 (16.4%)	12 (12.2%)	18 (18.4%)	35 (35.7%)	18 (18.4%)	15 (15.3%)
2	Feeling that you were unable to control the important things in your life	10 (18.2%)	9 (16.4%)	18 (32.7%)	9 (16.4%)	9 (16.4%)	27 (27.6%)	25 (25.5%)	22 (22.4%)	15 (15.3%)	9 (9.2%)
3	Feeling nervous and "stressed"	4 (7.3%)	4 (7.3%)	12 (21.8%)	11 (20.0%)	24 (43.6%)	10 (10.2%)	9 (9.2%)	34 (34.7%)	17 (17.3%)	28 (28.6%)
4	Dealing successfully with day to day problems and annoyances	7 (12.7%)	6 (10.9%)	20 (36.4%)	10 (18.2%)	12 (21.8%)	10 (10.3%)	13 (13.4%)	35 (36.1%)	22 (22.7%)	17 (17.5%)
5	Feeling that you were effectively coping with important changes that were occurring in your life	7 (12.7%)	11 (20.0%)	8 (14.5%)	10 (18.2%)	19 (34.5%)	9 (9.2%)	11 (11.2%)	14 (14.3%)	29 (29.6%)	35 (35.7%)
6	Feeling confident about your ability to handle your personal problems	3 (5.6%)	3 (5.6%)	14 (25.9%)	17 (31.5%)	17 (31.5%)	5 (5.1%)	17 (17.3%)	23 (23.5%)	25 (25.5%)	28 (28.6%)
7	Feeling that things were going your way	11 (19.6%)	6 (10.7%)	20 (35.7%)	11 (19.6%)	8 (14.3%)	8 (8.2%)	20 (20.6%)	30 (30.9%)	25 (25.8%)	14 (14.4%)
8	Finding that you could not cope with all the things that you had to do	12 (21.8%)	10 (18.2%)	20 (36.4%)	7 (12.7%)	6 (10.9%)	30 (30.9%)	20 (20.6%)	28 (28.9%)	15 (15.5%)	4 (4.1%)
9	Able to control irritations in your life	7 (12.7%)	9 (16.4%)	17 (30.9%)	8 (14.5%)	14 (25.5%)	7 (7.2%)	15 (15.5%)	23 (23.7%)	27 (27.8%)	25 (25.8%)
10	Feeling that you were on top of things	4 (7.7%)	10 (19.2%)	16 (30.8%)	11 (21.2%)	11 (21.2%)	8 (8.3%)	15 (15.6%)	40 (41.7%)	20 (20.8%)	13 (13.5%)
11	Angered because of things that happened that were outside of your control	7 (13.0%)	6 (11.1%)	12 (22.2%)	16 (29.6%)	13 (24.1%)	13 (13.5%)	19 (19.8%)	19 (19.8%)	28 (29.2%)	17 (17.7%)
12	Finding yourself thinking about things that you have to accomplish	4 (7.4%)	6 (11.1%)	10 (18.5%)	11 (20.4%)	23 (42.6%)	5 (5.2%)	12 (12.5%)	20 (20.8%)	26 (27.1%)	33 (34.4%)
13	Able to control the way you spend your time	9 (16.1%)	12 (21.4%)	18 (32.1%)	4 (7.1%)	13 (23.2%)	7 (7.2%)	19 (19.6%)	29 (29.9%)	26 (26.8%)	16 (16.5%)
14	Feeling difficulties were piling up so high that you could not overcome them	13 (23.2%)	8 (14.3%)	13 (23.2%)	12 (21.4%)	10 (17.9%)	30 (30.9%)	22 (22.7%)	18 (18.6%)	15 (15.5%)	12 (12.4%)


Table 3: Responses of companions and administrative employees to the items of perceived stress scale (PSS-14).

$p < 0.001$) but a weak non-significant negative correlation with adaptive stress-coping strategies ($\rho = -0.05$, $p = 0.572$). The positive correlation between PSS-14 and maladaptive stress-coping strategies was higher among the companions compared with the administrative employees.

Discussion

Here, we report a slight increase in the level of stress among companions compared to administrative employees in a tertiary care hospital in Saudi Arabia. The slight increase in the level of stress among companions may be explained by the exposure to several caregiving-associated stressors such as worry about the patient, loss of personal

time, work absenteeism, multiple roles, and the hospital environment [12]. On the other hand, the administrative employees in the current study had more work-related stressors, which may have reduced the difference. Comparing the current findings to previous data is complicated by the lack of similar local studies and limited international data. A number of international studies showed higher burden, distress, and psychological disorders among caregivers of admitted patients [9-11]. An Italian longitudinal study of family caregivers of patients with prolonged disorders of consciousness ($n = 18$) showed a progressive and statistically significant increase of "emotional burden" during the hospital stay [22]. In Netherlands, relatives of ICU survivors could



	Companions (N=56)	Administrative employees (N=98)	Total (N=154)	p-value
Adaptive stress-coping	46.89 ± 7.73	45.98 ± 7.10	46.31 ± 7.32	0.458
Active coping (2 and 7)	5.79 ± 1.58	6.07 ± 1.58	5.97 ± 1.58	0.282
Instrumental support (10 and 23)	5.38 ± 1.54	5.60 ± 1.62	5.52 ± 1.59	0.396
Planning (14 and 25)	5.86 ± 1.72	6.06 ± 1.50	5.99 ± 1.58	0.442
Acceptance (20 and 24)	6.53 ± 1.43	6.17 ± 1.41	6.30 ± 1.42	0.141
Emotional support (5 and 15)	5.77 ± 1.57	5.22 ± 1.71	5.42 ± 1.68	0.053
Humor (18 and 28)	5.02 ± 2.05	4.79 ± 1.93	4.87 ± 1.97	0.484
Positive reframing (12 and 17)	6.13 ± 1.58	5.93 ± 1.52	6.00 ± 1.54	0.449
Religion (22 and 27)	6.67 ± 1.44	6.13 ± 1.77	6.33 ± 1.67	0.055
Maladaptive stress-coping	27.73 ± 4.34	25.87 ± 5.03	26.55 ± 4.86	0.021
Behavioral disengagement (6 and 16)	3.95 ± 1.69	3.66 ± 1.51	3.77 ± 1.58	0.286
Denial (3 and 8)	4.05 ± 1.60	3.37 ± 1.39	3.62 ± 1.50	0.006
Self-distraction (1 and 19)	6.18 ± 1.57	5.43 ± 1.67	5.70 ± 1.67	0.007
Self-blame (13 and 26)	5.91 ± 1.73	5.63 ± 1.85	5.73 ± 1.80	0.359
Substance use (4 and 11)	2.02 ± 0.13	2.12 ± 0.71	2.08 ± 0.57	0.276
Venting (9 and 21)	5.63 ± 1.48	5.65 ± 1.51	5.64 ± 1.50	0.911

Table 4: Differences in stress-coping strategies between companions and administrative employees.




Figure 2: Correlation between perceived stress scale (PSS-14) score and stress-coping strategies scores among companions and administrative employees.

experience strain 3 months after hospital discharge and are at risk of developing post-traumatic stress disorder –related features [23]. In a systematic review of psychosocial outcomes in informal caregivers of the critically ill, found that depressive symptoms were the most

prevalent in informal caregivers of survivors of intensive care who were ventilated for more than 48 hours and persist at 1 year with a prevalence of 22.8-29.0%, which is comparable with caregivers of patients with dementia [24]. Nevertheless, there is a major difference in the role of the companions in the current study and the companions/sitters in the Western model. For example, the sitters in Western hospitals are typically non-relative trained personnel who have specific caregiving roles to reduce the need and the cost of nursing care among patients with mainly psycho-geriatric conditions [25,26]. On the other hand, the companions in Saudi hospitals are typically family members/friends who provide mainly psychological support to the admitted patients with almost no caregiving role other than helping the patient with eating and walking [3].

The current study showed an interaction between the gender and the study groups with significantly higher stress among the companions that was observed only in males. This interaction was even more apparent in the adjusted model indicating that it is not influenced by gender-specific demographic, clinical, or work-related differences. Interestingly, there were higher stress levels in females compared with males in both the companions and the administrative employees.

Similar to the current findings, gender-specific differences in the level of stress among caregivers have been reported, with more stress reported among females than males [27-30]. This has been partially explained by gender differences in the care recipient's psychosocial adjustment, with less caregiver's esteem observed among female caregivers [29]. Additionally, hospital administrative jobs have been shown to be less stressful than clinical jobs and have been used as a control group when examining stress among healthcare workers with caregiving (clinical) responsibilities [13]. A recent review, found the risk factors for caregiver burden to include female sex, low educational attainment, residence with the care recipient, higher number of hours spent caregiving, depression, social isolation, financial stress, and lack of choice in being a caregiver [27]. However, there is a lack of data examining gender-specific differences in the stress levels among hospital administrators as well as a lack of data comparing this in relation to companions.

Exposure to caregiving-associated stress without protective stress-coping strategies may lead to psychological morbidity, poorer quality of life, and a sense of hopelessness [9,27,28]. This may be translated into a significant proportion of companions who require psychological support [9]. Compared with the administrative employees, the companions in the current study adopted more maladaptive stress-coping strategies that were moderately correlated with the level of stress. This may explain the slight increase in the level of stress among these companions. Given the low prevalence of stress management training in the current study, the current findings may highlight the need and the importance of developing psychological interventions to reduce and prevent high stress outcomes among companions [10]. Unfortunately, comparing the current findings is limited by a lack of similar studies that used the same stress-coping tool in a controlled study design. However, in a Korean study to identify attitudes about stress and coping among 33 primary caregivers of Hemodialysis HD patients using Q-methodology, three discrete factors emerged as follows: Factor I (they reduced their stress by participating in religious activities; religious sublimation), Factor II (they always worried about the caregiving situations and about the patients' conditions; nervousness), and Factor III (they thought it better to accept their stressful situations; leading handler). These three factors accounted for 44.5% of all the variance, including Factor I (26.0%), Factor II (10.1%), and Factor III (8.4%) [31].

The current study had many advantages: controlled design; using well-validated tools for perceived stress and stress-coping; and examining pathophysiologic changes. Nevertheless, we acknowledged a number of limitations: the possibility of a reporting bias cannot be excluded because the current study was a self-reported study; the case-control design precluded the detection of any causal associations between potential risks and perceiving high stress levels; and, finally, being a single-center study, the findings should be generalized to Saudi companions with caution.

Conclusion

In conclusion, the companions had a slight (marginally significant) increase in the level of stress compared with the administrative employees after adjusting for socio-demographic, clinical, and work-related characteristics. Females had higher stress levels than males in both of the study groups. Maladaptive stress-coping strategies that were moderately correlated to perceived stress were more frequently used by companions. Companions may need psychological support to reduce and prevent high stress outcomes.

Acknowledgment

The authors would like to acknowledge the support they received

from the SABIC Psychological Health Research and Applications Chair, Deanship of Scientific Research, King Saud University, Riyadh, Saudi Arabia. Furthermore, the authors would like to express their gratitude to Dr. Aiman El-Saed (Asst. Professor of Epidemiology and Biostatistics) for his assistance in data analysis and writing and Ms. Ahoud Alluhaim for assistance in translation.

Competing Interests

The authors declare that they have no competing interests.

References

1. Blumenfeld M, Milazzo J, Orlowski B (2000) Constant Observation in the General Hospital. *Psychosomatics* 41: 289-293.
2. Worley LL, Kunkel EJ, Gitlin DF, Menefee LA, Conway G (2000) Constant observation practices in the general hospital setting: a national survey. *Psychosomatics* 41: 301-310.
3. Al-Asmary SM, Al-Shehri A-SA, Al-Omari FK, Farahat FM, Al-Otaibi FS (2010) Pattern of use and impact of patient sitters on the quality of healthcare in Saudi Arabia. *Saudi Med J* 31: 688-694.
4. Al-Mandeeel HM, Almufleh AS, Al-Damri AJ, Al-Bassam DA, Hajr EA, et al. (2013) Saudi womens acceptance and attitudes towards companion support during labor: should we implement an antenatal awareness program? *Ann Saudi Med* 33: 28-33.
5. Carr FM (2013) The role of sitters in delirium: an update. *Can Geriatr J* 16: 22-36.
6. Figved N, Myhr KM, Larsen JP, Aarsland D (2007) Caregiver burden in multiple sclerosis: the impact of neuropsychiatric symptoms. *J Neurol Neurosurg Psychiatry* 78: 1097-1102.
7. Leiknes I, Tysnes OB, Aarsland D, Larsen JP (2010) Caregiver distress associated with neuropsychiatric problems in patients with early Parkinson's disease: the Norwegian ParkWest study. *Acta Neurol Scand* 122: 418-424.
8. Labiano-Fontcuberta A, Mitchell AJ, Moreno-García S, Benito-León J (2015) Anxiety and depressive symptoms in caregivers of multiple sclerosis patients: The role of information processing speed impairment. *J Neurol Sci* 349: 220-225.
9. Luchetti L, Porcu N, Dordoni G, Gobbi G, Lorigo A (2012) [Burden of caregivers of aged patients at a hospital acute care unit and role of the psychologist in the management of the "fragile" caregiver]. *G Ital Med Lav Ergon* 34: A34-A40.
10. Nabors LA, Kichler JC, Brassell A, Thakkar S, Bartz J, et al. (2013) Factors related to caregiver state anxiety and coping with a child's chronic illness. *Fam Syst Health* 31: 171-180.
11. Klarić M, Frančišković T, Pernar M, Nemčić Moro I, Miličević R, et al. (2010) Caregiver Burden and Burnout in Partners of War Veterans with Post-traumatic Stress Disorder. *Coll Antropol* 34: 15-21.
12. Bialon LN, Coke S (2012) A study on caregiver burden: stressors, challenges, and possible solutions. *Am J Hosp Palliat Care* 29: 210-218.
13. Golshiri P, Pourabdian S, Najimi A, Zadeh HM, Hasheminia J (2012) Job stress and its relationship with the level of secretory IgA in saliva: a comparison between nurses working in emergency wards and hospital clerks. *J Pak Med Assoc* 62: S26-S30.
14. Andreou E, Alexopoulos EC, Lionis C, Varvogli L, Gnardellis C, et al. (2011) Perceived Stress Scale: Reliability and Validity Study in Greece. *Int J Environ Res Public Health* 8: 3287-3298.
15. Cohen S, Kamarck T, Mermelstein R (1983) A Global Measure of Perceived Stress. *J Heal Soc Behav* 24: 385-396.
16. Cohen S (1988) Perceived stress in a probability sample of the United States.
17. Carver CS (1997) You want to measure coping but your protocol's too long: consider the brief COPE. *Int J Behav Med* 4: 92-100.
18. Kasi PM, Naqvi HA, Afghan AK, Khawar T, Khan FH, et al. (2012) Coping styles in patients with anxiety and depression. *ISRN Psychiatry* 2012:128672.
19. Ezzati A, Jiang J, Katz MJ, Sliwinski MJ, Zimmerman ME, et al. (2014) Validation of the Perceived Stress Scale in a community sample of older adults. *Int J Geriatr Psychiatry* 29: 645-652.
20. Katsarou A, Panagiotakos D, Zafeiropoulou A, Vryonis M, Skoularigis I, et al. (2012) Validation of a Greek version of PSS-14: a global measure of perceived stress. *Cent Eur J Public Health* 20: 104-109.

21. Montel S, Albertini L, Desnuelle C, Spitz E (2012) The impact of active coping strategies on survival in ALS: The first pilot study. *Amyotrophic Lateral Sclerosis. Amyotroph Lateral Scler* 13: 599-601.
22. Moretta P, Estraneo A, De Lucia L, Cardinale V, Loreto V, et al. (2014) A study of the psychological distress in family caregivers of patients with prolonged disorders of consciousness during in-hospital rehabilitation. *Clin Rehabil* 28: 717-725.
23. van den Born-van Zanten SA, Dongelmans DA, Dettling-Ihnenfeldt D, Vink R, van der Schaaf M (2016) Caregiver strain and posttraumatic stress symptoms of informal caregivers of intensive care unit survivors. *Rehabil Psychol* 61: 173-178.
24. Haines KJ, Denehy L, Skinner EH, Warrillow S, Berney S (2015) Psychosocial outcomes in informal caregivers of the critically ill: a systematic review. *Crit Care Med* 43: 1112-1120.
25. Boswell DJ, Ramsey J, Smith MA, Wagers B (2001) The cost-effectiveness of a patient-sitter program in an acute care hospital: a test of the impact of sitters on the incidence of falls and patient satisfaction. *Qual Manag Health Care* 10: 10-16.
26. Rochefort CM, Ward L, Ritchie JA, Girard N, Tamblyn RM (2012) Patient and nurse staffing characteristics associated with high sitter use costs. *J Adv Nurs* 68: 1758-1767.
27. Adelman RD, Tmanova LL, Delgado D, Dion S, Lachs MS (2014) Caregiver Burden A Clinical Review. *JAMA* 311: 1052-1060.
28. Ingersoll-Dayton B, Raschick M (2004) The Relationship Between Care-Recipient Behaviors and Spousal Caregiving Stress. *Gerontologist* 44: 318-427.
29. Kim Y, Loscalzo MJ, Wellisch DK, Spillers RL (2006) Gender differences in caregiving stress among caregivers of cancer survivors. *Psychooncology* 15: 1086-1092.
30. Schrank B, Ebert-Vogel A, Amering M, Masel EK, Neubauer M, et al. (2015) Gender differences in caregiver burden and its determinants in family members of terminally ill cancer patients. *Psychooncology*.
31. Yeun EJ, Bang HY, Kim EJ, Jeon M, Jang ES, et al. (2016) Attitudes toward stress and coping among primary caregivers of patients undergoing hemodialysis: A Q-methodology study. *Hemodial Int*.