

Oral Hypoglycemic as Attractive Alternative to Insulin for the Management of Diabetes Mellitus during Pregnancy

Magdy A Mohamed^{1*}, Allam M Abdelmonem¹, Mostafa A Abdellah¹ and Adel A Elsayed²

¹Obstetrics & Gynecology Department, Sohag University, Egypt

²Internal Medicine Department, Sohag University, Egypt

Abstract

Objective: To study the efficacy and safety of glibenclamide/metformin combination as a treatment option for pregnant women with either gestational diabetes or type-2 diabetes mellitus compared to conventional insulin therapy.

Methods: Eighty-four pregnant women with singleton pregnancies (65 women with gestational diabetes & 19 with type-2 diabetes mellitus) were included. They were randomly allocated to receive either glibenclamide/metformin combination or insulin. The primary end point was the achievement of the desired level of glycemic control. The presence or absence of maternal, fetal and/or neonatal complications was measured as secondary end points.

Results: There were no differences between the oral hypoglycemic-treated group and insulin-treated group as regard the mean blood glucose levels. It was 125.64 ± 18.15 mg/dL in the former group where it was 124.62 ± 9.29 mg/dL in the latter one. There was no significant difference as regard the amniotic fluid volume, gestational age at delivery and mode of delivery. Also, there were no significant difference between the two groups as regard fetal outcomes; the perinatal mortality was 7.1% in oral hypoglycemic group compared to 4.8% in insulin group, and the rate of congenital anomalies was 4.9% in oral group compared to 9.5% in insulin group. Also there was no statistically significant difference as regard neonatal birth weight, neonatal umbilical blood glucose, incidence and duration of admission to NICU.

Conclusion: The treatment of gestational diabetes and type-2 diabetes mellitus pregnant ladies with Glibenclamide/metformin combination or insulin seems to be equivalent for both the mother and newborns.

Keywords: Diabetes; Metformin; Glibenclamide

Introduction

Very tight glycemic control is the main requirement for improvement of pregnancy outcome in diabetic ladies [1]. Insulin is the only approved pharmacological therapy to control hyperglycemia during pregnancy [2,3]. To achieve tight glycemic control required during pregnancy by the use of insulin, the pregnant lady has to have multiple insulin injections of insulin pump. In developing countries lack of both affordability and education are the main barriers to achieve this level of tight glycemic control by insulin use. The use of oral therapy to control glycemia during pregnancy, if possible, would be an excellent alternative to overcome these barriers.

Problems with the use of oral treatment of DM in pregnancy result from the ability of many antidiabetic oral agents to cross the placenta and cause fetal hyperinsulinemia. The fear of the possible teratogenic effects of oral antidiabetic drugs is another obstacle to their use in pregnancy. In 1991, Elliott et al. demonstrated experimentally that minimal glyburide was detectable crossing the placenta in an in vitro placental perfusion model [4]. In 2000, Langer et al. published the results of their controlled randomized clinical trial; where glyburide was compared with insulin in treatment of diabetes mellitus [5]. Reanalysis of the results of this study revealed that the use of large dose of Glyburide >10 mgm was accompanied with a trend to increased neonatal size [6]. Metformin on the other hand, in spite of its significant ability to cross the placenta, does not cause hyperinsulinemia and was reported to be effective and safe in several clinical studies [7-12]. However, most of the published studies are small and retrospective.

Aim of the Study

To compare Glibenclamide/Metformin combination with insulin in treatment of diabetes mellitus during pregnancy; whether gestational or type-2.

Patient and Methods

Study design

Randomized controlled clinical trial.

This study was conducted on women attending outpatient clinic of obstetrics and gynecology department, Sohag university hospital.

Inclusion criteria

- Gestational Diabetes Mellitus (GDM): diagnosed according to WHO criteria by using Oral Glucose Tolerance Test (OGTT) that consists of glucose load of 75 g and a blood sample was measured 2 hrs later. The diagnosis of GDM was established if plasma glucose level was more than 140 mg/dL [13].

- Type 2 diabetes: where the beginning of the recruitment was before pregnancy.

Exclusion criteria

- Women with preexisting type 1 diabetes mellitus.
- Women with history of diabetic ketoacidosis.
- Multi-fetal pregnancy.

***Corresponding author:** Magdy A Mohamed, Obstetrics & Gynecology Department, Sohag University, Egypt, Tel: +20 93 4601807; E-mail: magdyelkardosy@yahoo.com

Received July 31, 2013; **Accepted** December 28, 2013; **Published** January 07, 2014

Citation: Mohamed MA, Abdelmonem AM, Abdellah MA, Elsayed AA (2014) Oral Hypoglycemic as Attractive Alternative to Insulin for the Management of Diabetes Mellitus during Pregnancy. Gynecol Obstet (Sunnyvale) 4: 193 doi:[10.4172/2161-0932.1000193](https://doi.org/10.4172/2161-0932.1000193)

Copyright: © 2014 Mohamed MA, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

- Hypersensitivity to the used medications.
- Underlying vascular disease or medical condition known to affect fetal growth or drug clearance such as: chronic hypertension, systemic lupus erythematosus, chronic renal insufficiency, hepatic disease, antiphospholipid antibody syndrome or thrombophilia.
- Fetal anomalies identified on ultrasound prior to initiation of therapy.
- Diagnosis of GDM made after 32 weeks gestation.
- Patient refusal.

The study protocol was approved by the faculty scientific research ethics committee; all women who agree to participate in the study, by informed written consent, were randomly allocated into two equal groups. The 1st group received Glibenclamide/Metformin combination; the beginning dose was 2.5 mg Glibenclamide and 500 mg Metformin orally with the morning meal. The dosage was increased gradually according to blood glucose level to a maximum daily dose of 10 mg Glibenclamide and 2 gm Metformin. If this maximum daily dose of the combination did not result in achieving the target values for two week period, insulin was given [5].

The second group received standard insulin therapy. Dosing was based upon subcutaneous two shot combined dose of intermediate acting and short acting insulin given prior to breakfast and dinner. The starting dose was 0.7 unit per kilogram of the body weight at admission and increased weekly as necessary [5].

All women were provided with standard nutritional instructions for three daily meals. Adherence to the dietary regimen was evaluated and reinforced at weekly visits to the clinic. The diets were designed to provide 25 kcal per kilogram of body weight for the obese women and 35 kcal per kilogram for the non-obese ones, with 40 to 45 percent of the calories from carbohydrates [5].

The sample size was determined using the following formula ($n=2 * C_p, \text{ power}/d^2$), where n is the number of subjects required in each group, d is the standardized difference and $C_p, \text{ power}$ is a constant defined by values chosen for the p value and power required (for p value 0.05 & power 90% the constant 10.5).

The standardized difference (d) equal Target difference/standard deviation= $15/21=0.71$. $n=2 * 10.5/(0.71)^2=41.65$. So sample size in each group 42 & total sample size 84.

Method of randomization

Patients were serially ranked. Random tables were used to randomize those patients into two equal groups (using Glibenclamide/metformin was labeled as letter “A”, and the patents who used insulin therapy was labeled as letter “B”. Both “A” and “B” labels were put in the opaque envelopes according to serial ranking obtained from the random tables. After counseling women about the nature of the study and after obtaining a written consent, the corresponding serial envelop was opened and the patient then received the corresponding drug according to the type of label inside “A” or “B”.

The goals of treatment was the achievement of a mean blood glucose concentration of 90 to 105 mg per deciliter (5.0 to 5.9 mmol per liter), a fasting blood glucose concentration of 60 to 90 mg per deciliter (3.4 to 5.0 mmol per liter), a pre-prandial blood glucose concentration of 80 to 95 mg per deciliter (4.5 to 5.3 mmol per liter), and a postprandial blood glucose concentration of less than 120 mg per deciliter (6.7 mmol per liter) and hemoglobin A_{1c} levels of less than 6.0% [1,5].

After establishment of optimum dose in both groups, the patients were followed up for:

- Weekly fasting and after two hour blood glucose level
- Development of polyhydramnios.
- Assessment of fetal wellbeing.

Upon admission for delivery both groups were compared as regard:

- Gestational age at delivery
- Mode of delivery
- Serum glycosylated hemoglobin A_{1c}.
- Birth weight: macrosomia was defined as birth weight 4000 gm or more [5,14].
- Fetal umbilical cord blood glucose level.
- Neonatal hyperbilirubinemia which was defined as a total bilirubin 12 mg/dl or more within the first 7 days of the birth.
- Admission to Neonatal Intensive Care Unit (NICU), and the duration of the admission.
- Neonatal outcome.

The results were statistically analyzed using independent sample student’s t-Test to compare numerical value & chi-square test or Fisher exact test to compare categorical data. P-value <0.05 was considered statistically significant.

Results

Eighty-four patients were included in this study; forty-two patients received oral hypoglycemic in the form of glibenclamide/metformin and the other forty-two patients received insulin therapy as a comparison group. The following table shows that both groups were comparable at the time of recruitment (Table 1).

There was no statistically significant difference between the two groups as regard mean maternal blood glucose, amniotic fluid index, glycosylated HbA_{1c} before delivery and reduction in glycosylated HbA_{1c} (Table 2).

There was no statistically significant difference between two groups

	Oral hypoglycemic group	Insulin group	p-value
Age	33.19 ± 4.94	32.14 ± 5.73	0.189
Number of previous pregnancies	3.72 ± 1.79	3.17 ± 1.98	0.097
Gestational age	22.14 ± 7.33	24.48 ± 6.33	0.063
Glycosylated HbA _{1c}	6.88 ± 0.62	6.96 ± 0.53	0.258

Table 1: Clinical characteristics of patients included in the study at admission (mean ± SD).

	Oral hypoglycemic group	Insulin group	p-value
Mean maternal blood glucose	125.64 ± 18.15	124.62 ± 9.29	0.374
Amniotic fluid index	17.05 ± 2.91	17.67 ± 2.85	0.167
Gestational age at delivery	38.05 ± 1.89	38.26 ± 1.24	0.272
Glycosylated HbA _{1c} before delivery	5.54 ± 0.64	5.43 ± 0.49	0.205
Reduction in glycosylated HbA _{1c}	1.34 ± 0.77	1.53 ± 0.73	0.130

Table 2: Effect of treatment and degree of control of diabetes (mean ± SD).

	Oral hypoglycemic group	Insulin group	p-value
Birth weight	3507.1 ± 462.5	3527.4 ± 489.4	0.424*
Fetal umbilical cord blood glucose	60.49 ± 12.87	59.52 ± 11.33	0.361*
Neonatal hypoglycemia	5 (12.1%)	4 (9.5%)	0.695**
Neonatal jaundice	18 (43.8%)	20 (47.6%)	0.620**
Admission to Neonatal Intensive Care Unit (NICU)	26 (63.4%)	29 (69%)	0.432**
Duration of admission to NICU	3.31 ± 1.77	3.45 ± 2.67	0.412*
Perinatal mortality	3 (7.1%)	2 (4.8%)	0.320***
Congenital malformations	2 (4.9%)	4 (9.5%)	0.249***

(*) using Independent sample T test.

(**) using Chi square test.

(***) using Fisher exact test.

Table 3: Neonatal outcomes in both groups.

as regard fetal birth weight, fetal blood glucose, neonatal jaundice, neonatal hypoglycemia, admission to NICU, duration of admission to NICU, Perinatal mortality and congenital malformations (Table 3).

Discussion

In the current study glibenclamide/metformin combination was comparable to insulin treatment as regard mean maternal blood glucose, amniotic fluid volume, glycosylated HbA1c before delivery and reduction in glycosylated HbA1c. These results are in agreement with Langer et al. who used only glibenclamide in comparison with insulin [5]. Also in agreement with Jacobson et al. and Ramos et al. in their retrospective studies comparing glibenclamide and insulin; as regard maternal blood glucose [14,15]. On the other hand, results of the current study agreeing with studies comparing metformin only with insulin like that of Janet et al. in their prospective study in gestational diabetes (MIG trial) and also with that of Hickman et al. [16,17]. In all these studies only one oral hypoglycemic drug was used compared to two drug combination in our study but the maximum dose in some of these studies of glibenclamide was 20 mg compared to 10 mg in our study.

The birth weight was comparable between the two groups in the current study (3507.1 ± 462.5 gm in oral hypoglycemic group versus 3527.4 ± 489.4 gm in insulin group). This was resembling the results of Jacobson et al. (3661 ± 629, 3559 ± 650 for glibenclamide and insulin respectively) and with Ramos et al. (3420 ± 786, 3524 ± 548 for glibenclamide and insulin respectively) [14,15]. Also our results agree with study comparing metformin and insulin such as Janet et al. (3372 ± 572, 3413 ± 569 for metformin and insulin respectively) and Kristiina et al. (3671 ± 598, 3759 ± 642 for metformin and insulin respectively). Langer et al. found that birth weight was comparable between glibenclamide and insulin treatment but the birth weight in both groups was slightly lower than our results (3256 ± 543, 33194 ± 598 for glibenclamide and insulin respectively) [5,16-18].

The umbilical blood glucose was comparable in both groups (60.49 ± 12.87 mg/dl in oral hypoglycemic group versus 59.52 ± 11.33 mg/dl in insulin group). This agrees with Jean et al. who compared metformin and glibenclamide (57.9 ± 20.3 versus 54.7 ± 15.4 respectively) [19]. Neonatal hypoglycemia occurred in 12.1% in oral hypoglycemic group and was comparable to 9.5% in insulin group. These results agree with Langer et al. whereas neonatal hypoglycemia encountered in 9% in glibenclamide group and 12% in insulin group and agree with Janet et al. (15.2% versus 18.6% with metformin and insulin respectively) [5,16]. However our results disagree with Ramos et al. they reported increased risk of neonatal hypoglycemia in the glibenclamide group

34% compared 14% in insulin group [15]. This can be explained by retrospective nature of this study and there was no maximum dose for glibenclamide. In our study the maximum dose of glibenclamide was 10 mg in addition of using metformin in combination to glibenclamide to decrease the dose of glibenclamide.

Both groups were comparable as regards admission to Neonatal Intensive Care Unit (NICU). About 63% of neonates admitted in oral hypoglycemic group whereas 69% admitted in insulin group. This agrees with Kristiina et al. (42.2% versus 62.2% in metformin & insulin group respectively), Janet et al. (18.7% versus 21.1% in metformin & insulin group respectively) [16-18]. Also duration of admission was similar in our study (3.31 ± 1.77 days versus 3.45 ± 2.67 days in oral hypoglycemic & insulin group respectively). This agrees with Kristiina et al. (2.4 days versus 3.9 days), However our results not agree with & Jacobson et al. who found higher frequency of admission in insulin group & longer duration of admission (15% versus 24%, P=0.008), (4.3 days versus 8.01 days, P=0.008) in glibenclamide & insulin group respectively [14,18]. The explanation of this difference in this retrospective study that insulin group had a higher mean body mass index & a significantly higher mean fasting blood glucose before treatment.

Both groups were comparable as regard congenital malformation & perinatal mortality. About 4.9%, 7.1% respectively in oral hypoglycemic group in comparison to 9.5%, 4.8% in insulin group. These results agree with Janet et al. who reported similar rate of congenital malformations between metformin & insulin group (3.1% versus 4.9% respectively) [16]. Also it agrees with Langer et al. who reported nearly the same rate of congenital malformation (2%) & perinatal mortality (1%) between glibenclamide & insulin groups [5]. The above two studies include only gestational diabetes cases.

Conclusion

The use of the Glibenclamide/metformin combination appears as attractive alternative to traditional insulin treatment in controlling diabetes during pregnancy. It has advantage of being cheaper, more easily to administer & more accepted by patients especially in developing countries. Further studies are needed before wide spread use of oral hypoglycemic for pregnant ladies.

References

1. John Kitzmiller L, Jennifer Block M, Florence Brown M, Patrick Catalano M, Deborah Conway L, et al. (2008) Managing Preexisting Diabetes for Pregnancy. *Diabetes Care* 31: 1060-1079.
2. American Diabetic Association (2004) Gestational diabetes mellitus. *Diabetes care* 1: S88-S90.
3. American College of Obstetricians and Gynecologists Committee on Practice Bulletins--Obstetrics (2001) ACOG Practice Bulletin. Clinical management guidelines for obstetrician-gynecologists. Number 30, September 2001 (replaces Technical Bulletin Number 200, December 1994). *Gestational diabetes. Obstet Gynecol* 98: 525-538.
4. Elliott BD, Langer O, Schenker S, Johnson RF (1991) Insignificant transfer of glyburide occurs across the human placenta. *Am J Obstet Gynecol* 165: 807-812.
5. Langer O, Conway DL, Berkus MD, Xenakis EM, Gonzales O (2000) A comparison of glyburide and insulin in women with gestational diabetes mellitus. *N Engl J Med* 343: 1134-1138.
6. Langer O, Yogev Y, Xenakis EM, Rosenn B (2005) Insulin and glyburide therapy: dosage, severity level of gestational diabetes, and pregnancy outcome. *Am J Obstet Gynecol* 192: 134-139.
7. Elliott BD, Schenker S, Langer O, Johnson R, Prihoda T (1994) Comparative placental transport of oral hypoglycemic agents in humans: a model of human placental drug transfer. *Am J Obstet Gynecol* 171: 653-660.
8. Charles B, Norris R, Xiao X, Hague W (2006) Population pharmacokinetics of metformin in late pregnancy. *Ther Drug Monit* 28: 67-72.

9. Coetzee EJ, Jackson WP (1979) Metformin in management of pregnant insulin-independent diabetics. *Diabetologia* 16: 241-245.
10. Coetzee EJ, Jackson WP (1985) The management of non-insulin-dependent diabetes during pregnancy. *Diabetes Res Clin Pract* 1: 281-287.
11. Hague WM, Davoren PM, Oliver J, Rowan J (2003) Contraindications to use of metformin. Metformin may be useful in gestational diabetes. *BMJ* 326: 762.
12. Hughes RC, Rowan JA (2006) Pregnancy in women with Type 2 diabetes: who takes metformin and what is the outcome? *Diabet Med* 23: 318-322.
13. Forsbach-Sánchez G, Tamez-Peréz HE, Vazquez-Lara J (2005) Diabetes and pregnancy. *Arch Med Res* 36: 291-299.
14. Jacobson GF, Ramos GA, Ching JY, Kirby RS, Ferrara A, et al. (2005) Comparison of glyburide and insulin for the management of gestational diabetes in a large managed care organization. *Am J Obstet Gynecol* 193: 118-124.
15. Ramos GA, Jacobson GF, Kirby RS, Ching JY, Field DR (2007) Comparison of glyburide and insulin for the management of gestational diabetics with markedly elevated oral glucose challenge test and fasting hyperglycemia. *J Perinatol* 27: 262-267.
16. Rowan JA, Hague WM, Gao W, Battin MR, Moore MP, et al. (2008) Metformin versus insulin for the treatment of gestational diabetes. *N Engl J Med* 358: 2003-2015.
17. Hickman MA, McBride R, Boggess KA, Strauss R (2013) Metformin compared with insulin in the treatment of pregnant women with overt diabetes: a randomized controlled trial. *Am J Perinatol* 30: 483-490.
18. Tertti K, Ekblad U, Vahlberg T, Rönnemaa T (2008) Comparison of metformin and insulin in the treatment of gestational diabetes: a retrospective, case-control study. *Rev Diabet Stud* 5: 95-101.
19. Silva JC, Pacheco C, Bizato J, de Souza BV, Ribeiro TE, et al. (2010) Metformin compared with glyburide for the management of gestational diabetes. *Int J Gynaecol Obstet* 111: 37-40.

Citation: Mohamed MA, Abdelmonem AM, Abdellah MA, Elsayed AA (2014) Oral Hypoglycemic as Attractive Alternative to Insulin for the Management of Diabetes Mellitus during Pregnancy. *Gynecol Obstet (Sunnyvale)* 4: 193 doi:[10.4172/2161-0932.1000193](https://doi.org/10.4172/2161-0932.1000193)

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 300 Open Access Journals
- 25,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission/>

