

Oxygen Uptake and Spectrophotometric Measurement of Superoxide, Hydrogen Peroxide and Hydroxyl Radicals

Hideharu Shintani*

Chuo University, School of Science, 1-13-27, Kasuga Bunkyo 112-0003 Tokyo, Japan

Keywords: Oxygen uptake; Spectrophotometric measurement; Superoxide anion radicals; Hydrogen peroxide; Hydroxy radicals

Introduction

The uptake or evolution of oxygen during a reaction can be monitored by use of a Clark-type electrode. The method is particularly useful when turbidity or interfering chromogens preclude the use of spectrophotometry [1].

Protocol

Electrode calibration

1. Add deionized water (2 mL) to the temperature-controller reaction chamber with magnetic stirring, and apply 0.7 V across the electrodes.
2. An output of 0.1 to 10 mV is used with a chart recorder set to a value of 100%.
3. Leave electrode to settle for 5 min.
4. Add a few crystals of sodium dithionate to remove O_2 .
5. This value represents 0% O_2 . The electrode is now calibrated from 0 to 100% oxygen.
6. Apply test samples to chamber after carefully washing each time.

Calculation

The concentration of oxygen can be calculated by use of the information that air-saturated water at 30°C contains 0.460 $\mu\text{mol } O_2$.

Spectrophotometric Measurement of Superoxide Anion Radicals

First applied to neutrophil studies in 1968, the nitroblue tetrazolium (NBT) test provides a simple assay for $\cdot O_2^-$ production. NBT is reduced by O_2^- to a blue formazan product which is measured at 620 nm. Spectrophotometric assay techniques are covered in full [2,3]. Pulse radiolysis techniques are also described [4].

Spectrophotometric Measurement of Hydrogen Peroxide

Any system producing $\cdot O_2^-$ will also produce H_2O_2 by the dismutation reaction. H_2O_2 is an uncharged molecule that can readily enter cells as a water look-alike. Most assays utilize a peroxidase with coupled oxidation of a detector molecule.

H_2O_2 can also be determined by measuring the loss of scopoletin fluorescence (460 nm), an assay covered in detail [5] or on the basis of its reaction with cytochrome-c peroxidase to form a stable complex with absorbance at 419 nm (A_{419}) [1]

Herein we describe a method using phenol red to measure hydrogen peroxide in breath condensates [6].

Protocol

1. Dilute stock hydrogen peroxide (30% v/v) 1:1000 with Chelex resin treated distilled water.
2. Calculate the concentration of H_2O_2 by measuring absorbance at 240 nm using a molar absorption coefficient of 43.6 M/cm.
3. Prepare buffered phenol red-peroxidase reagent containing NaCl (40 mM), potassium phosphate (pH 7.0, 10 mM), phenol red (0.1 g/L), and horseradish peroxidase (HRPO; 8.5 unit/mL), adding phenol red and HRPO to buffer just before use.
4. To 1.0 mL phenol red reagent add 0.5 mL breath condensate. Hydrogen peroxide standards are similarly treated.
5. Mix and stand for 5 min at 25°C.
6. Add NaOH (1 M, 10 μL) to give a pH of 12.5 and a stable purple color.
7. Measure absorbance at 610 nm.

Calculation

Include a blank containing H_2O_2 -free water and calculate values from an H_2O_2 standard curve.

Spectrophotometric Measurement of Hydroxy Radical Damage

The hydroxyl radical is much too reactive to be measured directly by simple spectrophotometric techniques. Evidence characteristic of its damage is, therefore, usually sought by use of a detection molecular for such damage. Here we describe the use of the sugar 2-deoxy-D-ribose [7-9].

Protocol

1. Into clean glass tubes place 5-deoxy-D-ribose (5 mM, 0.2 mL) and sodium phosphate buffer (0.024 M containing 0.15 M NaCl, pH 7.4, 0.2 mL).
 2. Add OH⁻ generating system, i.e.;
- [A] 0.1 mL of ferrous salt (freshly prepared at pH <6.0) or

*Corresponding author: Hideharu Shintani, Chuo University, School of Science, 1-13-27, Kasuga Bunkyo 112-0003 Tokyo, Japan, Tel: +81425922336; E-mail: shintani@mail.hinocatv.ne.jp

Received May 21, 2013; Accepted June 12, 2013; Published June 15, 2013

Citation: Shintani H (2013) Oxygen Uptake and Spectrophotometric Measurement of Superoxide, Hydrogen Peroxide and Hydroxyl Radicals. Pharm Anal Acta S7: 005. doi:10.4172/2153-2435.S7-005

Copyright: © 2013 Shintani H. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

[B] ferric-EDTA + ascorbate or

[C] hypoxanthine + xanthine oxidase + ferric salt.

3. Incubate reactants for 30 min at 37°C.
4. Add thiobarbituric acid (1% w/v, 0.5 mL) and trichloroacetic acid (2.8% W/V, 0.5 mL).
5. Heat tube contents at 100°C for 10 min.
6. Cool and read absorbance at 532 nm or fluorescence at 553 nm after excitation at 532 nm.

Calculation

The assessment of OH \cdot damage can be used to test the effectiveness of scavengers and inhibitors, and to determine second-order rate constants for reaction of OH \cdot with various scavenger.

Comments

OH \cdot generating systems [A] and [B] are not inhibited by SCD, whereas [C] is. All three -OH-generating systems should be inhibited by catalase or other H₂O₂ removing enzymes. If solutions are turbid extract chromogen into butan-1-ol.

References

1. Trudgill PW (1985) Oxygen Consumption. In Cell CRC Handbook of Methods of Methods for Oxygen Radical Research, CRC Press, Boca Raton, USA 329-42.
2. Rice Evans CA, Diplock AT, Symons MCR. (1991) Techniques in Free Radical Research, Elsevier Amsterdam 85-91.
3. Greenwald RA (1995) CRC Handbook of Methods for Oxygen Radical Research. CRC Press, Boca Raton, USA 85-91.
4. PUNCHARD NA, Kelly FJ (1996) Free Radicals: A Practical Approach, IRL-Press, Oxford 11-24.
5. Rice-Evans CA, Diplock AT, Symons MCR (1991) Techniques in Free Radical Research. Elsevier, Amsterdam 91-96.
6. Pick E, Keisari Y (1980) A simple colorimetric method for the measurement of hydrogen peroxide produced by cells in culture. J Immunol Methods 38: 161-170.
7. Gutteridge JM (1981) Thiobarbituric acid-reactivity following iron-dependent free-radical damage to amino acids and carbohydrates. FEBS Lett 128: 343-346.
8. Halliwell B, Gutteridge JM (1981) Formation of thiobarbituric-acid-reactive substance from deoxyribose in the presence of iron salts: the role of superoxide and hydroxyl radicals. FEBS Lett 128: 347-352.
9. Gutteridge JM (1987) Ferrous-salt-promoted damage to deoxyribose and benzoate. The increased effectiveness of hydroxyl-radical scavengers in the presence of EDTA. Biochem J 243: 709-714.

Citation: Shintani H (2013) Oxygen Uptake and Spectrophotometric Measurement of Superoxide, Hydrogen Peroxide and Hydroxyl Radicals. Pharm Anal Acta S7: 005. doi:10.4172/2153-2435.S7-005

This article was originally published in a special issue, **Drug Target and Drug Design** handled by Editor(s). Dr. Paulo Sérgio Monzani, University of North Paraná, Brazil

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 250 Open Access Journals
- 20,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: www.omicsonline.org/submission

