

## Write Brief Detail on Cervical Biopsy and Their Details

Pierre Fermat\*

Department of Health Science and Cancer Centre, Albania

### Abstract

A cervical biopsy is a procedure in which a small sample of tissue is taken from the cervix, the lower part of the uterus that connects to the vagina, for microscopic examination. It is usually done as part of a diagnostic workup when abnormal cervical cells are detected on a Pap test or other screening test.

**Keywords:** Cervical cells; Microscopic; Potential malignancies; Cervical biopsy

### Introduction

The cervix plays a crucial role in reproductive health and is susceptible to various conditions, including precancerous or cancerous changes. A cervical biopsy helps to obtain tissue samples from the cervix for laboratory analysis, enabling healthcare providers to assess the presence of abnormal cells or potential malignancies [1-3].

There are different types of cervical biopsies, including punch biopsy, cone biopsy, and endocervical curettage (ECC). The specific type of biopsy performed depends on the purpose of the procedure and the suspected or observed abnormalities.

During a cervical biopsy, a healthcare provider will typically administer local anaesthesia to numb the cervix, ensuring the procedure is as comfortable as possible. The chosen biopsy technique involves the removal of a small sample of cervical tissue, which is then sent to a laboratory for microscopic examination by a pathologist.

The pathologist analyses the tissue samples to identify any abnormal cell changes, such as precancerous or cancerous cells. The biopsy results play a crucial role in determining the appropriate treatment or further diagnostic procedures required, if necessary [4].

While a cervical biopsy is generally considered a safe procedure, some women may experience mild discomfort, cramping, or light bleeding afterward. Your healthcare provider will provide you with specific instructions on post-biopsy care and any precautions you may need to take.

Regular cervical screenings, including Pap smears and follow-up biopsies if indicated, are crucial for early detection and intervention in cases of cervical abnormalities. These screenings significantly contribute to the prevention and early treatment of cervical cancer, improving women's overall health outcomes [5].

### There are three types of cervical biopsy

**Punch biopsy:** A small tissue sample is taken using a biopsy instrument that punches out a small round piece of tissue from the cervix. This is the most common type of cervical biopsy.

**Cone biopsy:** A cone-shaped wedge of tissue is removed from the cervix using a scalpel or laser. This type of biopsy is usually done when abnormal cells are found in the cervical canal or end cervix.

**End cervical curettage (ECC):** A small spoon-shaped instrument called a curette is used to scrape cells from the cervical canal. This procedure is often done in combination with a punch biopsy.

Cervical biopsies are typically performed in a gynecologist's office

or outpatient clinic, and local anaesthesia is usually used to minimize discomfort. After the biopsy, the tissue sample is sent to a laboratory for analysis by a pathologist, who examines it under a microscope to look for signs of abnormal or cancerous cells.

Depending on the results of the biopsy, further testing or treatment may be recommended, such as repeat Pap tests, colposcopy, or a cone biopsy to remove any abnormal cells. If cervical cancer is diagnosed, treatment options may include surgery, radiation therapy, or chemotherapy, depending on the stage and extent of the cancer. Regular cervical cancer screening and follow-up care are important for early detection and treatment of abnormal cervical cells and cancer [6-8].

### Symptoms of cervical biopsy

After a cervical biopsy, it is common to experience some mild discomfort, spotting, and cramping. These symptoms usually subside within a few days and can be managed with over-the-counter pain relievers and rest. However, if you experience any of the following symptoms, it is important to contact your healthcare provider right away:

Heavy bleeding or bleeding that lasts longer than a week

Severe pain or cramping that is not relieved by over-the-counter pain medications

Foul-smelling discharge or fever, which may indicate an infection

Signs of cervical stenosis, such as difficulty passing urine or having bowel movements

Signs of an allergic reaction, such as hives, swelling, or difficulty breathing

While it is rare, there is also a small risk of more serious complications from a cervical biopsy, such as heavy bleeding, infection, or perforation of the uterus or bowel. Be sure to discuss any concerns or questions you may have about the procedure with your healthcare provider before and after the biopsy, and seek medical attention promptly if you experience

\*Corresponding author: Pierre Fermat, Department of Health Science and Cancer Centre, Albania, E-mail: Pierre7475@gmail.com

**Received:** 03-April-2023, Manuscript No. Ccoa-23-97425; **Editor assigned:** 05-April 2023, Preqç No. PQ- Ccoa-23-97425; **Reviewed:** 19-April-2023, QC No Ccoa-23-97425; **Revised:** 22-April-2023, Manuscript No Ccoa-23-97425 (R); **Published:** 29-April-2023, DOI: 10.4172/2475-3173.1000154

**Citation:** Fermat P (2023) Write Brief Detail on Cervical Biopsy and Their Details. Cervical Cancer, 8: 154.

**Copyright:** © 2023 Fermat P. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

any unusual symptoms or complications [9].

### Causes for Cervical biopsy

A cervical biopsy is a medical procedure that involves the removal of a small sample of tissue from the cervix, which is the lower part of the uterus that connects to the vagina. The most common causes of cervical biopsies are abnormal cervical cell changes that have been detected through a Pap test or other screening test. These cell changes may be caused by an infection with the human papillomavirus (HPV), a sexually transmitted infection that can lead to the development of cervical cancer [10].

### Other causes of abnormal cervical cell changes and the need for a cervical biopsy may include

**Cervical dysplasia:** This is a precancerous condition in which the cells of the cervix become abnormal but have not yet developed into cancer.

**Cervical cancer:** This is a type of cancer that develops in the cells of the cervix and can spread to other parts of the body if left untreated.

**Inflammation or infection of the cervix:** This can be caused by a variety of factors, such as a bacterial or fungal infection or a sexually transmitted infection.

**Hormonal changes:** Fluctuations in hormone levels, such as those that occur during pregnancy or menopause, can sometimes cause abnormal cervical cell changes.

**Other factors:** Certain lifestyle factors, such as smoking or a weakened immune system, may also increase the risk of abnormal cervical cell changes and the need for a cervical biopsy.

While a cervical biopsy is typically a safe and low-risk procedure, it is important to discuss any concerns or questions you may have with your healthcare provider before the procedure. Regular cervical cancer screening and follow-up care are also important for early detection and prevention of cervical cancer.

### Prevention for cervical biopsy

Cervical biopsies are typically safe and relatively low-risk procedures. However, as with any medical procedure, there are some potential risks and complications, such as bleeding, infection, or cervical stenosis (narrowing of the cervical opening).

### Here are some tips to help prevent complications and ensure a safe and comfortable cervical biopsy

Avoid sexual activity, vaginal douching, and tampon use for at least 24 hours before the procedure.

Let your healthcare provider know if you are pregnant or have a bleeding disorder, as this may affect the procedure or require special precautions [10].

Discuss any medications you are taking with your healthcare provider, as some medications may increase the risk of bleeding or other complications.

Be sure to inform your healthcare provider if you have a history of abnormal cervical cells, cervical cancer, or other gynaecological conditions.

Ask your healthcare provider about any pre-procedure instructions or preparations, such as fasting or medication adjustments.

Consider taking a mild pain reliever, such as ibuprofen, before the procedure to help reduce discomfort.

Arrange for someone to accompany you to the appointment and drive you home, as you may experience some mild cramping or dizziness after the procedure.

Follow any post-procedure instructions carefully, such as avoiding strenuous activity or sexual activity for a specified period of time, and report any unusual symptoms or complications to your healthcare provider.

Regular cervical cancer screening, including Pap tests and HPV testing, is also important for early detection and prevention of cervical cancer. Consult with your healthcare provider to determine the best screening schedule and approach for your individual needs and risk factors.

### Conclusion

A cervical biopsy is a medical procedure that involves the removal of a small sample of tissue from the cervix for examination under a microscope. It is usually performed as part of a diagnostic process to investigate abnormal cervical cell changes or detect the presence of cervical cancer. The most common cause of cervical biopsies is abnormal cervical cell changes that have been detected through a Pap test or other screening test, often as a result of infection with the human papillomavirus (HPV).

While a cervical biopsy is generally a safe and low-risk procedure, there is a small risk of complications, such as bleeding, infection, or cervical stenosis. It is important to follow any pre-procedure and post-procedure instructions carefully and report any unusual symptoms or complications to your healthcare provider right away.

Regular cervical cancer screening, including Pap tests and HPV testing, is also important for early detection and prevention of cervical cancer. Consult with your healthcare provider to determine the best screening schedule and approach for your individual needs and risk factors.

### References

1. Trevor EA, Erik KA (2019) Thyroid Nodules and Thyroid Cancer in the Pregnant Woman. *Endocrinol Metab Clin North Am* 48: 557-567.
2. Anthony C, Juan R (2020) Radioactive Iodine Therapy in Differentiated Thyroid Cancer: 2020 Update. *AJR Am J Roentgenol* 215: 285-291.
3. Sylvia LA (2019) The Current Histologic Classification of Thyroid Cancer. *Endocrinol Metab Clin North Am* 48: 1-22.
4. Mary KD (2020) Thyroid Cancer: Implications of Genomics for Care and Practice. *Clin J Oncol Nurs* 24: 483-487.
5. Bryan RH (2017) 2015 American Thyroid Association Management Guidelines for Adult Patients with Thyroid Nodules and Differentiated Thyroid Cancer: What is new and what has changed?. *Cancer* 123: 372-381.
6. Antonio M, Federica F, Ferruccio S, Rossella E (2020) Obesity as a risk factor for thyroid cancer. *Curr Opin Endocrinol Diabetes Obes* 27: 358-363.
7. Daniela VPA, Archie B, Juan PB (2017) Thyroid cancer in adolescents and young adults. *Future Oncol* 13: 1253-1261.
8. Daniel V, Mark Z (2021) Surgical Considerations in Thyroid Cancer: What the Radiologist Needs to Know. *Neuroimaging Clin N Am* 31: 327-335.
9. Mahsa S, Electron K (2019) Long noncoding RNAs in thyroid cancer. *Curr Opin Endocrinol Diabetes Obes* 26: 275-281.
10. Noor J, Volkert BW (2018) Aggressive differentiated thyroid cancer. *Eur J Surg Oncol* 44: 367-377.