



An Ecological Risk Assessment of Water Quality's Nutrient Over-Enrichment

Jyoti Singha Pandey*

Department of Marine Sciences, School of life sciences, India

Abstract

The lives of organisms in shallow marine environments continue to be threatened by the natural (seasonality) and anthropogenic (e.g., coastal developments and changes in land-use practices) drivers that compromise water quality. In three Kenyan coral reef ecosystems—the Tana Delta, Watamu, and Shimoni Mupunguti Reef—water and sediment samples from healthy, living *Porites* spp. colonies were used in a case study to evaluate the effects of these drivers on nutrient over-enrichments. For (N-(NO₃-+ NO₂-)) and (P-PO₄-), the assessed nutrient over-enrichment levels were higher than the universal reference background concentrations of 0.01 and 0.003 mg/L, respectively.

Keywords: Ecosystem; Water quality; Coastal cities

Introduction

It is commonly acknowledged that nutrient over-enrichment caused by both natural and human-induced drivers is endangering the water quality of the shallow marine environment. Anthropogenic activities, such as maritime operations, the growing tourism sector, and altered land-use practices, are currently putting pressure on the tropical and sub-tropical coastal environments. The growing number of people living in coastal cities as a result of maritime activity and the growing tourism sector has increased the amount of urban domestic sewage that overburdens coral habitats. It has been observed globally that, in addition to naturally occurring drivers like erosion, runoff, and riverine discharge, other drivers like coastal developments and modifications to land-use practices account for the majority of nutrient emissions [1,2].

Methodology

It is noteworthy that, according to, nutrient over-enrichment has continued to have a negative effect on benthic communities, such as coral reefs, on a global scale. This could have long-term socioeconomic repercussions. It has already been observed that the rate of pollution in the marine environment is concerning. Thirty percent of the global nutrient over-enrichment in coral reef habitats by 2006 came from natural and human sources. The fact that there are more and more "coastal dead zones" raises serious concerns because they are thought to be caused by an unchecked influx of nutrients brought on by human activity.

A concentration of 0.03-0.1 mg/L of N or P is sufficient to establish macro-algae bloom dominancy, according to. Because macro-algae blooms are linked to a structural community shift in benthic communities—that is, from a coral reef community to macro-algae dominance—they can be harmful in the absence of grazers and herbivores. Concerns regarding the effect of human activity on marine pollution have been highlighted in earlier studies conducted by a number of marine scientists reported nutrient over-enrichment from Peri-Urban Creeks and estuaries above the universal reference background concentrations of 0.003 and 0.01 mg/L of P and N, respectively [3].

It should be understood that, especially during high tides, ephemeral environments like creeks and estuaries are constantly in communication with shallow marine environments. Strong tidal waves of pelagic origin in nature resuspension nutrient-rich sediment from these environments during high tides. Through desorption

processes, the water column of the shallow coral reef environment becomes nutrient enriched with the help of the intricate and dynamic hydrological circulation of the marine environment. Consequently, it is important and merits attention to regularly assess nutrient over-enrichment in the shallow environment, as suggested by a number of marine researchers. It is possible to address marine pollution through routine assessment. Moreover, the information derived from routine evaluations can serve as a guide for the development of policies aimed at conserving and protecting marine life [4-6].

Even though routinely assessing nutrient over-enrichment is a crucial effort to combat marine pollution, recent developments in pollution assessment have shown that quantifying nutrients alone—without taking sediment quality into account as one of the water quality parameters, for example—is a less accurate method of assessing nutrient over-enrichment. According to Shen et al. (2021), an evaluation of pollution that disregards the ecological pollution index is considered inadequate, less illuminating, and thus untrustworthy. Therefore, it is believed that the best method is to use sediment quality as a criterion tool for aquatic water quality assessment. This method's strength is its capacity to assess and identify the degree of the risk associated with nitrogen exposure [7,8].

Results

In order to assess the effects of anthropogenically induced and natural drivers on the ecological risk exposure of total dissolved inorganic nitrogen (TIN) and phosphorous (TIP) on coral reef ecosystems, the current study attempts to take sediment quality into account. By comparing the levels of nitrogen and phosphorous in sediments and water columns across the coral reef ecosystems under investigation, the current study aims to accomplish the following two goals: (1) to identify nutrient over-enrichment; and (2) to assess the susceptibility of nutrient over-enrichment in Kenyan coral reef ecosystems using nitrogen and

*Corresponding author: Jyoti Singha Pandey, Department of Marine Sciences, School of life sciences, India, E-mail: jyoti33@yahoo.com

Received: 03-Nov-2023, Manuscript No: jee-23-121345; **Editor assigned:** 06-Nov-2023, Pre-QC No: jee-23-121345 (PQ); **Reviewed:** 20-Nov-2023, QC No: jee-23-121345; **Revised:** 22-Nov-2023, Manuscript No: jee-23-121345 (R); **Published:** 29-Nov-2023, DOI: 10.4172/2157-7625.1000468

Citation: Pandey JS (2023) An Ecological Risk Assessment of Water Quality's Nutrient Over-Enrichment. J Ecosys Ecograph, 13: 468.

Copyright: © 2023 Pandey JS. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

phosphorous pollution index. By addressing marine pollution, this study is in line with the 14th Sustainable Development Goal (SDG), "Life below water [9,10].

Discussion

When assessing pollution, it is crucial to use reference background concentration, particularly when attempting to pinpoint pollution hotspots. Table 6 presents a comparison between the current findings and comparable published research concerning the background concentrations for coastal waters. The critical nutrients N-(NO₃+NO₂-) and P-PO₄³⁻, whose mean concentrations were measured in the water column, fell between the reported concentration ranges from Kenyan estuaries.

Conclusion

There were nutrient over-enrichments above the universal background reference concentrations in both the shallow marine settings (coral reef ecosystems) and the transitional environments (creeks and estuaries). The N/P ratios in both seasons were less than the Renfield ratios of (N/P = 16), suggesting that there was an excess of phosphorous in the water columns of all the reefs under investigation. The nutritional enrichment temporal trend showed that both naturally occurring and man-made drivers.

References

1. Breman JG, Henderson DA (2002) Diagnosis and management of smallpox. *N Engl J Med* 346: 1300-1308.
2. Damon IK (2011) Status of human monkeypox: clinical disease, epidemiology and research. *Vaccine* 29: D54-D59.
3. Ladnyj ID, Ziegler P, Kima E (2017) A human infection caused by monkeypox virus in Basankusu Territory, Democratic Republic of the Congo. *Bull World Health Organ* 46: 593.
4. Olson VA, Laue T, Laker MT, Babkin IV, Drosten C, et al. (2019) Real-time PCR system for detection of orthopoxviruses and simultaneous identification of smallpox virus. *J Clin Microbiol* 42: 1940-1946.
5. MacNeil A, Reynolds MG, Braden Z, Carroll DS, Bostik V, et al (2009) Transmission of atypical varicella-zoster virus infections involving palm and sole manifestations in an area with monkeypox endemicity. *Clin Infect Dis* 48: 6-8.
6. Di Giulio DB, Eckburg PB (2004) Human monkeypox: an emerging zoonosis. *Lancet Infect Dis* 4: 15-25.
7. Ježek Z, Szczeniowski M, Paluku KM, Moomba M (2000) Human monkeypox: clinical features of 282 patients. *J Infect Dis* 156: 293-298.
8. Kulesh DA, Loveless BM, Norwood D, Garrison J, Whitehouse CA, et al. (2004) Monkeypox virus detection in rodents using real-time 3'-minor groove binder TaqMan assays on the Roche LightCycler. *Lab Invest* 84: 1200-1208.
9. Breman JG, Steniowski MV, Zanotto E, Gromyko AI, Arita I (1980) Human monkeypox, 1970-79. *Bull World Health Organ* 58: 165.
10. Karem KL, Reynolds M, Braden Z, Lou G, Bernard N, et al. (2005) Characterization of acute-phase humoral immunity to monkeypox: use of immunoglobulin M enzyme-linked immunosorbent assay for detection of monkeypox infection during the 2003 North American outbreak. *Clin Diagn Lab Immunol* 12: 867-872.