

Understanding Bipolar Manic Depression Symptoms, Causes and Treatment

Vivian James*

Department of Mental Health, Jimma University, Ethiopia

Abstract

Bipolar disorder, formerly known as manic depression, is a mental health condition characterized by extreme mood swings that include emotional highs (mania or hypomania) and lows (depression). These mood swings can be intense and unpredictable, significantly impacting a person's daily life, relationships, and ability to function. In this article, we delve into the intricacies of bipolar manic depression, exploring its symptoms, causes, and available treatments.

Keywords: Bipolar disorder; Mania; Depression; Mood swings; Treatment

Introduction

Bipolar disorder manifests in different ways, with individuals experiencing varying degrees of mood swings. The two primary mood episodes associated with bipolar disorder are manic and depressive episodes. During a manic episode, individuals may feel unusually energetic, euphoric, or irritable. They may exhibit erratic behaviour, such as engaging in risky activities, talking rapidly, and experiencing racing thoughts. Additionally, heightened creativity and a decreased need for sleep are common during manic episodes [1].

Methodology

Conversely, depressive episodes are characterized by overwhelming sadness, loss of interest or pleasure in activities once enjoyed, fatigue, changes in appetite or weight, feelings of worthlessness or guilt, and thoughts of death or suicide [2]. Depressive episodes can severely impair functioning and may last for weeks or even months.

Causes

The exact cause of bipolar disorder remains unclear, but it is believed to result from a combination of genetic, biological, and environmental factors. Family history plays a significant role, with individuals having a close relative with bipolar disorder at a higher risk of developing the condition themselves [3]. Biological factors, such as chemical imbalances in the brain involving neurotransmitters like dopamine and serotonin, may also contribute to the onset of bipolar disorder. Additionally, stressful life events, trauma, substance abuse, and certain medications can trigger or exacerbate mood episodes in susceptible individuals.

Treatment

While bipolar disorder is a chronic condition, effective treatment can help manage symptoms and improve quality of life. Treatment typically involves a combination of medication, psychotherapy, and lifestyle changes. Medications commonly prescribed for bipolar disorder include mood stabilizers, antipsychotics, and antidepressants [4]. Mood stabilizers, such as lithium and valproate, help control manic and depressive episodes, while antipsychotic medications can alleviate symptoms of mania and psychosis. Antidepressants may be prescribed cautiously, as they can trigger manic episodes in some individuals [5]. Psychotherapy, such as cognitive-behavioral therapy (CBT) and interpersonal therapy (IPT), can help individuals better

understand their condition, identify triggers, and develop coping strategies to manage mood swings effectively. Family therapy may also be beneficial in improving communication and support networks [6]. In addition to medication and therapy, adopting a healthy lifestyle can play a crucial role in managing bipolar disorder. Regular exercise, adequate sleep, stress reduction techniques (such as mindfulness and relaxation exercises), and avoiding alcohol and recreational drugs can help stabilize mood and reduce the frequency and severity of mood episodes [7]. Bipolar manic depression is a complex mental health condition characterized by extreme mood swings that can significantly impact an individual's life. While there is no cure for bipolar disorder, effective treatment can help individuals manage symptoms and lead fulfilling lives [8]. With a combination of medication, psychotherapy, and lifestyle changes, individuals with bipolar disorder can achieve stability and maintain overall well-being. It's essential for those affected by bipolar disorder to seek professional help and support from loved ones to effectively manage their condition and navigate the challenges it presents. Bipolar depression is a multifaceted mental health condition that presents significant challenges to individuals and their loved ones [9]. Characterized by extreme mood swings between manic and depressive episodes, bipolar disorder can disrupt daily life, relationships, and overall well-being. However, with proper diagnosis and treatment, individuals with bipolar disorder can effectively manage their symptoms and lead fulfilling lives. Treatment for bipolar depression typically involves a combination of medication, psychotherapy, and lifestyle adjustments. Mood stabilizers, antipsychotics, and antidepressants are commonly prescribed to help control mood swings and alleviate symptoms. Psychotherapy, such as cognitive-behavioral therapy (CBT) and interpersonal therapy (IPT), provides individuals with valuable tools to understand their condition, identify triggers, and develop coping strategies [10]. Additionally, adopting a healthy lifestyle that includes regular exercise, adequate sleep, stress reduction techniques,

*Corresponding author: Vivian James, Department of Mental Health, Jimma University, Ethiopia, E-mail: vivianj395@yahoo.com

Received: 01-Jun-2024, Manuscript No: ppo-24-139188, **Editor assigned:** 03-Jun-2024, PreQC No: ppo-24-139188 (PQ), **Reviewed:** 18-Jun-2024, QC No: ppo-24-139188, **Revised:** 25-Jun-2024, Manuscript No: ppo-24-139188 (R), **Published:** 31-Jun-2024, DOI: 10.4172/ppo.1000207

Citation: Vivian J (2024) Understanding Bipolar Manic Depression Symptoms, Causes and Treatment. Psychol Psychiatry 8: 207.

Copyright: © 2024 Vivian J. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

and avoiding substance abuse can complement medical treatment and contribute to mood stability. It's crucial for individuals with bipolar depression to seek professional help and support from loved ones. With the guidance of mental health professionals and a strong support network, individuals can navigate the challenges of bipolar disorder and learn to effectively manage their symptoms. Education about the condition, open communication with healthcare providers, and adherence to treatment plans are essential components of long-term management.

Conclusion

While living with bipolar depression may present ongoing challenges, it's important to recognize that recovery is possible. By taking proactive steps to manage their condition and prioritize self-care, individuals with bipolar disorder can achieve stability and pursue their goals and aspirations. With understanding, empathy, and appropriate support, individuals with bipolar depression can lead meaningful and fulfilling lives, emphasizing the importance of compassion and destigmatization within society.

Acknowledgement

None

Conflict of Interest

None

References

1. Renaud J, Kenneth H, Matthew H, Jerome B, Flavie W, et al. (2016) Are

Hallucinations Due to an Imbalance Between Excitatory and Inhibitory Influences on the Brain?. *Schizophr Bull* 42: 1124-1134.

2. Marcella M, Pantelis L, Charles F, Flavie W, Frank L, et al. (2021) A Review of Multimodal Hallucinations: Categorization, Assessment, Theoretical Perspectives, and Clinical Recommendations. *Schizophr Bull* 47: 237-248.
3. Kenneth H (2017) Auditory Hallucinations as Translational Psychiatry: Evidence from Magnetic Resonance Imaging. *Balkan Med J* 34: 504-513.
4. Gras A, Amad A, Thomas P, Jardri R (2014) [Hallucinations and borderline personality disorder: a review]. *Encephale* 40: 431-438.
5. Franco DM, Cesare D, Gabriella G, Andrea P (2015) Hallucinations in the psychotic state: Psychoanalysis and the neurosciences compared. *Int J Psychoanal* 96: 293-318.
6. Shusaku O, Hidetomo M, Tomotaka S, Keiko B, Tadashi U, et al. (2021) Risk factors for minor hallucinations in Parkinson's disease. *Acta Neurol Scand* 143: 538-544.
7. Kirk R, John F, Frances S (2011) Auditory hallucinations: a review of assessment tools. *Clin Psychol Psychother* 18: 524-534.
8. Akgor U, Erdem F, Burcu S, Canan U, Murat C, et al. (2021) Anxiety, depression and concerns of pregnant women during the COVID-19 pandemic. *Arch Gynecol Obstet* 304: 125-130.
9. Areej EJ, Jamie MJ, Ashley MN, Lara T, Joseph AG, et al. (2020) Multimodal psychosocial intervention for family caregivers of patients undergoing hematopoietic stem cell transplantation: A randomized clinical trial. *Cancer* 126: 1758-1765
10. Loretta H, Alexandra W, Steven H, Sarah AO, Mark AL, et al. (2021) Clinical psychology PhD students' admission experiences: Implications for recruiting racial/ethnic minority and LGBTQ students. *J Clin Psychol* 77: 105-120.