

Heavy Metals Contamination in Vegetables and its Growing Soil

Tasrina RC^{1*}, Rowshon A², Mustafizur AMR², Rafiqul I³ and MP Ali^{4*}

¹Chemistry Division, Atomic Energy Center, Dhaka-1000, Bangladesh

²Department of Applied Chemistry and Chemical Engineering, Dhaka University, Bangladesh

³Department of Zoology, Rangpur Government College, Rangpur, Bangladesh

⁴Laboratory of Biotechnology, Graduate School of Science and Technology, Shizuoka University, Japan

Abstract

Dietary exposure to several heavy metals including Ni, Cd, Cr, Co, Pb, As, Hg, Zn and Cu, has been recognized as a risk to human health through the consumption of vegetable crops. This study investigates the source and magnitude of heavy metal contamination in soil and various kinds of vegetables including potato, red amarantha, spinach, amarantha, carrot, cabbage, tomato and brinjal at Pakshi, Bangladesh is a mix of commercial and residential vegetables growing areas. The concentration of As and Fe in all soil sample tested is higher than that of permissible limit of different international standard. The concentration of Co in Chor Ruppur, Pb in Pakshi and Diarpur, Mn in Chor Ruppur and Zn in Chor Ruppur and Pakshi soil are also higher than that of standard level. Other tested metals are lower than that of standard level. The lead in vegetables in all tested areas was higher level than that of the permissible limits of different International standards. Except lead, it is concluded that nearly all the samples did not exceed the Intentional Food Standards maximum level (ML) in vegetables at Pakshi region in Bangladesh. If contaminated soil and vegetables contribute to a progressive gathering of these metals in food chain there might possible to deep alternations of ecosystem with possible noxious effect on human health. Our study highlights that vegetables growing soil containing higher amount of metals that could be transferred into edible parts of the plant, so study area should be monitored regularly to avoid health risk of human being due to exposure of toxic level.

Keywords: Heavy metals, Spectroscopy; Soil; Vegetables; Contamination.

Introduction

Heavy metals are hazardous contaminants in food and the environment and they are non-biodegradable having long biological half-lives [1]. The implications associated with metal (embracing metalloids) contamination are of great concern, particularly in agricultural production systems [2] due to their increasing trends in human foods and environment. Metals most often found as contaminants in vegetables include As, Cd and Pb. These metals can pose as a significant health risk to humans, particularly in elevated concentrations above the very low body requirements [3]. So, the metals must be controlled in food sources in order to assure public health safety [4]. Excessive amount of heavy metals in food cause a number of diseases, especially cardiovascular, renal, neurological, and bone diseases [5]. These metals could reach food chain through various biochemical process and ultimately biomagnified in various trophic levels and eventually threaten the health of human. The contamination of soil and vegetables by heavy metals is also a global environmental issue. They are ubiquitous in the environment through various pathways, due to natural and anthropogenic activities [6]. Under certain environmental conditions metals may accumulate to toxic concentration and they cause ecological damages [7,8].

Source of anthropogenic contamination include the addition of manures, sewage sludge, fertilizers and pesticides to soils, several studies identifying the risks in relation to increased soil metal concentration and consequent plant uptake [9,10]. Both commercial and residential growing areas are also vulnerable to atmospheric pollution, in the form of metal containing aerosols. These aerosols can penetrate the soil and be absorbed by vegetables, or alternatively be deposited on leaves and adsorbed. Analysis of vegetables grown in locations close to industry has reported elevated levels of heavy metals contamination [2,11] studied the impact of atmospheric pollution from industry on heavy metal contamination in vegetables grown in Greece. The results of the study indicated significantly higher levels of metal accumulation in leafy vegetables as compared with root vegetables. This partitioning of Cd is well known, with accumulation of greater concentrations in the edible leafy portions of crops, than the storage organs or fruit [12,13].

As the present study area is free from industrial pollution, the major sources of soil contamination with heavy metals might be due to the waste water irrigation, solid waste disposal, sludge applications, vehicular exhaust and agrochemicals. Excessive accumulation of heavy metals in agricultural soils through the use of agrochemicals and by other sources may not only result in soil contamination but also lead to elevated heavy metal up-take by vegetables and thus affect food quality and safety [14]. Heavy metals are easily accumulated in the edible parts of leafy vegetables, as compared to grain or fruit crops [15]. Vegetables take up heavy metals and accumulate them in their edible [16] and inedible parts in quantities high enough to cause clinical problems both to animals and human beings when they consume these metal-rich plants [17]. Intake of toxic metals in a chronic level through soil and vegetables has adverse impacts on human, plants and the associated harmful impacts become apparent only after several years of exposure [16,18]. However, the consumption of heavy metal-contaminated food can seriously deplete some essential nutrients in the body that are further responsible for decreasing immunological defenses, such as intrauterine growth retardation, impaired psycho-social facilities, disabilities associated with malnutrition and high prevalence of upper gastrointestinal cancer rates [19,20].

In this study we investigated the concentrations of As, Pb, Cd, Ni, Mn, Co, Cu, Fe, Zn, and Hg in both soil and vegetable crops within vegetable growing regions of Pakshi, Pabna, Bangladesh; and evaluated their contamination status with respect to EU and international food standard guidelines.

***Corresponding authors:** MP Ali, Laboratory of Biotechnology, Graduate School of Science and Technology, Shizuoka University, Japan, Tel: 08036504222; E-mail: panna_ali@yahoo.com

Tasrina RC, Chemistry Division, Atomic Energy Center, Dhaka-1000, Bangladesh, Tel: 08036504222; E-mail: tsarina_rabia@yahoo.com

Received May 13, 2015; Accepted June 19, 2015; Published June 30, 2015

Citation: Tasrina RC, Rowshon A, Mustafizur AMR, Rafiqul I, MP Ali (2015) Heavy Metals Contamination in Vegetables and its Growing Soil. J Environ Anal Chem 2: 142. doi:10.4172/2380-2391.1000142

Copyright: © 2015 Tasrina RC et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Materials and Methods

Sampling site

Soil and plant samples were collected from 6 sites within four regions across Pakshi, Bangladesh (Figure 1). The soil samples were collected from six sites within four regions (Pakshi, Dear Baghail, Rooppur Dear and Chor Rooppur) and the vegetable samples from the Char Rooppur field (the land where the nuclear power plant will be set up) of Pakshi union, Ishwardi, Pabna. The following vegetables are considered for heavy metals investigation: potato, red amarantha, spinach, amarantha, carrot, cabbage, tomato and brinjal Bangladesh (Figure 2).

The sites were a mixture of commercial vegetables farms and private residential vegetable gardens. Currently, Bangladesh Government has decided to establish a nuclear power plant for electricity generation in Pakshi, Ishwardi, Pabna. Therefore a monitoring of the environmental components like vegetables and soil at Pakshi area was done for the determination and measurement of heavy metals as background study for safe and secured introduction of nuclear power plant in Bangladesh and also these results will provide important information for health purpose.

Soil and vegetable sampling



Figure 1: The location of sampling region.




Figure 2: Comparison of heavy metals content in vegetables and its growing soil (Chor Rooppur soil).

Soil samples were collected from one side of the Padma River within 0-7 km and locations were maintained at about 1 km distance from one sampling point to another. The samples were about 500 g in weight. Immediately after collection, the samples were placed in the polythene bag which were washed with deionized water repeatedly for 3 times

and taken to the laboratory as soon as possible. In the laboratory, the collected soil samples were dried in electrical oven at a temperature around 90°C to remove the moisture and then homogenized in mortar-pestle and reduced size to a fine powder. On the other hand, the collected vegetable samples (about 500 g) were thoroughly washed with fresh water in order to remove the adhering dirt and finally with deionized water. Then approximately 10 g of each sample were taken to a mortar for grinding to obtain a homogenous mass. Finally the samples were taken to the small airtight polythene bags and then kept to the refrigerator for further analysis.

Chemicals and equipment used

Individual standard solutions (Spectro Pure, USA) of target elements were supplied by Varian Inc, USA with highest purity level (99.98%), supra pure nitric acid, potassium iodide and hydrochloric acid were purchased from E. Merck, Germany; sodium borohydride (Acros Organics, USA). All other chemicals were extra pure or supra pure received from E. Merck, Germany. The analytical

instruments used in this study were Flame Atomic Absorption Spectrometer (FAAS), Varian Analytical Instrument, model Varian AA 280Z (Zeeman AAS) for Ni, Cr, Cd, Co, Cu, Pb, Fe, Zn and Mn. On the other hand, Cold Vapor Atomic Absorption Spectrometer (CVAAS), model ANALYTIK JENA (novAA350) for Hg and Hydride Generation Atomic Absorption Spectrometer (HGAAS), model AA 240 for arsenic determination. A microwave accelerator reaction system (Model No; MARS 5) was used for the digestion of the samples.

Digestion of soil and vegetable samples for metal analysis

To determine the concentration of heavy metals in soil and vegetable samples, aliquot (about 0.3 g) amount of the ground dried samples were taken in a vessel (Model no XP-1500) of the microwave oven where 4 ml of concentrated HNO_3 was poured. The heating program employed was the one proposed in the user's manual. The three steps temperature programs were applied - 180°C with a ramping time of 10 min; holding time 15 min and cooling time 10 min. After completion of digestion, it is necessary to cool down rotor at a temperature of 60°C and then after cooling, loosen the upper screw of the vessel carefully with a torque wrench to release pressure under fume hood. The samples thus obtained were filtered and leveled up to the mark with deionized water in a 10 ml volumetric flask. Finally the samples were examined with AAS for heavy metals estimation.

Experimental procedure for metals determination

The working standard solutions were prepared for heavy metal determination except arsenic and mercury by diluting a stock solution containing 1000 ppm of single element Atomic Absorption Spectrometer (AAS) grade standard with ultra-pure water. Cold Vapor Atomic Absorption Spectrometer (CVAAS) technique was used to determine the concentration of mercury (Hg) in the samples. 4 ml of the digested sample from the mother stock was taken in a test tube; 1 ml concentrated hydrochloric acid (HCl) and 15 ml de-ionized water was added into it as modifier for Hg. There were two chambers, reductant chamber with 0.3% diluted solution of NaBH_4 and 0.1% diluted solution of sodium hydroxide (NaOH) and another was hydrochloric acid (HCl) acid chamber with 3% concentrated HCl. The wave length of mercury was 253.7 nm.

Hydride Generation Atomic Absorption Spectrometer (HGAAS) technique was used to determine the concentration of arsenic in the samples. 2 ml of digested sample was taken in attest tube; 3M concentrated HCl with a flow rate of 7 ml/min and 1% diluted solution of KI, 1% of diluted solutions of ascorbic acid were added into the test tube for arsenic as modifier with reaction time 3.5 hr. The total volume of the liquid sample was 20 ml. The wave length of arsenic was 193.7 nm.

The standard solutions of mercury and arsenic and other heavy metal's standard solution were used to construct the calibration curves with the help of AAS. Quality assurance measures included the calculation of method detection limit, inclusion of recovery and analysis of standard reference material. A blank reading was also taken and necessary correlation was made during the calculation of concentration of different elements.

Results and Discussion

Total element analysis of soil

The level of heavy metals in the soils was compared to the Indian Standard [21-25]. The guidelines identify ecological investigation

levels (EILs), based on total metal concentration, considerations of phytotoxicity. The results of heavy metals concentration in the soil samples are presented in (Table 1). It was found that the Hg in the sampling station was below the detection limit (<0.03 mg/kg) and the concentration of Ni, Cu, Cd, Pb, Cr, Co were below the permissible limits recommended by Indian Standard Awasthi and European Union, [21-25] respectively which were shown in the Table 1. Whereas, the concentration of the zinc in the same soil samples were found in the ranged from 40.708 mg/kg to 448.469 mg/kg. The highest concentration of zinc (448.469 mg/kg) was observed in the soil of Pakshi station and the lowest amount of zinc was (7.639 mg/kg) was found in Diar Rooppur station. From the study we see that, the concentration of Zn (102.506 mg/kg and 448.409 mg/kg) in Chor Rooppur and paksh is higher than that of the permissible limit (50-100 mg/kg) recommended by WHO and Encyclopedia of Environmental Science. So, it can be concluded that the soils of Chor Rooppur and Pakshi contaminated by zinc. Although the soil of Paskhi is heavily contaminated as compared with Chor Rooppur.

The concentration of arsenic (As) in the collected soil samples were in the ranged from 2700 mg/kg to 5460.060 mg/kg. The highest concentration of arsenic (5460.060 mg/kg) was found in the soil of Pakshi and the lowest concentration (2700 mg/kg) was found in the soil of Diar Baghail. The WHO permissible limit for arsenic in agricultural soils is 0.5 mg/kg [25]. This limit of arsenic was exceeded by all the soil samples which were analyzed. The soils of the study areas contain large amount of arsenic, which indicates that soils are polluted by arsenic. Therefore it is assumed that the vegetables that are grown in these soils will absorb more arsenic from the soil and thereby polluted by arsenic. Plant arsenic concentrations tend to increase with increasing soil arsenic and then stabilize at some maximal value at higher concentrations in soil, which is alarming to the people of that area. The content of iron (Fe) was in the ranged from 11809.818 mg/kg to 25626.379 mg/kg. The highest concentration of iron (25626.379 mg/kg) was observed in the soil of Chor Rooppur and the lowest concentration was (11809.818 mg/kg) found in the soil of Diar Baghail. The maximum allowable limit of iron recommended by WHO (World Health Organization) is 150 mg/kg. The concentration of iron was too much high in the study areas compare with the maximum allowable limit of WHO.

The sources of iron, zinc and arsenic in the study area are mainly due to burning of fossil fuel and anthropogenic activities such as waste water irrigation, solid waste disposal and sludge applications. So from the above results, it is concluded that the soil of the study area was highly polluted by zinc, arsenic and iron; Pakshi and Chor Rooppur are most polluted area. Therefore as the plants in these areas usually uptake more amount of heavy metals, thereby these plants will certainly affect the human and other animals when these plants will intake by them.

Total element analysis in vegetables

The all metals concentrations are expressed on a plant fresh weight basis (FW). Metal concentrations calculated in a dry weight basis (DW) were converted to a FW basis by diluting the metal concentration according to the ratio of FW to DW. The results of heavy metals concentration in the vegetable samples are presented in (Table 2). The contamination of heavy metals in vegetables indicate that the concentration of Ni, Cd, Cr, Cu, Co, As (<0.1 mg/kg) and As (<0.03 mg/kg) were obtained below the detection limits; the concentrations of Fe, Mn, Zn were below the permissible limit recommended by WHO [26,27] respectively which were shown in the Table 2. But only the

Metals mg/kg	Soil 1	Soil 2	Soil 3	Soil 4	Soil 5	Soil 6	Standard
Ni	36.774 ± 1.55	22.899 ± 0.65	18.277 ± 0.376	22.969 ± 3.343	15.051 ± 0.177	23.607 ± 5.353	75-150 ^a
Cd	<0.1	0.806 ± 0.02	0.633 ± 0.036	0.387 ± 0.007	0.833 ± 0.012	0.467 ± 0.004	0.07-1.1
Cr	28.194 ± 0.17	17.1 ± 0.11	12.065 ± 0.326	17.567 ± 0.105	11.733 ± 0.035	15.333 ± 0.230	65 ^c
Cu	27.481 ± 2.09	14.935 ± 0.04	9.970 ± 0.556	54.326 ± 4.437	9.07 ± 0.472	12.849 ± 0.306	6-60 ^b
As	4200 ± 16.80	4346.667 ± 65.2	2700 ± 75.6	5460 ± 60.06	2940 ± 73.5	4353.33 ± 165.4	0.5 ^d
Hg	<0.03	<0.03	<0.030	<0.03	<0.03	<0.03	-
Co	15.129 ± 0.38	9 ± 0.207	7.748 ± 0.091	8.833 ± 0.141	7 ± 0.007	8.733 ± 0.183	10 ^c
Pb	21.290 ± 0.47	22 ± 0.440	21.290 ± 0.213	116.667 ± 0.233	116.089 ± 2.206	115.743 ± 1.620	10-70 ^b
Fe	25626.38 ± 1855.117	15491.539 ± 74.975	11889.922 ± 573.052	13569.992 ± 1221.374	11809.818 ± 99.499	15720.552 ± 40.034	150 ^c
Mn	504.839 ± 1.5	317.822 ± 2.225	107.102 ± 2.142	307.333 ± 1.229	202.327 ± 0.6-7	261.578 ± 1.308	437 ^c
Zn	60.839 ± 0.27	102.506 ± 4.204	45.177 ± 2.077	448.469 ± 25.021	40.708 ± 0.032	42.003 ± 3.328	50-100 ^e

Soil 1 and 2 - Chor Rooppur; Soil 3 - Diar Baghail; Soil 4 - Pakshi, Soil 5 and 6 - Diar Rooppur.

^aIndian standard awashthi and European Union, 2002; ^bFAO/WHO, codex general standard for contaminants and toxins in foods, 1996; ^cWorld Health Organization, 2000; ^dWorld Health organization, 2004; ^eWHO and encyclopedia environmental science.

Table 1: The concentration of heavy metals in soil using FAAS, CVAAS and HGAAS analytical method.

Metals mg/kg	Potato	Red amarantha	Spinach amarantha	Carrot	Cabbage	Tomato	Brinjal	Standard
Ni	<0.1	0.840 ± 0.01	0.54 ± 0.02	<0.1	<0.1	0.16 ± 0.01	<0.1	1.5 ^a
Cd	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	0.1 ^b 0.3 ^c
Cr	<0.1	<0.1	<0.1	<0.1	0.495 ± 0.01	0.75 ± 0.01	0.436 ± 0.01	2.3 ^b
Cu	ND	ND	ND	ND	ND	ND	ND	40 ^d
As	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	0.2 ^e
Hg	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	10 ^d
Co	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	0.05-0.1 ^f
Fe	68.671 ± 4.53	136.3 ± 7.07	58.094 ± 1.3	8.824 ± 1.68	7.276 ± 0.14	6.444 ± 1.8	6.933 ± 0.708	425 ^b
Mn	1.22 ± 0.016	5.720 ± 0.017	5.280 ± 0.063	1.257 ± 0.014	0.99 ± 0.005	1.41 ± 0.001	1.446 ± 0.033	500 ^b
Zn	3.093 ± 0.643	11.305 ± 0.57	8.487 ± 1.171	1.206 ± 0.365	2.652 ± 1.228	2.818 ± 1.86	3.213 ± 0.983	60 ^e
Pb	0.377 ± 0.02	1.036 ± 0.01	1.596 ± 0.01	0.304 ± 0.01	0.119 ± 0.01	0.161 ± 0.01	0.465 ± 0.01	0.3 ^b 0.2 ^g

^aWHO/FAO (Codex Alimentarius Commission. Joint FAO/WHO, 2007) and indian standard

awashthi; ^bWHO (Codex Alimentarius Commission, Joint FAO/WHO, 2001 and codex alimentarius commission, 1994); ^cEuropean Union (EU), 2006; ^dWHO/FAO (FAO/WHO, codex general standard for contamination and toxin in foods, 1996); ^eWHO (codex alimentarius commission, 1991); ^fAgency for toxic substance disease registry (ATSDR, 1994a); ^gChina food hygiene standard, 1994.

Table 2: Heavy metal concentration in vegetables using FAAS, CVAAS and HGAAS analytical method [32].

concentration of lead (Pb) in vegetables was found in toxic level. Which were varied from 0.119 mg/kg to 1.596 mg/kg. The highest lead content was found in spinach amarantha (1.596 mg/kg) while in cabbage it was lowest in concentration (0.119 mg/kg). According to China food hygiene standard, 1994 the standard limit of lead for vegetables and fruit is 0.2 mg/kg while it is 0.3 mg/kg for WHO (Codex Alimentarius Commission. Joint FAO/WHO). It is found from Table 2 that in all vegetables except carrot, lead concentration is more than permitted level, so they are not suitable for consumption. In the study area lead concentration that was found in the vegetables is a result of human activities such as waste water irrigation, solid waste disposal and sludge applications, solid waste combustion, agrochemicals and vehicular exhausted. The lead in fuels can contribute to the air pollution. It is highly unlikely that in that region, the traffic is so voluminous that the air pollution could convert to soil pollution in short term.

Lead is a toxic element that can be harmful to plants, although plants usually show ability to accumulate large amounts of lead without visible change in their appearance or yield. In many plants lead accumulation can exceed several hundred times the threshold of maximum level permissible for human [28]. The introduction of lead into the food chain may affect human health and may cause

disruption of the biosynthesis of hemoglobin and anemia, rise in blood pressure, kidney damage, miscarriages and subtle abortions, disruption of nervous systems and brain damage. Thus studies concerning lead accumulation in vegetables have increased important [29].

From the overall study of the heavy metals in vegetables it has been found that the results obtained for different parameters investigated in each category of vegetable samples were at normal levels except lead, which is harmful for human. Thus immediate actions are necessary to take to keep it within the permissible limit.

Relationship of heavy metals in soil and vegetables

Heavy metals bioavailability to plants is strongly related to the concentration and specification of the element in the soil solution because this is where the plants get the heavy metals that they take up. Typically, plants only take up one or two forms of heavy metals from the soil solution. The accumulation of metals from soils to plants depends on many factors such as metal forms, plant species and parts and soil properties. The solubility and consequently the plant uptake of the trace metals in the soils vary considerably with pH and the redox potential within the soil or the root system. The associated anion and the particle size also have a great influence on growth and

metal uptake by plants. Uptake of heavy metals by plants tends to increase with increasing concentration, as long as it is within a certain range. When the concentration goes beyond the range the uptake will decrease because plant roots are injured, thus leading to a lower absorbing ability. Therefore it is easy to make error if the soil pollution status of an area is determined simply from the contents of pollutants in the seeds. The enrichment factors quantify the relative differences in bioavailability of metals to plants and is a function of both soil and plant properties.

The present study showed that the concentrations of metals in vegetables were generally lower than that of the corresponding soils. This might be attributed to the root which seems to act as a barrier to the translocation of metals. In order to evaluate the accumulating capacity of heavy metals from soils to plant, a quantitative evaluation of the relationship between metals concentration in vegetable and in corresponding soils was made by calculating the transfer factor for the soil/plant system. Generally transfer factor varies from one plant to another plant, suggesting a selectivity of the plants in absorbing elements from soils.

Soil-vegetable transfer coefficients

The transfer coefficient quantifies the relative differences in bioavailability of metals to plants and is a function of both soil and plant properties. The coefficient is calculated by dividing the concentration of a metal in a vegetable crop (DW) by the total metal concentration in the soil. Higher transfer coefficient represents relatively poor retention in soils or greater efficiency of plants to absorb metals. Low coefficient demonstrates the strong sorption of metals to the soil colloids [30].

Soil-to-plant transfer is one of the key components of human exposure to metals through food chain. Transfer Factor (TF) or Plant Concentration Factor (PCF) is a parameter used to describe the transfer of trace elements from soil to plant body and it is also a function of both soil and vegetables properties. The transfer coefficient was calculated by dividing the concentration of heavy metals in vegetables by the total heavy metal concentration in the soil [2].

$$TF = \frac{C_{plant}}{C_{soil}}$$

Where, C_{plant} : metal concentration in vegetable tissue, mg kg⁻¹ and C_{soil} : metal concentration in soil, mg kg⁻¹.

In the present study, the TF of different heavy metal from soil to vegetable are presented in Table 3. Higher transfer factors reflect relatively poor retention in soils or greater efficiency of vegetables to absorb metals. Low transfer factor reflects the strong sorption of metals to the soil colloids [29]. The TF or PCF value ranges were: Cr

0.019-0.033, Pb 0.005-0.073, Fe 0.0003-0.0006, Mn 0.003-0.014 and Zn 0.015-0.138 and the trend of TF for heavy metal in vegetable samples studied were in order: Zn>Pb>Cr>Mn>Fe.

The mobility of metals from soil to plants is a function of the physical and chemical properties of the soil and of vegetable species, and is altered by innumerable environmental and human factors [31]. The highest TF value was found 0.077 and 0.039 for Zn and Pb. These might be due to higher mobility of these heavy metals with a natural occurrence in soil [17] and the low retention of them in the soil than other toxic cations [32]. According to the soil to plant transfer factor (TF) calculated for tested metals and leafy vegetables consumed by local residents, it can be concluded that Pb and Zn was high accumulator among the investigated metals. However, the higher concentrations of these heavy metals are due to the waste water irrigation, solid waste disposal and sludge applications, solid waste combustion, agrochemicals and vehicular exhausted.

Conclusion

The present study reveals the determination of the heavy metals in soil and vegetables. The result clearly indicate that some heavy metals like Zn, Fe, As and Pb have been build up in soil and thereby in plants mainly vegetables are responsible for contamination. But without the above four heavy metals contamination, the soil and vegetables in the study area were found to free from other heavy metal contamination. So, the soil in these areas is quite safe for cultivation and also the vegetables are safe for eating. But dietary intake of these vegetables result in long term low level body accumulation of heavy metals and the detrimental impact becomes apparent only after several years of exposure. Thus this study area is one of the more vegetable growing areas in Bangladesh. So regular monitoring of these toxic heavy metals in soil, in vegetables and other food materials is essential to prevent excessive build-up in the food chain. Furthermore, it is recommended that the study of heavy metals in environmental components in the proposed area of Rooppur nuclear power plant site of Bangladesh should be repeated at least two times in every year to know either any changes in contamination levels are occurring or not. The remediation of the contamination of soil and vegetables is necessary not only to preserve soil and vegetables but also to safeguard ecosystem.

Therefore, the main objective of this study was the assessment of heavy metals (Ni, Cd, Cu, Co, Pd, Fe, Mn, Zn, As, Hg) speciation in soil and vegetables in Pakshi area. Finally, the result thus presented in this paper is the only database available for the specification of heavy metals in soil and vegetable samples of this study area that will certainly help in better resources management, contributing to the effective monitoring of both environmental quality and will also provide information for background levels of metals of this studied area.

Metals	Ni	Cd	Cr	Cu	As	Hg	Co	Pb	Fe	Mn	Zn
Potato	-	-	-	-	-	-	-	0.017	0.003	0.003	0.037
Red amarantha	0.016	-	-	-	-	-	-	0.048	0.006	0.014	0.138
Spinach amarantha	0.015	-	-	-	-	-	-	0.073	0.003	0.013	0.104
Carrot	-	-	-	-	-	-	-	0.014	0.0004	0.003	0.015
Cabbage	-	-	0.022	-	-	-	-	0.005	0.0004	0.003	0.033
Tomato	0.015	-	0.033	-	-	-	-	0.007	0.0003	0.003	0.035
Brinjal	-	-	0.019	-	-	-	-	0.021	0.0003	0.004	0.039
Maximum	0.016	-	0.033	-	-	-	-	0.073	0.0003	0.014	0.138
Minimum	-	-	0.019	-	-	-	-	0.005	0.0006	0.003	0.015
Mean	0.008	-	0.026	-	-	-	-	0.039	0.00045	0.0085	0.077

Table 3: Transfer factors of heavy metal from Chor Rooppur soils into the vegetable samples.

Acknowledgments

The author wants to show gratitude to Azizul Maksud, (EO) and Shahidur Rahman Khan, JEO of the Analytical Chemistry Laboratory, Chemistry Division, Atomic Energy Center, Dhaka, Bangladesh for helping in laboratory work during the research.

References

- Heidarieh M, Maragheh MG, Shamami MA, Behgar M, Ziaei F, et al. (2013) Evaluate of heavy metal concentration in shrimp (*Penaeus semisulcatus*) and crab (*Portunus pelagicus*) with INAA method. Springerplus 2: 72.
- Kachenko AG, Singh B (2006) Heavy metals contamination in vegetables grown in urban and metal smelter contaminated sites in Australia. Water Air Soil Pollut 169: 101-123.
- Gupta UC, Subhas C, Gupta MD (2008) Selenium in soils and crops, its deficiencies in livestock and humans: Implications for management. Commun Soil Sci Plant Anal 29: 1791-1807.
- WHO (1995) Inorganic lead. Geneva, World Health Organization, International Programme on Chemical Safety. Environmental Health Criteria 165.
- Chailapakul O, Korsrisakul S, Siangproh W, Grudpan K (2007) Fast and simultaneous detection of heavy metals using a simple and reliable microchip-electrochemistry route: An alternative approach to food analysis. Talanta 74: 683-689.
- Wilson B, Pyatt FB (2007) Heavy metal dispersion, persistence, and bioaccumulation around an ancient copper mine situated in Anglesey, UK. Ecotoxicol Environ Saf 66: 224-231.
- Jofferries DJ (1984) Chemical analysis some coarse fish from a suffolk River carried out part of the preparation for the first release of captive-bred otters. J Otter Trust 1: 17-22.
- Freedman B (1989) Environmental Ecology: The impacts of pollution and other stresses on ecosystem structure and function. New York: Academic Press Inc.
- Whitmuff MS (2002) Applying biosolids to acid soil in New South Wales: Are guideline soil metal limits from other countries appropriate? Aust J Soil Res 40: 1041-1056.
- McBride MB (2003) Toxic metals in sewage sludge-amended soils: has promotion of beneficial use discounted the risks? Adv Environ Res 8: 5-19.
- Voutsas D, Grimanis A, Samara C (1996) Trace elements in vegetables grown in an industrial area in relation to soil and air particulate matter. Environ Pollut 94: 325-335.
- Jinadasa KBPN, Milham PJ, Hawkins CA, Cornish PSD, Williams PA, et al. (1997) Survey of cadmium levels in vegetables and soils of greater Sydney, Australia. J Environ Qual 26: 924-933.
- Lehoczy E, Szabo L, Horvath S, Marth P, Szabados I (1998) Cadmium uptake by lettuce in different soils. Commun Soil Sci Plant Anal 28: 1903-1912.
- Muchuweti M, Birkett JW, Chinyanga E, Zvauya R, Scrimshaw MD, et al. (2006) Heavy metals content of vegetables irrigated with mixture of waste water and sewage sludge in Zimbabwe: Implication for human health. Agric Ecos Environ 112: 41-48.
- Mapanda F, Mangwayana EN, Nyamangara J, Giller KE (2005) Impacts of sewage irrigation on heavy metals distribution and contamination. Environ Intern 31: 05-812.
- Bahemuka TE, Mubofu EB (1991) Heavy metals in edible green vegetables grown along the sites of the Sinza and Msimbazi Rivers in Dar es Salaam, Tanzania. Food Chemistry 66: 63-66.
- Alam MG, Snow ET, Tanaka A (2003) Arsenic and heavy metal contamination of vegetables grown in Samta village, Bangladesh. Sci Total Environ 308: 83-96.
- Ikeda M, Zhang ZW, Shimbo S, Watanabe T, Nakatsuka H, et al. (2000) Urban population exposure to lead and cadmium in east and south-east Asia. Sci Total Environ 249: 373-384.
- Iyengar GV, Nair PP (2000) Global outlook on nutrition and the environment: meeting the challenges of the next millennium. Sci Total Environ 249: 331-346.
- Türkdoğan MK, Kilicel F, Kara K, Tuncer I, Uygan I (2003) Heavy metals in soil, vegetables and fruits in the endemic upper gastrointestinal cancer region of Turkey. Environ Toxicol Pharmacol 13: 175-179.
- Awashthi SK (2000) Prevention of Food Adulteration Act no. 37 of 1954. Central and State Rules as Amended for 1999, Ashoka Law House, New Delhi.
- European Union (2000) Heavy Metals in Wastes, European Commission on Environment.
- Codex Alimentarius Commission (1996) Joint FAO/WHO food standards programme. Codex General Standard for Contaminants and Toxins in Foods. Doc No. Cx/FAC 96/17.
- WHO (2000) Safety evaluation of certain food additives and contaminants. International Programme on Chemical Safety. WHO Food Additive Series 52.
- WHO (2004) Task group on environmental health criteria for arsenic and Arsenic Compounds 18, World Health Organization, Geneva, Switzerland 1-174.
- Chinese Department of Preventive Medicine (1994) Threshold for food hygiene. Beijing:
- China Standard press (In Chinese).
- Codex Alimentarius Commission. Joint FAO/WHO (2001) Food additives and contaminants. Joint FAO/WHO food Standards program; ALINORM 01/12A: 1-289.
- Wierzbicka M (1995) How lead loses its toxicity to plants. Acta Soc Bot Pol 64: 81-90.
- Coutate TP (1992) Food, the chemistry of its component (2nd edn) Cambridge: Royal society of chemistry Pp. 265.
- Alloway BJ, Ayres CD (1997) Chemical principals of environmental pollution, (2nd edn) Blackie Academic and Professional, London.
- Zurera G, Estrada B, Rincón F, Pozo R (1987) Lead and cadmium contamination levels in edible vegetables. Bull Environ Contam Toxicol 38: 805-812.
- Lokeshwari H, Chandrappa GT (2006) Impact of heavy metal contamination of Bellandur lake on soil and cultivated vegetation. Curr Sci 91: 622-627.