

## Insects as Biological Weapons

Fahad Nazir Chaudhry\*, Muhammad Faheem Malik, Mubashar Hussain and Nayab Asif

Department of Zoology, University of Gujrat, Hafiz Hayat Campus, Pakistan

\*Corresponding author: Fahad Nazir Chaudhry, Department of Zoology, University of Gujrat, Hafiz Hayat Campus, Gujrat, Pakistan, Tel: + 410-735-6409; E-mail: fahadnazirch55@gmail.com

Received date: April 22, 2017; Accepted date: July 26, 2017; Published date: July 31, 2017

Copyright: © 2018 Chaudhary FN, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

### Abstract

This study was conducted at University of Gujrat during 2017 to 2018 as a term paper for Master of Philosophy. The data regarding use of Insects as Biological Weapons was reviewed and compiled as a review paper from various published articles of international reputed journals annual/environmental reports of recognized organization and e-books. Use of entomological weapons is a wide field of research. Insects were used as biological weapons from ancient times and this concept is very old. Different insects can be used for this purpose by different ways. Insects can be used as biological weapons against crops, animals and human beings. Insects may be used as direct pests against crop or they may be used as vectors to spread diseases in animals and human beings. In both cases their effects are devastating. House Flies, Oriental Rat Flea, Colorado potato beetle, Ticks, screw worm larvae are the examples of some insects which can be used as biological weapons. Use of biological weapons in war against humans is however illegal.

**Keywords:** Biological weapons; Entomological warfare; Agro terrorism and Legal status

### Introduction

This concept of using insects as biological weapons is very old and even today research on this topic is carried out by many countries. In 14th century in Asia Minor plague which was spread through fleas also known as black death was the on the of earliest event of using insect as biological weapons it was used in Crimean against the city of Kaffa [1]. Another earliest use of insects as biological weapons by earliest humans was the use of bees for direct attack [2,3]. Japan and many other countries were accused of using Entomological warfare in World War 2. The only verified use of Entomological warfare on large scale is by Japanese against Chinese in World War 2. It is type of biological warfare [4]. In EW insect are used as a direct attack or used as vectors to spread disease through specific biological agent [5,6].

### History

Insects are being used as biological weapons since ancient times. In 14th century fleas are used to spread plague against the city of Kaffa in Asia Minor [1]. In World War 2 Germans used Colorado beetle against enemy crops [7]. Japanese also used plague infected fleas and cholera infected flies against Chinese in World War 2 [5,8]. Many techniques are developed by soviets during cold war era to transmit diseases like foot and mouth disease using ticks. But however they did not used them against any country [9]. Similarly during Cold war USA also developed a laboratory capable of producing millions of yellow fever infected mosquitos to attack soviets and they also did experiment on their survivability if dropped from plane [7,10].

### Biological Weapons

It is use of Living organisms like Viruses, Bacteria, Fungi, protozoa and insects or their toxins to cause disease or kill humans, animals and plants.

### Entomological Warfare

It is use of insects to attack enemy. It is type of biological warfare in which insects are used as biological weapons. Different insects can be used by different ways in entomological warfare [4].

### Different ways to use insects as biological weapons

EW can be used by 3 different ways [5]. For this purpose insects are first infected with pathogens then they are dispersed on an area which is to be targeted. These insects act as a vector and infect particular animal or people of targeted area. In second type insects are directly used to destroy crops. In this case insects may not cause damage by pathogens or by acting as a vector. In third type uninfected insects are directly used as a direct attack on enemy like bees [5].

### Agro Terrorism

It is the destruction or disruption of food and agriculture industry by using pant pathogens or pests.

### Colorado Potato Beetle

It is major potato pest and its female can lay up to 800 eggs. It has orange and yellow body with five strips on its elytra [11]. It is similar to false potato beetle. It is native to Colorado and Mexico [12]. With the passage of time Colorado potato beetle have become resistant to and dichloro-diphenyl-trichloroethane, other major pesticides classes [11,13]. Colorado potato beetle was also produced in large number by Germans during World War 2 to destroy enemy food sources [7]. They released 54000 Colorado beetles in south Frankfurt to check their

effect [14]. French entomologists also concluded that Colorado beetle can be used as biological weapon against enemy crops [7]. During Cold war CIA was also blamed to use it against soviet crops [15,16]. 15000 Colorado beetles are sent by USA to Britain in 1942 to study them as a biological weapon [14].

### Oriental Rat Flea

It is the vector of murine typhus and bubonic plague, and these diseases are transmitted from one flea generation to the next by their eggs [17]. Japan used these fleas on large scale as a biological weapon against Chinese in World War 2. This program was led by Lt. General Shirō Ishii [5]. These plague fleas were dropped by low flying planes on the city of Changde and caused plague which resulted in 500,000 deaths of Chinese [5,8].

### House Fly

House fly can act as a lethal biological weapon. It is vector of up to 100 different kinds of pathogens which can cause diseases like cholera, typhoid, salmonellosis, anthrax, tuberculosis, ophthalmic, bacillary dysentery and parasitic worms. It is also an important vector of different viruses like poliomyelitis, enteroviruses and viral hepatitis [18-20]. Some strains of house fly are also resistant to many kinds of pesticides commonly used [21,22]. Japanese also used cholera-infected flies as an entomological weapon against Chinese to spread disease during World War 2 [5].

### Ticks

Ticks can also be used as biological weapons to spread different diseases in poultry and livestock and humans. They are the carriers of many bacteria, protozoans and viruses [23]. Aegyptianellosis is a common disease found in poultry caused by *Argas* spp. of ticks. It spreads from one bird to another [24]. Similarly, cattle fever tick is a vector of *Babesia bigemina* which causes cattle fever. *Babesia bovis* can also be transmitted by this tick. Babesiosis causes \$3 billion annual losses to the U.S. cattle industry every year in the early 20th century [23]. During the Cold War era, the Soviet Union developed techniques to transmit diseases like foot and mouth disease using ticks. Avian ticks are also used by them to transmit *Chlamydia psittaci* in chickens [9]. In humans, they are responsible for causing rickettsialpox, typhus, African tick bite fever, Boutonneuse fever, Rocky Mountain spotted fever, Queensland tick typhus, Flinders Island spotted fever, Colorado tick fever, Q fever, tick-borne meningoencephalitis and Ehrlichiosis [25].

### Mosquitos

Mosquitos can also be used as biological weapons to cause disease in birds, animals and humans. They are the vectors of different viruses and parasites. Viral diseases include Dengue fever, chikungunya and yellow fever which are mostly caused by *Aedes aegyptus* [26]. Similarly, malaria is caused by a protozoan called *Plasmodium* [27]. During the cold war, the USA made a laboratory which can produce 100 million yellow fever-infected mosquitos to attack the Soviet Union [7]. In 1955, the United States dropped 300,000 yellow fever-infected mosquitos during an operation to check their survivability [10].

### Caterpillars

Caterpillar is the larvae of order Lepidoptera members which include butterflies and moths. Sawfly larvae are also known as

caterpillar [28]. Most caterpillars are herbivorous and agricultural pests [29]. They cause damage by eating leaves and many species have become resistant to many pesticides [30]. In 1990, the USA spent 6.5 million dollars on research to use caterpillars as biological weapons against crops [31].

### Black Flies

They are 3 to 6 mm in size, small, robust flies with short wings and humped thorax. They are also known as turkey gnats and buffalo gnats. They feed on the blood of poultry and cattle. If flies attack in large numbers, they cause the death of animals. Death is caused in many cases due to acute toxemia. Productivity is even reduced by low population of flies. They are also vectors of protozoans (*Leucocytozoon*) which cause leucocytozoonosis in poultry and filarial nematodes (*Onchocerca*) which cause bovine onchocerciasis [23,32-34]. In humans, black flies transmit onchocerciasis; they are vectors of parasitic nematode *Onchocerca volvulus*. The parasite lives in the skin of humans and is transmitted when flies take a meal during feeding, so they can also be used as biological weapons to transmit disease in humans, poultry and cattle [35].

### Biting Midges

They are 1 to 4 mm long flies which belong to order Diptera. They are also known as punkies, sand-flies and no-see-ums [36]. They are persistent and vicious biters [23]. They are vectors of arboviruses and different non-viral pathogens [37,38]. They are vectors of a virus that causes blue tongue in cattle and in sheep. In cattle, epizootic haemorrhagic disease is also caused by them. In poultry, they spread blood protozoans and equine onchocerciasis [23]. So, artificially infected biting midge colonies can be used to spread disease among cattle and poultry.

### Horse Flies and Deer flies

They can also be used as biological weapons as they are livestock's major pest, especially cattle, transmitting different diseases in them. They are robust and large in size. They have blade-like mouth parts which are designed by nature to inflict deep bleeding wounds [23,39]. They can transmit diseases like bovine leukemia, infectious anemia, hog cholera, *Trypanosoma* sp. Protozoans and *Elaeophora* sp. Nematodes. [23,40].

### Warble Fly

They are large flies which are parasites of cattle. They are also known as gad flies, bomb flies or heel flies. Their larva is called warbles or cattle grub. Some species of larvae also invade human tissues. They are large, hairy, orange to yellow in colour and bee-like in appearance. The adults are free living and have vestigial mouth parts [41].

Warble fly larvae which are known as cattle grubs can be used as biological weapons as in 1976, the U.S. alone faced an estimated \$360 million loss due to cattle grubs. When tissue-invading cattle grub larvae reach the oesophagus and spinal cord, they cause paralysis [23]. In humans, *H. bovis* species larvae cause a disease called intracerebral myiasis in which invasion of intracerebral tissues takes place. Symptoms of this disease include intracerebral hematoma and convulsions [42]. *H. tarandi*, which is a parasite of caribou, is responsible for human eye myiasis which causes glaucoma, uveitis, and retinal detachment [43]. So, warble

flies if produced in large scale in laboratory can be used as anti-livestock biological weapon.

### Screw Worm Fly

Screw worm fly can also be used as biological weapon against livestock animals. It is parasitic fly whose larvae infests on open wounds of warm blooded animals [44]. Their larvae attack only healthy living tissues of their host. It is major pest of cattle and other livestock animals [23]. Screw worm can cause tissue loss, vital organ destruction and even death in extreme cases [45]. Screw worm female can lay up to 3000 eggs in her lifecycle and travel up to 200km in the search of host [46].

### Legal status of entomological weapons

Insect vectors are not mentioned in the text of Biological and Toxic Weapons Convention (BWC) of 1972. However vectors are covered in treaty [47]. Use of vectors in armed conflict for hostile purposes is banned by article 1 of BWC [47,48]. So BWC covers insect vectors [49]. However the use of uninfected insects against crops is not clear [50].

### Conclusion and Recommendations

Use of insects as biological weapons is very cheap and effective warfare. They can easily be used to spread disease among enemy and to destroy enemy crops and livestock. But unlike conventional weapons they act slowly and their use in war is illegal and considered as war crime so there should be laws and regulations to prohibit their proliferation or to be used in war. There also should be check and balance to prevent biological weapons going into hands of terrorists and to be used to spread terrorism. Airport security and export and import security should also be increased so that biological weapons may not be smuggled from one country to another for this purpose there should be entomological experts in anti-terrorism investigation teams, borders and airport security teams.

### References

1. Kirby Reid (2005) Using the flea as weapon. *Army Chemical Review*.
2. Pete B (2008) Warfare gets the creepy-crawlies. *Laramie Boomerang*.
3. (2005) UW Professor Examines Biological Setting of Egyptian Plagues (Press release), University of Wyoming.
4. Peterson RKD (1990) The Role of Insects as a Biological Weapon, Department of Entomology, Montana State University, notes based on seminar.
5. Lockwood JA (2008) Six-legged Soldiers Using Insects as Weapons of War, Oxford University Press, USA. 9–26.
6. (2002) An Introduction to Biological Weapons Their Prohibition, and the Relationship to Biosafety. The Sunshine Project.
7. Lockwood JA (2008) "Bug Bomb", *Boston Globe*.
8. Novick LF, John SM (2001) Public Health Issues Disaster Preparedness, Jones & Bartlett Publishers.
9. Jonathan B (2008) Agricultural Biological Warfare: An Overview. The Arena, Paper # 9, Chemical and Biological Arms Control Institute, via Memorial Institute for the Prevention of Terrorism.
10. William RH (1981) An Evaluation of Entomological Warfare as a Potential Danger to the United States and European NATO Nations, U.S. Army Test and Evaluation Command, Dugway Proving Ground, via thesmokinggun.com.
11. Capinera JL (2001) Handbook of Vegetable Pests. Academic Press, San Diego, USA.
12. Arnett Jr RH, Thomas MC, Skeppey PE, Frank JH (2002) American Beetles, Vol 2. CRC Press. Boca Raton, USA.
13. Alyokhin A, Baker M, Mota-Sanchez D, Dively G, Grafius E (2008) Colorado potato beetle resistance to insecticides. *American Journal of Potato Research* 85: 395–413.
14. Heather NW, Hallman GJ (2008) Pest Management and Phytosanitary Trade Barriers CABI, 17–18.
15. Croddy EA, Wirtz JJ (2005) Weapons of Mass Destruction: An Encyclopedia of Worldwide Policy, Technology, and History ABC-CLIO.
16. Daisy S (2014) What's Orange and Black and Bugging Ukraine?. *Radio Free Europe / Radio Liberty*.
17. Farhang-Azad A, Traub R, Baqar S (1985) Transovarial transmission of murine typhus rickettsiae in *Xenopsylla cheopis* fleas. *Science* 227: 543–545.
18. Ostrolenk M, Welch H (1942) The House fly as a vector of food poisoning organisms in food producing establishments. *Am J Public Health Nations Health* 32: 487–494.
19. Levine OS, Levine MM (1991) House flies (*Musca domestica*) as mechanical vectors of shigellosis. *Rev Infect Dis* 13: 688–696.
20. Förster M, Klimpel S, Sievert K (2009) The house fly (*Musca domestica*) as a potential vector of metazoan parasites caught in a pig-pen in Germany. *Veterinary Parasitology* 160: 163–167.
21. Georghiou GP, Hawley MK (1971) Insecticide resistance resulting from sequential selection of houseflies in the field by organophosphorus compounds. *Bulletin of the World Health Organization* 45: 43–51.
22. Keiding J (1975) Problems of housefly (*Musca domestica*) control due to multiresistance to insecticides. *Journal of Hygiene, Epidemiology, Microbiology, and Immunology* 19: 340–355.
23. *Veterinary entomology* (2010) Livestock and Companion Animals Ralph E. Williams, Purdue University West Lafayette, Indiana, U.S.A.
24. Euzéby JP (1997) List of Bacterial Names with Standing in Nomenclature: a folder available on the Internet. *Int J Syst Bacteriol* 47: 590–592.
25. Unsworth NB, Stenos J, Graves SR, Faa AG, Cox GE, et al. (2007) Flinders Island spotted fever rickettsioses caused by "marmionii" strain of *Rickettsia honei*, Eastern Australia. *Emerg Infect Dis* 13: 566–573.
26. WHO (2009) Dengue Guidelines for Diagnosis, Treatment, Prevention and Control. Geneva: World Health Organization.
27. Worrall E, Basu S, Hanson K (2005) Is malaria a disease of poverty? A review of the literature. *Trop Med Int Health* 10: 1047–1059.
28. Eleanor Anne Ormerod (1892) A text-book of agricultural entomology: being a guide to methods of insect life and means of prevention of insect ravage for the use of agriculturists and agricultural students. Simpkin, Marshall, Hamilton, Kent & Co.
29. Greene E (1989) A Diet-Induced Developmental Polymorphism in a Caterpillar. *Science*. 243: 643–646.
30. Redd JT, Voorhees RE, Török TJ (2007) Outbreak of lepidopterism at a Boy Scout camp. *J Am Acad Dermatol* 56: 952–955.
31. Irwin ME, GE Kampmeier (2008) Commercial products, from insects, (p. 6). In Resh, V.H. & Carde, R. eds. *Encyclopedia of Insects*, Academic Press, San Diego, via University of Illinois and Illinois Natural History Survey.
32. Daley B (2008) Black flies surge in Maine's clean rivers. *Boston Globe*.
33. Hough A (2010) Blandford fly: surge in 'infected' insect bites blamed on new superfly. *The Daily Telegraph*. London.
34. Mullen G, Durden L (2009) Medical and Veterinary Entomology. Academic Press.
35. Service MW (2008) Medical Entomology for Students. Cambridge University Press. 81–92.
36. Whelan P (2003) Biting Midges or 'Sand-flies' in the Northern Territory. *The Northern Territory Disease Control Bulletin*.

37. Carpenter S, Groschup MH, Garros C, Felipe-Bauer ML, Purse BV (2013) Culicoides biting midges, arboviruses and public health in Europe. *Antiviral Res* 100: 102–113.
38. Linley JR (1985) Biting Midges (Diptera: Ceratopogonidae) as Vectors of Nonviral Animal Pathogens. *J Med Entomol* 22: 589–599.
39. Kazimírová M, Sulanová M, Kozánek M, Takác P, Labuda M, et al. (2001) Identification of anticoagulant activities in salivary gland extracts of four horse-fly species (Diptera, Tabanidae). *Haemostasis*. 31: 294–305.
40. Wilkerson RC, Butler JF, Pechuman LL (1985) Swarming, hovering and mating behavior of male horse flies and deer flies (Diptera: Tabanidae)". *Myia* 3: 515–546.
41. Piper R (2007) *Extraordinary Animals: An Encyclopedia of Curious and Unusual Animals*. Greenwood Press
42. Kalelioğlu M, Aktürk G, Aktürk F, Komsuoğlu SS, Kuzeyli K, et al. (1989) Intracerebral myiasis from *Hypoderma bovis* larva in a child. Case report. *J Neurosurg* 71: 929–931.
43. Lagacé-Wiens PRS, Dookeran R, Skinner S, Leicht R, Colwell DD, et al. (2008) Human ophthalmomyiasis interna caused by *Hypoderma tarandi*, Northern Canada. *Emerg Infect Dis* 14: 64–66.
44. Curran J (2002) "Screw-Worm Fly". Government of Western Australia: Department of Agriculture Farmnotes.
45. (2000) California Department of Food and Agriculture Animal Health Branch. "Fact Sheet: Screwworm".
46. James MT (1947) *The Flies That Cause Myiasis in Man*. USDA Miscellaneous Publication No. 631.
47. Sims, Alan NR (2001) (Stockholm International Peace Research Institute). *The Evolution of Biological Disarmament*, Oxford University Press 45–50.
48. (2001) *Convention on the Prohibition of the Development, Production and Stockpiling of Bacteriological (Biological) and Toxin Weapons and on Their Destruction*.
49. Zanders JP (2004) *Research Policies, BW Development & Disarmament, Conference: "Ethical Implications of Scientific Research on Bioweapons and Prevention of Bioterrorism"*, European Commission, via Bio Weapons Prevention Project.
50. Regalado A (2016) *Top U.S. Intelligence Official Calls Gene Editing a WMD Threat*. MIT Technology.