

# Physico-Chemical Characterization and Analytical Development for Sodium Azumolene, a Potential Drug Designed to Fight Malignant Hyperthermia

Josilene Chaves Ruela Corrêa\*, Marcelo Augusto Cavaretti Hiene and Hérica Regina Nunes Salgado

Department of Pharmaceutics, School of Pharmaceutical Sciences, Univ Estadual Paulista, Brazil

## Abstract

Sodium azumolene is a drug designed to fight Malignant Hyperthermia (MH), which is characterized by genetic predisposition and triggered by the use of inhalational anesthetics. This drug is shown as a water-soluble analogue of dantrolene sodium, 30-folds more water soluble, which gives advantages for its emergency use. To our knowledge there is no analytical method for sodium zaumolene raw material or dosage form published so far. The objective of the present investigation was to develop and validate analytical methods to achieve sodium azumolene chemical identification and quantification. The sodium azumolene was characterized regarding its thermal behavior, by differential thermal analysis and thermogravimetric analysis; Visible, UV and infrared absorption. To accurately assess the sodium Azumolene content three different analytical methods (visible and UV spectrophotometry and high performance liquid chromatography) were developed and validated. All methods showed to be linear, accurate, precise and reliable. Azumolene has shown to be equipotent to dantrolene in the treatment and prevention of an MH crisis and the great advantage compared to dantrolene is better water solubility. This study has characterized the sodium azumolene and presents new analytical methods which have not been reported so far.

**Keywords:** Malignant hyperthermia; Sodium azumolene; UV-VIS Spectrophotometry; Liquid chromatography; Quality control

## Introduction

Malignant hyperthermia, which so far has no cure defined, is also known as malignant fever, malignant hyperpyrexia and anesthetic fever. It is hereditary (autosomal dominant) and latent, it is characterized by hypermetabolic response to volatile anesthetics (halothane, isoflurane, sevoflurane, isoflurane, enflurane and desflurane) and succinylcholine, initially from a derangement of intracellular calcium homeostasis [1-4].

It can occur at any time of anesthesia procedure and even some hours after. Its incidence has been reported with variations from 1/20.000 to 1/100.000 users of the triggering anesthetic agents cited above. The incidence of malignant hyperthermia in aesthetician procedure is of 1/15.000 in children and 1/50.000 in adults. There are studies reporting that 50% of cases occur in people less than 15 years [5-7].

Whereas the evolution of the disease can be fatal and occurs quickly, early diagnosis and specific treatment are of great importance, reducing mortality from 80% to less than 10% of the cases [8,9]. Thus, encountering the signs described for the disease associated with the use of volatile anesthetics agents, it must be considered a case of malignant hyperthermia and specific treatment protocol must be initiated [8,9]. There should be immediate cessation of the exposure to possible agents triggering of malignant hyperthermia with replacement by safe anesthetic drugs, followed by administration of dantrolene sodium or sodium azumolene, with the goal of correcting cellular metabolism of calcium, cardiac function, body temperature and respiration rate [7,10-12].

Although there is no central action, dantrolene, a hydantoin derivative, acts as a central muscle relaxant with the receptor binding site ryanodine, which acts through inhibitory mechanism [9,13]. Azumolene sodium has chemical structure, potency and efficacy similar to dantrolene sodium. It prevents or reverses malignant hyperthermia, which was confirmed by *in vitro* and *in vivo* studies [12,14]. According to Zhao and coworkers [15] both drugs are satisfactory and have equivalent applications to treat the irregular release of calcium by the endoplasmic reticulum.

The use of sodium azumolene presents advantages in relation to

its branch dantrolene, since the latter shows poor water solubility and requires a dilution of 20 mg of the lyophilized drug in 60 ml water for injection. Sodium azumolene is many times more soluble in its administration vehicle than dantrolene and provides an easier and faster drug administration procedure which is necessary for emergency situations [9,16,17].

The chemical structure of sodium azumolene (1-(((5-(4-bromophenyl)-2-oxazolyl)methylene)amino)-2,4-imidazolidinedione sodium) is shown at Figure 1.

To our knowledge there is no analytical method for sodium zaumolene raw material or dosage form published so far. There is no monograph for sodium azumole in any pharmacopeia. The aims of this




Figure 1: Chemical structure of sodium azumolene.

\*Corresponding author: Josilene Chaves Ruela Corrêa, Department of Pharmaceutics, School of Pharmaceutical Sciences, Univ Estadual Paulista, Rodovia Araraquara-Jaú, km 1, 14801-902, Araraquara, Brazil, Tel: +55 16 3301 4681; E-mail: [josilenechavescorrea@gmail.com](mailto:josilenechavescorrea@gmail.com)

Received November 20, 2013; Accepted December 19, 2013; Published December 21, 2013

Citation: Corrêa JCR, Hiene MAC, Salgado HRN (2013) Physico-Chemical Characterization and Analytical Development for Sodium Azumolene, a Potential Drug Designed to Fight Malignant Hyperthermia. J Anal Bioanal Tech 5: 177. doi:10.4172/2155-9872.1000177

Copyright: © 2013 Corrêa JCR, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

study were to provide reliable analytical methods as tools to analyze sodium azumolene for routine use.

## Material and Methods

### Chemicals

Sodium azumolene chemical reference (assigned purity of 100%) and the product as powder for injectable solution were kindly donated by Cristalia (Itapira, Brazil).

Methanol LC grade was purchased from Tedia (Fairfield, USA). Ultrapurified water was prepared in-house by using Direct-Q® water system (Millipore Corporation, Billerica, USA). Prior to use, mobile phase solvents were degassed in an ultrasonic bath for 30 min. Solvents were filtered through a 0.45 µm membrane filter.

### Apparatus

Liquid Chromatography apparatus (Waters Corporation, Milford, MA, USA), equipped with a Waters 1525 binary pump, a Rheodyne Breeze 7725i manual injector, a Waters 2487 UV-VIS wavelength detector was used. In addition a Liquid Chromatography apparatus HPLC analysis was conducted by using a RP C18 column (Waters, Milford, MA, USA), with 5 µm particle size; 4.6×250 mm.

For thermal analysis the SDT 2960 (TA Instruments) was used to do the differential scanning calorimetry and thermal gravimetry analyses simultaneously. For UV-VIS analysis a spectrophotometer UV mini-1240 Shimadzu (Columbia USA) was used. Spectrometer of infrared Shimadzu FTIR 3000 (Columbia, USA) was used.

### Validation of Methods

All LC, UV and VIS spectrophotometric methods were validated according to the International Conference on Harmonization (ICH) guidelines [18]. The following characteristics were considered for validation: Linearity, range, accuracy and precision.

Linearity was determined in the range 5.0 to 15.0 µg/mL (LC method); 7.0 to 12.0 µg/mL (UV spectrophotometric method) and 8.0 to 13.0 µg/mL (Vis spectrophotometric method). The least squared method was used to calculate the regression coefficient from peak area or absorption vs. concentration of standard solutions.

Accuracy was tested by calculating the percentage recoveries of sodium azumolene from sample solutions at different concentrations and by determining the relative standard deviation (RSD). Precision was assessed at different levels—repeatability (system repeatability by testing three curves constructed with five different standard solution concentrations, in the same day); and intermediate (by analyzing three different standard solution concentrations high, intermediate and low, in a different day, with two days interval between repeatability and intermediate).

The limits of detection (LOD) and quantification (LOQ) were calculated as 3 and 10  $\sigma/S$ , respectively, where S is the slope of the calibration curve and  $\sigma$  is the standard deviation of y-intercept of regression equation [18].

### LC conditions

Chromatographic analysis was performed in isocratic mode. Mobile phase consists of methanol-water (75:25, v/v, pH 3.0 adjust with formic acid) at a flow rate of 1 mL/min. The injection volume was 20 µL and the detection wavelength was 340 nm. The retention time of sodium azumolene was 4.25 min (Figure 2). All experiments were




Figure 2: Chromatogram of sodium azumolene diluted in mobile phase at 10.0 µg/mL and monitored at 340 nm.

performed at room temperature (25°C) and the total area of peak was used to quantify sodium azumolene.

**Preparation of standard and sample solutions to validate the LC method:** Six standard and sample aqueous solutions of sodium azumolene were prepared at the concentrations 5.0; 7.0; 9.0; 11.0; 13.0 and 15.0 µg/mL. All these solution were evaluated at 340 nm.

### UV Spectrophotometric conditions

The UV spectrum of sodium azumolene was obtained using an aqueous solution of this drug at 10 µg/mL. The absorption peak could be observed at 339 nm. This wavelength was used for the drug determination.

**Preparation of standard and sample solutions to validate the UV Spectrophotometric method:** Solutions of sodium azumolene were prepared at the concentrations 7.0; 8.0; 9.0; 10.0; 11.0 and 12.0 µg/mL. Standard and sample solutions were evaluated at 339 nm.

### VIS Spectrophotometric conditions

For the visible determination a purple color was obtained by addition of 0.1% chloranilic acid in a sodium azumolene solution prepared with acetonitrile at 10.0 µg/mL. This coloring reaction was described by Melby and coworkers [19]. After the staining process the visible spectrum of the solution was obtained and an absorption peak could be observed at 507 nm.

Charge transfer reactions can be ideal for visible analysis methodology. The chloranilic acid is an acceptor of electrons and it is widely used for photometric analysis in visible region of spectre to penicillins [20], cephalosporins [21], antimalarials [22], antifungals [23] and propranolol [24]. In a literature search of sodium azumolene it was not found the association of the drug with the chloranilic acid.

The mix of chloranilic acid and acetonitrile shows a VIS absorption peak near to 390 nm. However after addition of this mix to sodium azumolene the color immediately changes from yellow to purple and the absorption peak also changes to 507. This reaction may be explained by the occurrence of ion pairing [25,26].

**Preparation of standard and sample solutions to validate the VIS Spectrophotometric method:** Standard and sample solutions of sodium azumolene were prepared at 8.0; 9.0; 10.0; 11.0; 12.0 and 13.0 µg/mL using 0.1% of chloranilic acid in acetonitrile. All these solution were evaluated at 507 nm.

## Results and Discussion

### Validation of LC Method

The LC method validated is fast (retention time=4.25 min) and optimized method. Good symmetry was obtained by using endcapped

column (C<sub>18</sub> 4.6×250 mm; 5 μm) and mix of methanol and water (75:25, v/v, pH 3.0 adjust with formic acid) as mobile phase.

Good linearity was obtained at the concentration range tested. It was evaluated by the analysis of variance (ANOVA) and results showed regression statistically significant (F<sub>calculated</sub><F<sub>critical</sub>; P=0.05). The equation of the curve was y=44299X+137924 with a good coefficient of determination (r<sup>2</sup>=0.9990).

The precision of the method was determined by repeatability (intra-day) and intermediate precision (inter-day) and was expressed as R.S.D. (%) of a series of measurement. The repeatability test shows R.S.D. equal to 0.08 and the inter-day variability was equal to 1.23% (Table 1). The accuracy of the method was determined by the mean recovery, by adding standard to sample. The mean result was 98.14 % (Table 1) indicating an agreement between the true value and the value found.

LOD and LOQ were determined based on the standard deviation of the response and the slope, based on the calibration curve [18], and the values were 0.95 and 2.89 μg/mL, respectively (Table 1).

To evaluate the robustness of the proposed method low and deliberate changes were made on the flux rate of mobile phase. Sample and standard solutions were submitted at the following flux rate 0.9; 1.0 and 1.1 mL/min. No interference over assay test was found.

### Validation of UV Spectrophotometric Method

The developed UV spectrophotometric method has an easy procedure; moreover it is fast and low cost. The UV method shows to be linear at the concentration range tested (7.0 to 12.0 mg/mL). The equation of the curve was y=0.0868X-0.0425 with a good coefficient of determination (r<sup>2</sup>=0.9995). The validity of the linearity test was evaluated by analysis of variance (ANOVA) which shows the statistically significance of the test.

The precision of the method was determined by repeatability (intra-day) and intermediate precision (inter-day) and was expressed as R.S.D. (%) of a series of measurement. The result obtained shows R.S.D. equal to 0.83%, indicating good intra-day precision. Inter-day variability was calculated from assays on 3 days and shows a mean R.S.D. of 2.22% (Table 1). The accuracy of the method was determined

Parameters	LC method	UV method	VIS method
Useful concentration, μg/mL	6	6	6
Concentration range, μg/mL	5.0-15.0	7.0-12.0	8.0-13.0
Analytical curve	44299X+137924	0.0868X-0.0425	0.0255X-0.0257
Determination coefficient (r <sup>2</sup> )	0.9990	0.9995	0.9970
R.S.D. of repeatability (%)	0.08	0.83	0.73
R.S.D. intermediate precision (%)	1.23	2.22	0.88
Accuracy (%)	98.14	98.1	98.63
RSD of accuracy (%)	0.74	0.9	0.68
LOQ, μg/mL	2.89	0.73	3.82
LOD, μg/mL	0.95	0.24	1.26

Table 1: Validation parameters for chromatographic, visible and UV quantitation methods for azumolene.




Figure 3: Infrared spectrum of sodium azumolene sample (A) and standard (B) prepared in KBr.

by the mean recovery. The mean result was 98.10% (Table 1) indicating an agreement between the true value and the value found. The LOD and LOQ were calculated at the same way used for LC method. Their results were 0.24 and 0.73  $\mu\text{g/mL}$  respectively.

### Validation of VIS Spectrophotometric Method

At the same way of the UV method, the developed VIS method is fast and low cost. It shows to be linear at the concentration range tested (8.0 to 13.0  $\mu\text{g/mL}$ ). The equation of the curve was  $y=0.0255X-0.0257$  with a good coefficient of determination ( $r^2=0.9970$ ). The validity of the linearity test was evaluated by analysis of variance (ANOVA) which shows the statistical significance of the test.

The precision of the method was determined by repeatability (intra-day) and intermediate precision (inter-day) and was expressed as R.S.D. (%) of a series of measurement. The result obtained shows R.S.D. of 0.73% (Table 1). Inter-day variability was calculated from assays on 3 days and shows a mean R.S.D. of 0.88% (Table 1). The mean recovery found was equal to 98.63% (Table 1). The LOD and LOQ values were 1.26 and 3.82  $\mu\text{g/mL}$  respectively.

### Sodium azumolene characterization and assay testing–A method comparison

Sodium azumolene sample and standard were well characterized




Figure 4: UV absorption spectrum for sodium azumolene sample (A) and standard (B) prepared at 10.00  $\mu\text{g/mL}$  using purified water as diluter.




Figure 5: Overlay of visible spectrum (350-800 nm) of sodium azumolene sample and standard prepared at 10.0  $\mu\text{g/mL}$  using a 1.0% of chloranilic acid in acetonitrile as diluter.





Figure 6: TG and DTA curves for sodium azumolene sample (A) and standard (B).


by infrared spectroscopy and UV-VIS spectrophotometry. The infrared (IR) spectrum of sodium azumolene standard (A) and sample (B) are shown in Figure 3. The UV and VIS spectrum are shown in Figures 4 and 5.

The extinction coefficient was determined by measurement of the spectrophotometric absorbance at 339 nm of sodium azumolene aqueous solution. The result was equal to 0.246  $\text{M}^{-1}$ .

Moreover, sodium azumolene sample and standard were evaluated for its thermal behavior. A simultaneous differential scanning calorimetry (DSC) analysis and thermogravimetric (TG) analysis were made in atmosphere of oxygen and heating rate of 20°C/min. The results are shown in Figure 6.



**Figure 7:** Infrared spectrum of sodium dantrolene published by Japanese Pharmacopoeia (2001).



**Figure 8:** Ultraviolet spectrum of sodium dantrolene published by Japanese Pharmacopoeia (2001).

Method	Assay (%)	R.S.D. (%)
LC	98.80	0.85
UV	98.86	0.62
VIS	98.15	0.66

**Table 2:** Results of the determination of sodium azumolene applying the three validated methods.

Despite the similarity between the drugs sodium azumolene and sodium dantrolene it is possible to distinguish them by IR spectroscopy and UV espectrophotometry. The IR and UV spectra of sodium dantrolene were published by Japanese Pharmacopoeia and they are shown in Figures 7 and 8. It is quite different from sodium azumolene spectra.

The results for all sample tests (IR, UV, VIS and thermal) were consistent with the properties and concentrations of the sodium azumolene standards. In addition, sodium azumolene sample was evaluated using the three validated methods. The assay results from all methods were similar (Table 2).

All validated methods showed to be precise, accurate and linear at the concentration ranges tested. The chromatographic method is the most expensive method, since it requires special equipment and expensive solvents. The UV-VIS spectrophotometric methods are easier, more economical and faster than the LC method, however the LC-method is more precise and can readily identify interfering peaks.

## Conclusion

For routine quality control of medicines, it is essential to employ well-characterized, fully validated analytical methods to obtain reliable results that can be satisfactorily interpreted. The proposed methods were used successfully to investigate the sodium azumolene content. All methods are simple and efficient; and they were validated showing accurate and excellent linearity and precision characteristics.

Our study leads to conclude that the three validated method can be used to determine sodium azumolene and applied as quality control tools. The choice of the method to use should be made considering cost, simplicity, equipment, solvents, speed, and application to large or small workloads.

## Acknowledgments

The authors wish to thank the pharmaceutical company Cristalia (Brazil) for the supply of the raw material and thank the Fapesp, CNPq and PADC – UNESP to financial support.

## References

- Merli GJ, Bell RD (1987) Tratamiento preoperatorio del paciente quirúrgico con enfermedad neurológica. Consulta preoperatoria. *The Medical Clinics of North America* 71: 516-517.
- Rosenberg H, Seilman D, Fletcher J (1992) Certain inherited disorders enhanced or instigated by drug administered by anesthesiologist. *Clinical Anesthesia* 589-613.
- Allen GC (1993) Malignant hyperthermia and associated disorders. *Curr Opin Rheumatol* 5: 719-724.
- Denborough M (1998) Malignant hyperthermia. *Lancet* 352: 1131-1136.
- Krivosic-Horber RM, ADNET PJ (1992) L'hyperthermie maligne anaesthesique: un syndrome dont Le taux de mortalite est inacceptable. *La Revue Du Praticien Medecine Generate* 8: 150-188.
- Strazis KP, Fox AW (1993) Malignant hyperthermia: a review of published cases. *Anesth Analg* 77: 297-304.
- Manica J (2004) Anestesiologia: princípios e técnicas. 3rd edition, Artmed, Porto alegre.
- Britt BA (1984) Dantrolene. *Can Anaesth Soc J* 31: 61-75.
- Krause T, Gerbershagen MU, Fiege M, Weissshorn R, Wappler F (2004) Dantrolene—a review of its pharmacology, therapeutic use and new developments. *Anaesthesia* 59: 364-373.
- Kaplan RF (1991) Malignant hyperthermia. Refresher course lectures. *Asa. Annual Meeting*. San francisco.
- Torpy JM, Lynn C, Glass RM (2005) JAMA patient page. Malignant hyperthermia. *JAMA* 293: 2958.
- Sudo RT, Carmo PL, Trachez MM, Zapata-Sudo G (2008) Effects of azumolene on normal and malignant hyperthermia-susceptible skeletal muscle. *Basic Clin Pharmacol Toxicol* 102: 308-316.
- Campos Suarez JM, Escolano Villen F, Catala Puigbo E, Villar Landeira JM, Sanz Mazo M (1985) [Sodium dantrolene]. *Rev Esp Anestesiol Reanim* 32: 179-183.
- Schwe PF (1989) Azumolene dosage form. United States norwich eaton pharmaceuticals. Pat. n° 4822629. 12 dec. 1986, 18 apr. 1989. Norwich, NY.
- (2006) Azumolene inhibits a component of store-operated calcium entry coupled to the skeletal muscle ryanodine receptor. *J Biol Chem* 281: 33477-33486.
- Dhillon DS, Pong SF, Moorehead TJ (1992) A comparison of the potency and specificity of the direct muscle relaxant activity of azumolene and dantrolene. *Drug Development Research* 25: 161-169.
- Ben Abraham R, Adnet P, Glauber V, Perel A (1998) Malignant hyperthermia. *Postgrad Med J* 74: 11-17.
- Revathi R, Ethiraj T, Thenmozhi P, Saravanan VS, Ganesan V (2011) High performance liquid chromatographic method development for simultaneous analysis of doxofylline and montelukast sodium in a combined form. *Pharm Methods* 2: 223-228.
- Melby LR, Harder RJ, Hertler WR, Mahler W, Menson RE, et al. (1962) Substituted Quinodimethans. II. Anion-radical Derivatives and Complexes of 7,7,8,8-Tetracyanoquinodimethan. *J Am Chem Soc* 84: 3374-3387.
- Liming D, Qingqin X, Jianmei Y (2003) Fluorescence spectroscopy determination of fluoroquinolones by charge-transfer reaction. *J Pharm Biomed Anal* 33: 693-698.
- Saleh GA, Askal HF, Radwan MF, Omar MA (2001) Use of charge-transfer

- 
- complexation in the spectrophotometric analysis of certain cephalosporins. *Talanta* 54: 1205-1215.
22. Onah JO, Odeiani JE (2002) Simultaneous spectrophotometric determination of sulfadoxine and pyrimethamine in pharmaceutical formulations. *J Pharm Biomed Anal* 30: 851-857.
23. Khashaba PY, El-Shabouri SR, Emara KM, Mohamed AM (2000) Analysis of some antifungal drugs by spectrophotometric and spectrofluorimetric methods in different pharmaceutical dosage forms. *J Pharm Biomed Anal* 22: 363-376.
24. El-Ries MA, Abou Attia FM, Ibrahim SA (2000) AAS and spectrophotometric determination of propranolol HCl and metoprolol tartrate. *J Pharm Biomed Anal* 24: 179-187.
25. Faelelbom KMS (2008) Analysis of certain tetracyclines and oxytetracyclines through charge transfer complexation. *Am J Pharmacol Toxicol* 3: 212-218.
26. Japanese Pharmacopoeia (2001) 14<sup>th</sup> edition, Tokio, Society of Japanese Pharmacopoeia.