

Visceral Fat and Association with Metabolic Risk Factors

Leonardo S. Roever-Borges^{1*}, Resende ES¹

¹Department of Clinical Research, Federal University of Uberlândia, San Paulo, Brazil.

*Corresponding Author: Leonardo S. Roever-Borges, MHS Av. Pará, 1720 - Bairro Umuarama, Uberlândia - MG - CEP 38400-902. Tel: +553488039878; E-mail: leonardoroever@hotmail.com

Rec date: Feb 13, 2015; Acc date: Feb 16, 2015; Pub date: Feb 20, 2015

Copyright: © 2015 Roever. L et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Editorial

Cardiovascular disease (CVD) is the leading cause of morbidity and mortality in the world. Globally, more than one billion adults are overweight (BMI \geq 25 kg/m²). At least 300 million are obese (BMI \geq 30 kg/m²) [1]. The visceral fat (VF) is a risk factor for multiple CVD risk factors, including endothelial dysfunction, hypertension, dyslipidemia, diabetes, impaired glucose metabolism, insulin resistance, metabolic syndrome (MetS), liver insulin resistance, non-alcoholic fatty liver disease, sleep apnea, increased predisposition to cancers of the colon, breast and prostate, and it is associated with prolonged hospital stays, increased incidence of infections and non-infectious complications, and increased mortality in hospital [2-5]. Age, gender, genetics, and ethnicity are factors contributing to variation in visceral adipose tissue accumulation. Specific mechanisms responsible for proportionally increased visceral fat storage may involve sex hormones, local cortisol production in abdominal adipose tissues, dietary fructose endocannabinoids and growth hormone. Physiological characteristics of abdominal adipose tissues such as adipocyte size and number, lipolytic responsiveness, lipid storage capacity, and inflammatory cytokine production are significant correlates and even possible determinants of the increased cardiometabolic risk associated with visceral fat [6].

Abdominal obesity itself is an independent component of MetS and VF accumulation also determines a comprehensive cardiovascular risk profile and increases the susceptibility to ischaemic heart disease and arterial hypertension [7-8].

The VF releases different bioactive molecules and hormones, such as adiponectin, leptin, tumour necrosis factor (TNF- α), resistin and interleukin (IL-6, IL-1 β), resistin, adipocyte fatty acid binding protein (A-FABP), omentin-1, and chemerin. Adipokines contribute to the modulation of adipogenesis, immune cell migration into adipose tissue, adipocyte metabolism and function. Adiponectin is inversely correlated with the amount of VF, while decreased concentrations of adiponectin are associated with hypertension, type 2 diabetes, elevated glucose levels, cardiovascular disease and certain malignancies. Hypertriglyceridemia is a central correlate of visceral obesity. It is caused by a combination of increased liver VLDL triglyceride production and impaired clearance from the circulation [9-13].

An important contribution to evaluation of the influence of obesity on cardiovascular risk is the InterHeart Study, which shows evidence that abdominal obesity makes a higher contribution than BMI to the probability of these events [14]. Yusuf et al. defining the results more accurately, showing that the association between abdominal adiposity and coronary heart disease risk is highly significant in all geographical areas in which InterHeart Study data were collected [15].

In a population of 1,498 Caucasian, already showed that there was a strong independent association with VF and both cardiovascular (odd

ratio (95% CI):2.45 (1.52–3.95)) and cerebrovascular events (odd ratio (95% CI):1.63 (1.06–2.50)); in the same study, a (ROC) analysis proved greater sensitivity and specificity of VF, compared to its individual components (WC, BMI, HDL, and TG) with regard to cardiovascular and cerebrovascular events [16]. In a large case-control study, a high VF is associated with elevated risk of CHD in Chinese men and women [17]. As visceral obesity is associated with poor prognosis, metabolic disturbances and degree of pathology also occur in several chronic diseases. The VF is metabolically active and associated with insulin resistance, atherosclerosis, metabolic syndrome and CVD. To develop novel aetiology based strategies for the prevention and treatment of these diseases, a better understanding of the molecular mechanisms underlying obesity and its relationship to metabolic and cardiovascular diseases is essential.

Therefore, the identification of VF that regulates the atherosclerotic process might provide new opportunities for developing more effective approaches for preventing and treatment of cardiovascular disease.

References

1. Go A, Mozaffarian D, Roger VL, Benjamin EJ, Berry JD, et al. (2015) Heart Disease and Stroke Statistics—2015 Update A Report From the American Heart Association. *Circulation* 131: e29-e322.
2. Ritchie SA, Connell JM (2007) The link between abdominal obesity, metabolic syndrome and cardiovascular disease. *Nutr Metab Cardiovasc Dis* 17: 319-326.
3. Fox CS, Massaro JM, Hoffmann U, Pou KM, Maurovich-Horvat P, et al. (2007) Abdominal visceral and subcutaneous adipose tissue compartments: association with metabolic risk factors in the Framingham Heart Study. *Circulation* 116: 39-48.
4. Oh TH, Byeon JS, Myung SJ, Yang SK, Choi KS, et al. (2008) Visceral obesity as a risk factor for colorectal neoplasm. *J Gastroenterol Hepatol* 23: 411-417
5. Von H P, Pina F, Pérez A, Tavares M, Barros H (2004) Visceral fat accumulation as a risk factor for prostate cancer. *Obes Res* 12: 1930-1935
6. Tchernof A, Després JP (2013) Pathophysiology of human visceral obesity: an update. *Physiol Rev* 93(1): 359-404.
7. Mathieu P, Pibarot P, Larose E, Poirier P, Marette A, et al. (2008) Visceral obesity and the heart. *Int J Biochem Cell Biol* 40: 821-836.
8. Lamarche B, Lemieux S, Dagenais GR, Després JP (1998) Visceral obesity and the risk of ischaemic heart disease: insights from the Québec Cardiovascular Study. *Growth Horm IGF Res* 8 Suppl B: 1-8.
9. Shuster A, Patlas M, Pinthus JH, Mourtzakis M (2012) The clinical importance of visceral adiposity: a critical review of methods for visceral adipose tissue analysis. *Br J Radiol* 85(1009): 1-10.
10. Kanaya AM, Harris T, Goodpaster BH, Tylavsky F, Cummings SR (2004) Adipocytokines attenuate the association between visceral adiposity and diabetes in older adults. *Diabetes Care* 27: 1375-1380.
11. Yatagai T, Nagasaka S, Taniguchi A, Fukushima M, Nakamura T, et al. (2003) Hypoadiponectinemia is associated with visceral fat accumulation

-
- and insulin resistance in Japanese men with type 2 diabetes mellitus. *Metabolism* 52: 1274-1278.
12. Ebrahimi-Mamaeghani M, Mohammadi S, Arefhosseini SR, Fallah P, Bazi Z (2015) Adiponectin as a potential biomarker of vascular disease. *Vasc Health Risk Manag* 11: 55-70.
 13. Taskinen MR, Adiels M, Westerbacka J, Soderlund S, Kahri J, et al. (2011) Dual metabolic defects are required to produce hypertriglyceridemia in obese subjects. *Arterioscler Thromb Vasc Biol* 31: 2144-2150.
 14. Yusuf S, Hawken S, Ounpuu S (2004) Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study. *The Lancet* 364(9438)(937): 952.
 15. Yusuf S, Hawken S, Ôunpuu S, Bautista L, Franzosi MG, et al. (2005) Obesity and the risk of myocardial infarction in 27 000 participants from 52 countries: a case-control study. *The Lancet* 366(9497): 1640-1649.
 16. Amato MC, Giordano C, Galia M (2010) Visceral adiposity index: a reliable indicator of visceral fat function associated with cardiometabolic risk. *Diabetes Care* 33(4): 920-922.
 17. Zhang X, Shu XO, Li H (2013) Visceral adiposity and risk of coronary heart disease in relatively lean Chinese adults. *International Journal of Cardiology* 168(3): 2141-2145.