

Potential of Microbial Inoculated Water Hyacinth Amended Thermophilic Composting and Vermicomposting in Biodegradation of Agro-Industrial Waste

Alkesh Patidar*, Richa Gupta and Archana Tiwari

School of Biotechnology, Rajiv Gandhi Proudyogiki Vishwavidhyalaya, State Technological University of Madhya Pradesh, Airport Bypass Road, Bhopal, India

Abstract

Potential of *Streptomyces viridosporus*, *Aspergillus niger* and *Moraxella osloensis* combination in thermophilic composting followed vermicomposting of jatropha seed cake with 2:1 ratio of Water Hyacinth (WH) and Cow Dung (CD) was tested. Significant decrement in MC, TOC, and C: N ratio and increment in temperature, TP was observed. TKN, pH and TK were first increases in composting and then decreased in vermin composting. Most stable and matured vermin compost was obtained in S1 substrate in terms of C: N ratio below 20 (19.50 ± 1.10). Maximum earthworm number (47 ± 6), maximum net biomass worm⁻¹ (150 ± 15.6) obtained in S3 substrate and maximum cocoons (23 ± 2) were counted S1 substrate. 20% (w/w) vermin compost amended soil induced the growth in root length, shoot length and 100% GI while >20% (w/w) was inhibitory for plant growth. This approach decreases time period of degradation and produce good quality vermin compost if mixed up to 20%.

Keywords: Jatropha seed cake; Vermicompost; *Perionyx excavates*; Inoculation

Introduction

India has huge biomass of agro-industrial processing waste viz. food processing, pulp and paper, rice husk, cotton dust, jatropha seed cake, viticulture and winery waste, sugarcane trash, guar gum industrial waste, but the capability of these organic resources is not fully utilized, and disposal of these wastes through incineration, open dumping and land filling, raises serious human health risk and environmental issue [1]. For future biodiesel production from the non-food biodiesel crop in southern Asia and Africa, Jatropha curcas is cited as one of the best candidates [2]. But the processing of jatropha seed produce 0.68 million tons of seed cake (at the rate of 65% oil recovery with 3 tons/ha oil yield), which is quite a significant amount. Jatropha seed cake cannot be used as animal feed because of its toxic properties. It has been reported to possess phorbol esters, a potential toxic compound to animals. Toxicity of *J. curcas* seeds has been studied extensively in different animals like goat, sheep, mice, rats and fish. Decrease in glucose level, lack of appetite, diarrhoea, dehydration and other hemorrhagic effects in different organs were the common observations when animals were fed on phorbol ester containing feed [3-5]. Generation of huge amount of biomass also pose serious threats to environment.

Due to high nitrogen content, jatropha seed cake can be processed into organic fertilizers via microbial thermophilic composting which is a technique that has been used to address the issues of environmental pollution, non-reliance on chemical fertilizers, sustainable natural soil fertility, and minimizing the development of new dumps and landfills [6]. In recent times, interest in the use of a closely-related technique, known as vermin composting (using earthworms to breakdown organic materials) has increased [7]. Industrial wastes that are rich in organic matter and free from toxic substances or ions could be suitable substrates for vermin composting [8]. A combine approach of composting and subsequent vermicomposting brings the advantages of both process and minimizes the adverse impact of waste on the environment. The product obtained with desirable characteristics and at a faster rate than either of the individual processes [9]. Through composting the required temperature for ensuring satisfactory pathogen kill would be achieved and product passes EPA rules for pathogen reduction. Through vermin

composting, the rate of decomposition of organic materials increased and microbially active casts would be achieved.

Although, the microbial community naturally present in wastes usually carries out the process satisfactorily, the inoculation of residues with lignocellulolytic microorganisms is a strategy that could potentially enhance the way the process takes place or the properties of the final product. Inoculation with bacteria and fungi which can break down ligno-cellulolytic material has been reported to be effective in composting [10]. Inoculation increases cellulase activity, promote biodegradation of organic matter and accelerate the composting process [11,12] showed that mixed inoculation with complex microorganisms and ligno-cellulolytic microorganisms during composting of municipal solid waste had a clear advantage over inoculation with complex microorganisms or lingo-cellulolytic microorganisms alone.

Water hyacinth (*Eichhornia crassipes* (Mart) Solms – Laubach; family: Pontederiaceae) has been listed as most troublesome weed in aquatic systems. One of the easiest ways to utilize water hyacinth is to subject it to composting. Water hyacinth compost has been shown to have positive effect on plant growth. Research conducted by Umsakul et al. [13] showed that agro-industrial waste should be mixed with water hyacinth for composting to facilitate the development of thermophilic phase during the composting and to improve the product.

This present study was conducted to investigate the efficiency of water hyacinth and mixed microbial inoculation in biodegradation of industrial waste (Jatropha seed cake) during integrated composting

***Corresponding author:** Alkesh Patidar, School of Biotechnology, Rajiv Gandhi Proudyogiki Vishwavidhyalaya, State Technological University of Madhya Pradesh, Airport Bypass Road, Bhopal (M.P.), India, Tel: +918269537372; Fax: 0755-2742002-3; E-mail: pati.ak9@gmail.com

Received April 27, 2013; Accepted June 24, 2013; Published June 26, 2013

Citation: Patidar A, Gupta R, Tiwari A (2013) Potential of Microbial Inoculated Water Hyacinth Amended Thermophilic Composting and Vermicomposting in Biodegradation of Agro-Industrial Waste. J Bioremed Biodeg 4: 191. doi:10.4172/2155-6199.1000191

Copyright: © 2013 Patidar A, et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

and vermin composting technology. The objective of this study was to evaluate physical, chemical and biological parameters to determine the stability and maturity of composts prepared by mixing water hyacinth and cow dung in agro-industrial wastes.

Materials and Methods

Microbial source

The fungal strains *A. niger*, *S. viridosporus* and the bacterial strain *M. osloensis* were procured from IMTECH, Chandigarh, India. The fungi *A. Niger* and *S. viridosporus* were maintained by subculturing them on czepek yeast extract agar and trypton yeast extract respectively whereas bacteria was subcultured on tryptone yeast extract medium.

Industrial wastes (IW)

Jatropha seed cake (IW) was obtained from biodiesel plant at RGPV Bhopal.

Water hyacinth, cow dung and earthworm

Water Hyacinth (WH) was collected from the Lake situated in Bhopal. Cow manure (CD) was procured from a nearby poultry farm. Earthworms, *Perionyx excavatus*, used in this study came from the experimental farm of the Agronomic.

Experimental set-up

Different proportion of six feed mixture of industrial waste (Jatropha seed cake) was mixed in 2:1 ratio of CD and WH including one with CD and WH only were established (Table 1). Five kilogram of mixture (dry weight) was put in each rectangular pit (1 m length×1 m width×1 m height). A combination of pure cultures of *A. niger* and *S. viridosporus* (500 g mycelium per ton substrate) and *M. osloensis* (50 ml/kg substrate having 106 cells per ml) were inoculated in each pit on first day of composting. These mixtures were turned over manually every 24 h for 20 days in order to eliminate volatile substances potentially toxic to animals. The substrate with different treatments was pre-decomposed, in triplicates, for 20 days.

Subsequent vermicomposting

After 20 days, 20 adult individuals of *Perionyx excavatus* were introduced into each pit and then subjected to vermin composting for 20 days. The moisture content was maintained at $70 \pm 10\%$ of water holding capacity by periodic sprinkling of an adequate quantity of water. A plastic pipe with upper cap was installed in the middle of each bed and brought out of the ground through the tied mouth of the bag, in order to measure the temperature during composting. The temperature was measured daily by placing the thermometer through the plastic pipe. The earthworms and cocoons were removed manually and the vermin compost was chemically analyzed.

Compost sampling

Homogenized samples of the feed material were drawn at 0, 20 and

Substrate	Cow dung: Water hyacinth (2:1) (Kg)	Jatropha seed cake (Kg)
S1	4 (80%)	1 (20%)
S2	3 (60%)	2 (40%)
S3	2 (40%)	3 (60%)
S4	1 (20%)	4 (80%)
S5	0 (0%)	5 (100%)
Control	5 (100%)	0 (0%)

Table 1: Substrates composition.

40 days from each pit. The zero days refers to the time of initial mixing of the industrial wastes with cow dung and water hyacinth before preliminary decomposition.

Compost analysis

The pH of the compost was determined in distilled water with 1:10 (w/v) compost: water ratio. The moisture content was measured by drying to constant weight in an oven at 105°C. Changes in the temperature of the compost pile were recorded using mercury thermometer. The mean of the pile temperature at two monitoring points is recorded. Total Kjeldahl Nitrogen (TKN) and Total Organic Carbon (TOC) of the pre-decomposed bio inoculated residue and the vermicompost were measured with the Micro-Kjeldahl method [14] and Walkely and Black [15] Rapid titration method (1934). Total phosphorus (TP) was determined spectrophotometrically [16]. Total potassium (TK) was detected by the flame emission technique by Flame photometer [17].

Germination assay

Impact of most stable and matured vermin compost in terms of C: N ratio below 20 was tested on germinated seeds of cress plant. The amendments of vermin compost with garden soil were carried out at different concentrations (0, 20, 40, 60 80 and 100%). A control of garden soil was also used. Different concentrations of vermicompost amended soils (2.5 kg) were filled in different plastic pots. The soaked and sterile 30 cress (*L. sativum* L.) seeds were evenly sown in each pot to a depth of 2 inch. Prior to sowing, the cress seeds were sterilized in 3% formalin for 5 min to avoid any fungal contamination, washed thrice with double distilled water and soaked overnight in water. Experiments were conducted in triplicate with a parallel set of control. All the pots were watered daily till seed germination. All the pots were kept in a natural condition with an average durable temperature (25–35°C) and low humidity (50–55%). The germination index (GI) was calculated by the method reported by Fuentes et al. [18]. Subsequently, the plants were harvested after 5 days of sowing and repeatedly washed with tap water to remove any attached particles. Furthermore, these were rinsed with 10 mmol CaCl solution along with washing with deionised water. Root length, shoots length was measured manually using an inch-scale immediately after harvesting.

Statistical analysis

Results are the means of three replicates. One way analysis of variance (ANOVA) was done using the SIGMASTAT programme. The objective of statistical analysis was to determine any significant differences among the parameters analyzed in different treatments during the composting and vermin composting process.

Results and Discussion

Physico-chemical parameter

The physico-chemical characteristics of industrial waste, water hyacinth and cow dung are given in Table 2.

Changes in Temperature and Percent moisture content during composting are given in Figures 1 and 2. An initial temperature in the range of 26–28°C was noted in all five substrates and in control. Final temperature of 54°C in S2 and 54°C in control was recorded at 20th day of composting. It has been previously reported that the increase of temperature in the early phase of composting may be the result of thermophilic phase of composting. The temperature rise as a

Parameter	Jatropha seed cake	Water hyacinth	Cow dung
pH	6.8 ± 0.20	7.2 ± 0.20	8.7 ± 0.20
Total organic carbon (TOC)	35.64 ± 0.21	42.6 ± 0.10	39.2 ± 0.21
Total Kjeldahl nitrogen (TKN)	0.84 ± 0.03	0.86 ± 0.05	0.63 ± 0.02
C: N ratio	42.42 ± 1.12	49.3 ± 1.01	62.2 ± 1.05
Total phosphorus (TP)	0.50 ± 0.02	0.82 ± 0.13	0.59 ± 0.14
Total potassium (TK)	0.57 ± 0.10	0.82 ± 0.05	0.62 ± 0.12

Table 2: Physico-chemical characteristics of the initial raw materials used in experiment.




Figure 1: Changes in temperature during jatropha seed cake composting (20 days).




Figure 2: Changes in % moisture content during jatropha seed cake composting (20 days).

consequence of the rapid breakdown of organic matter, mostly readily available substrate rich in carbon, by microbial metabolism [19].

Moisture content is an important factor in degradation because it influences the structural and thermal properties of the material, as well as the rate of degradation and metabolic process of microorganisms. Initial moisture content was in the range of 81.15-75 on 5th day of composting. Final moisture content was decreased and in the range of 63.93-69.49 in all the treatments. Lowest percentage was noted in S3 treatment. After composting of jatropha seed cake pH was increased in all the treatments and after vermin composting on 40th day it was decreased (Figure 3). Increase in pH during composting during

composting was due to mineralization of the nitrogen and phosphorus into nitrites/nitrates and orthophosphates. The decrease in pH during vermicomposting may be due to CO₂ and organic acids produced by microbial metabolism [20].

Total Organic Carbon (TOC) decreased significantly in all treatments during composting and subsequent vermin composting, extremely in S1 and S2 which contained lower concentrations of industrial wastes. It may be due to the mineralization of organic matter [21]. During composting maximum carbon loss of 11.84% was noted in S1 treatment and least was 1.98% in S5 treatment (Table 3). This decrement was due to combined potential of added microbes and water hyacinth both initiate the thermophilic phase of composting. During vermin composting maximum carbon loss of 7.9% was noted in S2 and least was 1.92 in S3. It was the action of Earthworms which break and homogenizes the ingested material through muscular action of their foregut and, also adds mucus and enzymes in ingested material, this increase the surface area for microbial action. Thus action of water hyacinth earthworms and microorganisms bring TOC loss from the substrates in the form of CO₂ [22].

Percentage loss in carbon during composting was fast as compared to that of the vermin composting and it was due to higher initial N concentration, which might have increased the microbial activity in composting phase [23].

During composting phase significant increment in TKN at 20th day was noted. Maximum percentage of TKN increment was noted in S1 treatment which receives the 80% of 2:1 ratio of cow dung and water hyacinth (Table 3). This increment might be due to potential of water hyacinth which facilitate and initiate the development of thermophilic phase during the composting process [13] and the inoculated microbes especially *moraxella osloensis* which fix the nitrogen present in the substrate. During subsequent vermin composting nitrogen content was decreased in all the treatments possibly due to ammonification, NH₃ volatilization and denitrification as reported by Bernal et al. [24].

C: N ratio-index of compost maturity

The C: N ratio of a substrate material reflects the organic waste mineralization and stabilization during the process of composting and vermin composting. The loss of carbon as CO₂ through microbial respiration and simultaneously addition of nitrogen by worms in the form of Mucus and nitrogenous excretory material lower the C: N ratio of the substrate [25]. C: N ratio is considered as a parameter to




Figure 3: Changes in pH during jatropha seed cake composting (20 days) and subsequent vermicomposting (20 days).

establish the degree of maturity of compost and its agronomic quality [26]. Decline of C: N ratio to less than 20 indicates an advanced degree of organic matter stabilization and reflects a satisfactory degree of maturity of organic wastes [27]. Microbial composting and vermin composting of jatropa seed cake resulted in a loss of carbon because of mineralization of organic waste (Figure 4). Initial C: N ratio was in the range of 37.89-65.47 at zero days in different substrates. The initial C: N ratio was more in those feed mixtures which had higher percentage of jatropa seed cake. After composting and vermincompostubg C: N ratio was in the range of 18.59-37.82 and 18.46-39.11 respectively. Maximum decrement in C: N ratio was observed in S1 substrate comparable to control (P<0.001). The decomposition of the waste during vermicomposting was slow compared to that of the composting

and might have been due to higher initial N concentration, which might have increased the microbial activity in the beginning, thus decreasing the C/N ratio [23].

Percentage of total phosphorus was increased significantly during composting and vermicomposting in all the treatments with respect to control (P<0.001). The overall percent increase in the TP content was maximum in the substrate S3 (1.71 ± 0.10), and minimum percent increase was in the substrate S5 (0.65 ± 0.20) (Table 4). Yadav and Garg [28] noticed 1.5-2-fold increase in total phosphorus from industrial sludge vermicomposting. Lee [29] states that when organic material passes through the gut of earthworm then insoluable phosphorus is converted to soluable form by phosphatase produced within the

Substrate	0 days		20 days		40 days	
	TOC	TKN	TOC	TKN	TOC	TKN
Control	36.00 ± 0.12*	0.95 ± 0.21	27.74 ± 0.01*	1.49 ± 0.10*	25.66 ± 0.20*	1.39 ± 0.01*
S1	45.50 ± 0.21*	0.92 ± 0.01	33.66 ± 0.15*	1.57 ± 0.12	27.70 ± 0.15*	1.42 ± 0.02
S2	53.40 ± 0.25*	0.93 ± 0.03	49.40 ± 0.18*	1.45 ± 0.10	41.50 ± 0.01*	1.42 ± 0.01
S3	55.00 ± 0.10*	0.84 ± 0.02	53.40 ± 0.20*	1.48 ± 0.10	51.48 ± 0.01*	1.36 ± 0.10
S4	47.52 ± 0.10*	0.86 ± 0.01	45.50 ± 0.05*	1.42 ± 0.20	43.56 ± 0.05*	1.32 ± 0.01
S5	35.64 ± 0.21	0.84 ± 0.03	33.66 ± 0.31*	0.89 ± 0.10*	31.68 ± 0.24*	0.81 ± 0.02*

All values are mean (n=3) ± S.D. and given in percentage.
*Significant (P<0.001).

Table 3: Percentage of total organic carbon (TOC), total Kjeldahl nitrogen (TKN) in different substrate composition during microbial composting (0–20 days) and subsequent vermicomposting (20 days).




Figure 4: Changes in C: N Ratio during jatropa seed cake composting (20 days) and subsequent vermicomposting (20 days).

Substrate	0 days	20 days	40 days
Control	37.89 ± 1.10*	18.59 ± 1.20*	18.46 ± 1.20*
S1	49.45 ± 1.10*	21.43 ± 1.02*	19.50 ± 1.10
S2	57.41 ± 1.20*	34.06 ± 1.12*	29.22 ± 1.20*
S3	65.47 ± 1.21*	36.08 ± 1.02*	37.85 ± 1.15*
S4	55.25 ± 1.02*	32.04 ± 1.12*	33.00 ± 1.10*
S5	42.42 ± 1.12*	37.82 ± 1.10*	39.11 ± 1.10*

All values are mean (n=3) ± S.D. and given in percentage.
*Significant (P<0.001).

Table 4: Changes in C/N ratio during composting and vermicomposting.

earthworm and secreted by inoculated microbes which is easily available to plants.

Data in Table 4 reported that total potassium during microbial composting increase significantly in all the treatments as well as in control ($P < 0.001$). Maximum increment was noted in S3 substrate (1.57 ± 0.15) and minimum was in S1 substrate (0.72 ± 0.10). However, we observed a marginal decrease in potassium during vermicomposting. Only S1 substrate showed slight increment in TK. This decrement in potassium was supported the result of Kumar et al. [30] who reported lower K content in sugar-cane waste by-products after vermicomposting. There are contradictory reports regarding the potassium content in vermicompost obtained from different substrates. Hait and Tare [31] have reported higher potassium content in the sewage sludge vermicompost.

This difference in potassium content can be attributed to the differences in the chemical nature of substrate. This might be also due to leaching of potassium by excess water that drained through the feeds. Benitez et al. [32] have reported that the leachate collected during vermicomposting process had higher potassium concentrations. It has been suggested that earthworm processed residue contains higher concentration of exchangeable K due to enhanced microbial activity during vermicomposting, which consequently enhances the rate of mineralization [25].

Total no. of earthworm, cocoons and biomass

The different pattern of earthworm number and cocoons production depending on the percentage of substrate in the feed mixtures was observed after vermicomposting of jatropa seed cake (Table 5). Maximum earthworm number was observed in substrate S3 and maximum cocoons production was noted in S1. The addition of industrial waste up to 40% in cow dung and water hyacinth was found significant in cocoon production, but concentration above this level was not able to support the cocoon production. The increase in earthworms' growth may also be attributed to a low C: N ratio of the microbial composted easily consumable water hyacinth added substrate [8]. Fayolle et al. [33] suggested that food source play an important role

on cocoon production rates by worms.

The maximum net biomass production by *Perionyx excavatus* was observed in substrate S2 (150 ± 15.6 mg earthworm⁻¹) and lowest in S5 substrate (53 ± 15.2 mg earthworm⁻¹). Increasing percentage of jatropa seed cake in the feed mixtures resulted in a decrease in biomass of earthworm. It is concluded from the results that addition of 40% jatropa seed cake to the water hyacinth and cow dung is acceptable during the vermicomposting in terms of growth and reproductive success of the earthworms.

Biological parameter of vermicompost analysis

The evaluation of industrial wastes toxicity by chemical characterisation and biological testing is extremely important for screening the suitability of vermicompost for land application. To evaluate the biological parameter of vermicompost, the amendment of most stable and matured compost on the basis of C: N ratio less than 20, substrate S1 with garden soil were carried out at different concentrations (20, 40, 60, 80 and 100%). In general, percentage of seed germination decreased with increase in vermicompost concentrations from S1a to S1e (Table 6). In present study, it was observed that up to 20% (w/w) concentration of vermicompost has no inhibitory effect on seed germination while at higher concentration (>40%), decrease in percent germination was recorded. This decrease in percent germination was due to phytotoxic effect produced by combination of several factors, rather than one [34]. These factors include the presence of heavy metals, ammonia, salts, and low molecular weight organic acids [18] all of which have been shown to have inhibitory effects.

The vegetative growth parameters of cress plant (shoot length and root length) at all tested concentrations showed significant inhibitory effect (except 20% vermicompost amended soil) versus the control (Table 7). Maximum root length was noted in S1a trial (3.6 ± 0.12). Similarly maximum shoot length was observed in S1a trial (5.6 ± 0.14) more than the control. As the concentration of vermicompost increased root and shoot length was decreased. It has been reported that high vermicompost content was suppressive for plant growth

Substrate	0 days		20 days		40 days	
	TP	TK	TP	TK	TP	TK
Control	$0.85 \pm 0.10^*$	$0.90 \pm 0.20^*$	0.98 ± 0.10	$1.70 \pm 0.15^*$	1.25 ± 0.10	$1.67 \pm 0.20^*$
S1	0.70 ± 0.10	$0.42 \pm 0.18^*$	0.88 ± 0.10	$0.72 \pm 0.10^*$	1.65 ± 0.10	$0.78 \pm 0.10^*$
S2	0.67 ± 0.01	0.80 ± 0.15	0.90 ± 0.10	$1.57 \pm 0.15^*$	1.70 ± 0.20	1.47 ± 0.25
S3	0.66 ± 0.02	0.77 ± 0.15	0.93 ± 0.20	1.65 ± 0.20	1.71 ± 0.10	1.57 ± 0.15
S4	$0.61 \pm 0.10^*$	0.70 ± 0.10	0.67 ± 0.15	1.35 ± 0.10	0.83 ± 0.15	1.46 ± 0.15
S5	$0.50 \pm 0.02^*$	0.60 ± 0.10	0.52 ± 0.20	$1.10 \pm 0.20^*$	$0.65 \pm 0.20^*$	$1.17 \pm 0.16^*$

All values are mean ($n=3$) \pm S.D. and given in percentage.

*Significant ($P < 0.001$).

Table 5: Percentage of total phosphorus (TP) and total potassium (TK).

Substrate	Initial No. of earthworms	Total earthworm	Cocoons no.	Mean Biomass worm ⁻¹ (mg)		Net biomass worm ⁻¹ (mg)
				Initial	Final	
Control	20	$31 \pm 4^*$	$11 \pm 4^*$	$235 \pm 2.5^*$	$375 \pm 12^*$	$140 \pm 11.2^*$
S1	20	28 ± 5	$23 \pm 2^*$	238 ± 1.7	368 ± 12	130 ± 5.6
S2	20	$42 \pm 4^*$	18 ± 4	240 ± 3.6	390 ± 14	150 ± 15.6
S3	20	$47 \pm 6^*$	07 ± 1	230 ± 1.7	$325 \pm 11^*$	$95 \pm 2.4^*$
S4	20	31 ± 2	04 ± 3	230 ± 1.6	$284 \pm 4.3^*$	$54 \pm 4.9^*$
S5	20	29 ± 5	03 ± 2	$225 \pm 1.7^*$	$278 \pm 11^*$	$53 \pm 15.2^*$

All values are mean ($n=3$) \pm S.D.

*Significant ($P < 0.001$).

Table 6: Changes in growth, cocoon production and net biomass of earthworms.

Vermicompost concentration (%)	GI (%)	Root length (cm)	Shoot length (cm)
Control (0)	100	3.6 ± 0.30	4.2 ± 0.10*
S1a (20)	100	3.6 ± 0.12	5.6 ± 0.14*
S1b (40)	96	3.2 ± 0.21	5.4 ± 0.13*
S1c (60)	70	3.2 ± 0.20	4.3 ± 0.10*
S1d (80)	70	3.0 ± 0.10	3.8 ± 0.12*
S1e (100)	60	2.6 ± 0.21*	3.2 ± 0.10*

All values are mean (n=3) ± S.D.

*Significant (P<0.001).

Table 7: Effect of vermicompost S1-amended soil on germination of cress seed.

hormone (s) (auxin and gibberline) which are responsible for the growth and development of plants [35]. The reduction in plant growth at high concentrations of vermicompost might be due to the entrance of the metal into the protoplasm resulting in the loss of intermediary metabolites which are essential for further growth and development of plants. These findings are very much in accordance with the earlier reported results [36].

Conclusion

The chemical analyses of the compost produced by microbial inoculated composting and vermicomposting of jatropa seed cake admixed with water hyacinth and cow dung, point towards its patentability. On the basis of C: N ratio below 20, earthworm biomass, growth and reproduction, results demonstrated that after the addition of jatropa seed cake (20%) in 2:1 ratio of water hyacinth, cow dung, and inoculated microbes, *S. viridosporus*, *A. niger* and *M. osolensis*, it can be used as a raw material. Germination index using cress seeds showed that concentration above 20% of vermicompost had inhibitory effect and cause phytotoxicity.

Acknowledgements

Author would like to thank the Indian institute of soil science for help with the compost analyzing.

References

- Raj D, Antil RS (2011) Evaluation of maturity and stability parameters of composts prepared from agro-industrial wastes. *Bioresour Technol* 102: 2868-2873.
- Sachs G (2010) Commodity prices and volatility: old answers to new questions. Goldman Sachs Global Economics, Commodities and Strategy Research, Global Economics Paper No: 194.
- Adam SE (1974) Toxic effects of *Jatropha curcas* in mice. *Toxicology* 2: 67-76.
- Adam SE, Magzoub M (1975) Toxicity of *Jatropha curcas* for goats. *Toxicology* 4: 347-354.
- Makkar HP, Becker K (1999) Nutritional studies on rats and fish (carp *Cyprinus carpio*) fed diets containing unheated and heated *Jatropha curcas* meal of a non-toxic provenance. *Plant Foods Hum Nutr* 53: 183-192.
- Papadimitriou EK, Balis C (1996) Comparative study of parameters to evaluate and monitor the rate of a composting process. *Compost Sci Utilization* 4: 52-61.
- Logsdon G (1994) Worldwide progress in vermicomposting. *Biocycle* 35: 63-65.
- Ndegwa PM, Thompson SA (2000) Effects of C-to-N ratio on vermicomposting of biosolids. *Bioresour Technol* 75: 7-12.
- Ndegwa PM, Thompson SA (2001) Integrating composting and vermicomposting in the treatment and bioconversion of biosolids. *Bioresour Technol* 76: 107-112.
- Nair J, Okamitsu K (2010) Microbial inoculants for small scale composting of putrescible kitchen wastes. *Waste Manag* 30: 977-982.
- Ghaffari S, Sepahi AA, Razavi MR, Malekzadeh F, Haydarian H (2011) Effectiveness of inoculation with isolated *Anoxybacillus* sp MGA110 on municipal solid waste composting process. *Afr J Microbiol Res* 5: 5373-5378.
- Wei Z, Xi B, Zhao Y, Wang S, Liu H, et al. (2007) Effect of inoculating microbes in municipal solid waste composting on characteristics of humic acid. *Chemosphere* 68: 368-374.
- Umsakul K, Dissara Y, Srimuang N (2010) Chemical, physical and microbiological changes during composting of the water hyacinth. *Pak J Biol Sci* 13: 985-992.
- Shaw J, Beadle LC (1949) A simplified ultramicro Kjeldahl method for the estimation of protein and total nitrogen in fluid samples of less than 1-0 µl. *J Exp Biol* 26: 15-23.
- Walkely JA, Black JA (1934) Estimation of organic carbon by the chronic acid titration method. *Soil Sci* 37: 29-31.
- Fiske CH, Subbarow Y (1925) The colorimetric determination of phosphorus. *J Biol Chem* 66: 375-400.
- Pearson RW (1952) Potassium – supplying power of eight Alabama Soils. *Soil Sci* 74: 301-310.
- Fuentes A, Lloréns M, Sáez J, Aguilar MI, Ortuño JF, et al. (2004) Phytotoxicity and heavy metals speciation of stabilised sewage sludges. *J Hazard Mater* 108: 161-169.
- Kutzner HJ (2000) Microbiology of composting technology. In: Rehm HJ, Reed G (eds) *Biotechnology Environmental Processes II*, John Wiley & Sons, USA.
- Elvira C, Sampedro L, Benítez E, Nogales R (1998) Vermicomposting of sludges from paper mill and dairy industries with *Eisenia andrei*: A pilot-scale study. *Bioresour Technol* 63: 205-211.
- Kaushik P, Garg VK (2003) Vermicomposting of mixed solid textile mill sludge and cow dung with the epigeic earthworm *Eisenia foetida*. *Bioresour Technol* 90: 311-316.
- Edwards CA (1988) Breakdown of animal, vegetable and industrial organic wastes by earthworms. *Earthworms in Waste and in Environment management*. SPB Academic Publishing, The Netherlands.
- Eiland F, Klamer M, Lind AM, Leth M, Bååth E (2001) Influence of Initial C/N Ratio on Chemical and Microbial Composition during Long Term Composting of Straw. *Microb Ecol* 41: 272-280.
- Bernal MP, Navarro AF, Roig A, Cegarra J, Garcia D (2006) Carbon and nitrogen transformation during composting of sorghum bagasse. *Biol Fertil Soils* 22: 141-148.
- Suthar S (2008) Bioconversion of post harvest crop residues and cattle shed manure into value-added products using earthworm *Eudrilus eugeniae* Kinberg. *Ecological Engineering* 32: 206-214.
- Singh D, Suthar S (2012) Vermicomposting of herbal pharmaceutical industry solid wastes. *Ecological Engineering* 39: 1-6.
- Senesi N (1989) Composted materials as organic fertilizers. *Sci Total Environ* 81/82: 521-524.
- Yadav A, Garg VK (2009) Feasibility of nutrient recovery from industrial sludge by vermicomposting technology. *J Hazard Mater* 168: 262-268.
- Lee KE (1992) Some trends and opportunities in earthworm research or: Darwin's children--the future of our discipline. *Soil Biol Biochem* 24: 1765-1771.
- Kumar R, Verma D, Singh BL, Kumar U, Shweta (2010) Composting of sugar-cane waste by-products through treatment with microorganisms and subsequent vermicomposting. *Bioresour Technol* 101: 6707-6711.
- Hait S, Tare V (2011) Vermistabilization of primary sewage sludge. *Bioresour Technol* 102: 2812-2820.
- Benitez E, Nogales R, Elvira C, Masciandaro G, Ceccanti B (1999) Enzyme activities as indicators of the stabilization of sewage sludge composting with *Eisenia foetida*. *Bioresour Technol* 68: 297-303.
- Fayolle L, Michaud H, Cluzeau D, Stawiecki J (1997) Influence of temperature and food source on the life cycle of the earthworm *Dendrobaena veneta* (*Oligochaeta*). *Soil Biol Biochem* 29: 747-750.
- Hoekstra NJ, Bosker T, Lantinga EA (2002) Effects of cattle dung from farms with different feeding strategies on germination and initial root growth of cress (*Lepidium sativum* L.). *Agric Ecosys Environ* 93: 189-196.

35. Misra RN, Behera PK (1991) The effect of paper industry effluent on growth, pigments, carbohydrates and proteins of rice seedlings. *Environ Pollut* 72: 159-167.
36. Chandra R, Yadav S, Mohan D (2008) Effect of distillery sludge on seed germination and growth parameters of green gram (*Phaseolus mungo* L.). *J Hazard Mater* 152: 431-439.