

Promoting Awareness on Cerebrovascular Disease in Young Adults is a Work in Progress

Aleksandra Pikula*

Department of Neurology, Boston University School of Medicine, Boston, MA 02118, USA

Over the past decades, the overall stroke incidence has been declining, but among young adults the stroke incidence is increasing. Population and hospital-based studies have raised concerns about the increasing prevalence of traditional stroke risk factors among young adults and its consequence on stroke incidence, stroke recurrence and stroke related morbidity and mortality [1,2].

Stroke events in individuals younger than 50 years of age are considered "strokes in young adults". Based on the US Nationwide Inpatient Sample data from 1993-2008, hospital admissions among patients 14-44 years old with diagnosis of ischemic stroke have increased by ~50%. The same data show concurrent increase in prevalence of traditional stroke risk factors among young for obesity, dyslipidemia, hypertension and smoking [3,4]. Due to popular belief that stroke is a disease of elderly, stroke symptoms among young are commonly misdiagnosed in the acute settings. The young stroke patients are also less likely to use pre-hospital stroke pathway systems [5]. However, the severity of ischemic stroke is milder [6], and the prognosis after stroke in young adults is more favorable than in the elderly [7]. Nevertheless, the long-term psychosocial consequences are greater and the risk for recurrence due to longer lifetime expectancy is much higher [1,8-10].

With introduction of sensitive brain imaging techniques it had been established that the burden of cerebrovascular disease is far greater than that of overt clinical disease [11]. In large population-based studies, the MRI findings of subclinical vascular brain injury such as presence of covert brain infarcts (CBIs), white matter hyperintensities (WMHs) and cerebral microbleeds (CMBs) are common. The prevalence of CBIs and WMHs ranges from 10-20% in subjects <65 years of age, and frequency of CMBs while limited by number of studies available was 6.5-8% in <50 years old otherwise healthy subjects [12,11]. While these subclinical findings do not cause abrupt clinical symptoms, they are often linked to cognitive decline and motor function impairment, and have been associated with an increased risk of subsequent stroke and dementia [11]. The complexity of MRI techniques and inconsistency in the definitions of these MRI measurements limits systematic examination of their frequencies and its relevance on the functional and cognitive outcomes in young adults with ischemic stroke. As a result, there are only a few studies that evaluated this in some detail.

Utilizing quantitative scores that are based on presence or absence of SBIs and on the degree of WMHs (mild, moderate or severe) observed on brain MRI, the Helsinki Stroke Registry retrospectively examined 669 predominantly white patients with first-ever ischemic stroke. SBIs were present in 13% and WMHs in 7% of all patients. Most patients had mild to moderate degree of WMHs. The risk factors for higher prevalence of SBIs and WMHs were older age, type 1 DM and obesity, but not hypertension as usually observed in middle aged or elderly population. In multivariate analysis, the presence of SBIs independently increased the risk of recurrent stroke, and moderate to severe WMHs were independently associated with a greater risk of death (OR=3.43) [13].

From the single-center French retrospective study, among 170 adults (18-50 years old) with first-ever ischemic stroke, 28% patients had

silent ischemic lesions (SILs). Over the mean follow-up of ~25 months, 23% of patients with SILs had a recurrent stroke vs. 6.5% without SILs, $p=0.003$. After adjusting for vascular risk factors, recurrent stroke was associated with presence of SILs (HR 3.2, $p=0.002$), and more so with the presence of combination of SBIs and WMHs (HR 7.3, $p=0.003$) [14].

Data published in this month's issue of Neurology arrives from the largest prospective study of stroke in young adults with Fabry disease (SIFAP1 study) that enrolled young adults with first ever TIA or stroke aged 18-55 years (over 2000 subjects with ischemic stroke and TIA). They observed equally high prevalence of WMH (46% of all) with the severity positively associated with older age (>45 years). The old infarcts were present in 8% of patients younger than 35 years, while in the age groups 35-44 and older than 44, the prevalence of old infarcts was 12% and 25%, respectively.

Intriguingly, this rate was double the one of SBIs prevalence among young adults with first-ever ischemic stroke observed in Helsinki study. The prevalence of traditional risk factors was high and the prevalence of Fabry disease in the same study as reported earlier was 0.5% [15].

Interestingly, Fabry disease frequently presents with clinical symptoms and MRI lesions that are often misdiagnosed as multiple sclerosis [16]. In this study, the prevalence of WMH among patients diagnosed with Fabry disease was unclear. There was also no discussion on how this could influence the overall study findings. It is worth underlining that SBIs and WMHs are frequently seen in patients with other more common conditions such as migraine or multiple sclerosis, and/or with less common but other determined stroke etiologies such as CADASIL (cerebral autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy), MELAS (mitochondrial encephalomyopathy, lactic acidosis, and stroke-like episodes), lupus, various encephalopathies and vasculitides. So, when interpreting data from this and all of the mentioned studies, it is important to notice that patients with these conditions should have been accounted for or excluded from the analysis.

If we recognize these shortcomings, along with the differences in the study populations and designs, these reports still provide an intriguing observation of strikingly high prevalence of so-called silent

***Corresponding author:** Aleksandra Pikula, Assistant Professor, Department of Neurology, Boston University School of Medicine, 715 Albany Street, D-312, Boston, MA 02118-2526, USA, Tel: (617)-638-5435; Fax: (617) 638-8465; E-mail: alpkikula@bu.edu

Received November 28, 2013; **Accepted** November 30, 2013; **Published** December 07, 2013

Citation: Pikula A (2013) Promoting Awareness on Cerebrovascular Disease in Young Adults is a Work in Progress. J Neurol Neurophysiol 5: e115. doi:[10.4172/2155-9562.1000e115](http://dx.doi.org/10.4172/2155-9562.1000e115)

Copyright: © 2013 Pikula A. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

brain infarcts and WMHs among young adults with ischemic stroke. Given that presence of WMHs on MRI are associated with an increased risk of stroke in the general population and in high risk populations with a history of stroke [11], once documented during diagnostic investigations it should prompt extensive exploration of possible etiologies as well as aggressive treatment of concurrent risk factors.

Consequently, it is not only the increased incidence of stroke and stroke risk factors that we need to take into account, but also the high prevalence of covert cerebrovascular disease in young adults. Further research should focus on all aspects of overt and covert cerebrovascular disease, and on rising public awareness and providing better access to primary and secondary stroke preventive care for young adults.

References

1. Putaala J, Curtze S, Hiltunen S, Tolppanen H, Kaste M, et al. (2009) Causes of death and predictors of 5-year mortality in young adults after first-ever ischemic stroke: the Helsinki Young Stroke Registry. *Stroke* 40: 2698-2703.
2. Putaala J, Metso AJ, Metso TM, Konkola N, Kraemer Y, et al. (2009) Analysis of 1008 consecutive patients aged 15 to 49 with first-ever ischemic stroke: The helsinki young stroke registry. *Stroke; a journal of cerebral circulation.* 40:1195-1203.
3. George MG, Tong X, Kuklina EV, Labarthe DR (2011) Trends in stroke hospitalizations and associated risk factors among children and young adults, 1995-2008. *Ann Neurol* 70: 713-721.
4. Kissela BM, Khoury JC, Alwell K, Moomaw CJ, Woo D, et al. (2012) Age at stroke: temporal trends in stroke incidence in a large, biracial population. *Neurology* 79: 1781-1787.
5. Kuruvilla A, Bhattacharya P, Rajamani K, Chaturvedi S (2011) Factors associated with misdiagnosis of acute stroke in young adults. *Journal of stroke and cerebrovascular diseases: the official journal of National Stroke Association* 20: 523-527.
6. Toni D, Ahmed N, Anzini A, Lorenzano S, Brozman M, et al. (2012) Intravenous thrombolysis in young stroke patients: results from the SITS-ISTR. *Neurology* 78: 880-887.
7. Varona JF, Bermejo F, Guerra JM, Molina JA (2004) Long-term prognosis of ischemic stroke in young adults. Study of 272 cases. *J Neurol* 251: 1507-1514.
8. Rutten-Jacobs LC, Maaijwee NA, Arntz RM, Schoonderwaldt HC, Dorresteijn LD, et al. (2013) Long-term risk of recurrent vascular events after young stroke: The FUTURE study. *Ann Neurol* 74: 592-601.
9. Schaapsmeeders P, Maaijwee NA, van Dijk EJ, Rutten-Jacobs LC, Arntz RM, et al. (2013) Long-term cognitive impairment after first-ever ischemic stroke in young adults. *Stroke* 44: 1621-1628.
10. Rutten-Jacobs LC, Arntz RM, Maaijwee NA, Schoonderwaldt HC, Dorresteijn LD, et al. (2013) Long-term mortality after stroke among adults aged 18 to 50 years. *JAMA* 309: 1136-1144.
11. Debette S, Markus HS (2010). The clinical importance of white matter hyperintensities on brain magnetic resonance imaging: Systematic review and meta-analysis. *Bmj* 341: c3666.
12. Shoamanesh A, Kwok CS, Benavente O (2011) Cerebral microbleeds: histopathological correlation of neuroimaging. *Cerebrovasc Dis* 32: 528-534.
13. Putaala J, Kurkinen M, Tarvos V, Salonen O, Kaste M, et al. (2009) Silent brain infarcts and leukoariosis in young adults with first-ever ischemic stroke. *Neurology* 72: 1823-1829.
14. Gioia LC, Tollard É, Dubuc V, Lanthier S, Deschaintre Y, et al. (2012) Silent ischemic lesions in young adults with first stroke are associated with recurrent stroke. *Neurology* 79: 1208-1214.
15. Rolfs A, Fazekas F, Grittner U, Dichgans M, Martus P, et al. (2013) Acute cerebrovascular disease in the young: the Stroke in Young Fabry Patients study. *Stroke* 44: 340-349.
16. Böttcher T, Rolfs A, Tanislav C, Bitsch A, Köhler W, et al. (2013) Fabry disease - underestimated in the differential diagnosis of multiple sclerosis? *PLoS One* 8: e71894.

Citation: Pikula A (2013) Promoting Awareness on Cerebrovascular Disease in Young Adults is a Work in Progress. *J Neurol Neurophysiol* 5: e115.
doi:[10.4172/2155-9562.1000e115](https://doi.org/10.4172/2155-9562.1000e115)

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 300 Open Access Journals
- 25,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission>