

Approximate Solution of Second-Order Linear Differential Equation

Nadhem Echi*

Department of Mathematics, El Manar University, Tunis 1002, Tunisia

Abstract

This paper presents an efficient approach for determining the solution of second-order linear differential equation. The second-order linear ordinary differential equation is first converted to a Volterra integral equation. By solving the resulting Volterra equation by means of Taylor's expansion, different approaches based on differentiation and integration methods are employed to reduce the resulting integral equation to a system of linear equation for the unknown and its derivatives the approximate solution of second-order linear differential equation is obtained. Test example demonstrates the effectiveness of the method and gives the efficiency and high accuracy of the proposed method.

Mathematics subject classification: 34-XX; 41-XX

Keywords: Second-order linear differential equation; Volterra integral equation; Taylor's expansion; Error analysis

Introduction

Some second-order differential equations with variable coefficients can be solved analytically by various methods [1]. For general cases, one must appeal to numerical techniques or approximate approaches for getting its solutions [2,3]. The Adomian decomposition method for solving differential and integral equations, linear or nonlinear, has been subject of extensive analytical and numerical studies. In particular Adomian's decomposition method has been proposed for solving second-order linear differential equation.

In the present paper, we give an approach for determining the solution of second-order linear differential equation. The second-order linear ordinary differential equation is first converted to a Volterra integral equation. By solving the resulting Volterra equation by means of Taylor's expansion, different approaches based on differentiation and integration methods are employed to reduce the resulting integral equation to a system of linear equation for the unknown and its derivatives the approximate solution of second-order linear differential equation is obtained. By studying the estimation of the error give the efficiency and high accuracy of the proposed method [4-6].

Volterra Integrals Equations

We consider the following second-order differential equation :

$$(E) : y''(t) + p(t)y'(t) + q(t)y(t) = g(t) \tag{1}$$

with p, q and g are infinitely differential functions in open interval $I \subset \mathbb{R}$. We fix a point a of the interval I . We have, $\forall x \in I$,

$$\int_a^x \frac{(t-x)^{i+1}}{(i+1)!} y''(t) dt = y(a) \frac{(a-x)^i}{i!} - y'(a) \frac{(a-x)^{i+1}}{(i+1)!} + \int_a^x \frac{(t-x)^{i-1}}{(i-1)!} y(t) dt \tag{2}$$

and

$$\int_a^x \frac{(t-x)^{i+1}}{(i+1)!} p(t)y'(t) dt = -p(a)y(a) \frac{(a-x)^{i+1}}{(i+1)!} - \int_a^x \left[\frac{(t-x)^i}{i!} p(t) + \frac{(t-x)^{i+1}}{(i+1)!} p'(t) \right] y(t) dt \tag{3}$$

The differential equation (E) equivalent at integral equation :

$$(E_i) : \forall x \in I, \int_a^x h_{i,x}(t)y(t) dt = f_i(x) \tag{4}$$

with

$$h_{i,x}(t) = \frac{(t-x)^{i-1}}{(i-1)!} - \frac{(t-x)^i}{i!} p(t) + \frac{(t-x)^{i+1}}{(i+1)!} (q(t) - p'(t)) \tag{5}$$

$$f_i(x) = -y(a) \frac{(a-x)^i}{i!} + [y'(a) + p(a)y(a)] \frac{(a-x)^{i+1}}{(i+1)!} + \int_a^x \frac{(t-x)^{i+1}}{(i+1)!} g(t) dt \tag{6}$$

Linear System and Approximate Solution

Next following the method suggested in [1,6]. We fix one positive integer $n \geq 1$. We have:

$$y(t) = \sum_{k=0}^{n-1} y^{(k)}(x) \frac{(t-x)^k}{k!} + R_{n,x}(t) \tag{7}$$

where $R_{n,x}(t)$ denotes integral remainder

$$R_{n,x}(t) = \int_x^t \frac{(t-s)^{n-1}}{(n-1)!} y^{(n)}(s) ds \tag{8}$$

In particular, if the desired solution $y(t)$ is a polynomial of degree equal to or less than $n-1$, then $R_{n,x}(t) = 0$.

We put for all i and j positive integer ≥ 1 :

$$b_{ij}(x) = \int_a^x h_{i,x}(t) \frac{(t-x)^{j-1}}{(j-1)!} dt \tag{9}$$

For an integer $i \geq 1$, the function y is solution of (E) if and only if we have:

$$\sum_{j=1}^n b_{ij}(x) y^{(j-1)}(x) = f_i(x) \tag{10}$$

We consider the matrix:

$$B_n(x) = (b_{ij}(x))_{i,j=1, \dots, n} \tag{11}$$

and the column

*Corresponding author: Nadhem Echi, Department of Mathematics, El Manar University, Tunis 1002, Tunisia, E-mail: nadhemechi_fsg@yahoo.fr

Received October 02, 2012; Published February 12, 2013

Citation: Echi N (2013) Approximate Solution of Second-Order Linear Differential Equation. 2: 607 doi:[10.4172/scientificreports.607](https://doi.org/10.4172/scientificreports.607)

Copyright: © 2013 Echi N. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

$$F_n(x) = \begin{pmatrix} f_1(x) \\ \vdots \\ f_n(x) \end{pmatrix}, Y_n(x) = \begin{pmatrix} y(x) \\ y'(x) \\ \vdots \\ y^{(n-1)}(x) \end{pmatrix} \tag{12}$$

For all positive integer $n \geq 2$, the equation (E) equivalent at

$$\forall x \in I, B_n(x)Y_n(x) = F_n(x) \tag{13}$$

Application of Cramer's rule to the resulting system yields an approximate solution of equation (1). It is also noted that not only $y(x)$ but also $y(i)(x)$ ($1 \leq i \leq n-1$) are determined via solving the resulting system. The solution $y(x)$ can be approximately obtained to be:

$$y_n(x) \equiv \frac{\det(B_1(x))}{\det(B_n(x))} \tag{14}$$

where

$$B_1(x) = \begin{pmatrix} f_1(x) & b_{12}(x) & \dots & b_{1n}(x) \\ f_2(x) & b_{22}(x) & \dots & b_{2n}(x) \\ \vdots & \vdots & \ddots & \vdots \\ f_n(x) & b_{n2}(x) & \dots & b_{nn}(x) \end{pmatrix} \tag{15}$$

Error Analysis

Approximation of B_{ij}

We are going to use the explore Taylor's approximations, in a of b_{ij} at order $i+j+n-1$. Let: $i \geq 1$ and $j \geq 1$ For all $k = 0, \dots, i+j-2$, we have:

$$b_{ij}^{(k)}(x) = (-1)^{i+j} \int_a^x [C_{i+j-2}^{j-1} \frac{(x-t)^{i+j-k-2}}{(i+j-k-2)!} - C_{i+j-1}^{j-1} \frac{(x-t)^{i+j-k-1}}{(i+j-k-1)!} p(t)] dt + C_{i+j}^{j-1} \frac{(x-t)^{i+j-k}}{(i+j-k)!} (p'(t) + q(t)) dt \tag{16}$$

In particular, we have :

$$b_{ij}^{(i+j-2)}(x) = (-1)^{i+j} [C_{i+j-2}^{j-1} (x-a) - C_{i+j-1}^{j-1} \int_a^x (x-t)p(t) dt + C_{i+j}^{j-1} \int_a^x \frac{(x-t)^2}{2} (p'(t) + q(t)) dt] \tag{17}$$

Furthermore, we have:

$$b_{ij}^{(i+j-1)}(x) = (-1)^{i+j} C_{i+j-2}^{j-1} + (-1)^{i+j-1} C_{i+j-1}^{j-1} \int_a^x p(t) dt + (-1)^{i+j} C_{i+j}^{j-1} \int_a^x (x-t)(p'(t) + q(t)) dt \tag{18}$$

$$b_{ij}^{(i+j)}(x) = (-1)^{i+j-1} C_{i+j-1}^{j-1} p(x) + (-1)^{i+j} C_{i+j}^{j-1} \int_a^x (p'(t) + q(t)) dt = (-1)^{i+j} [C_{i+j}^{j-1} - C_{i+j-1}^{j-1}] p(x) - (-1)^{i+j} C_{i+j}^{j-1} p(a) + (-1)^{i+j} C_{i+j}^{j-1} \int_a^x q(t) dt \tag{19}$$

$$b_{ij}^{(i+j+1)}(x) = (-1)^{i+j} [C_{i+j}^{j-1} - C_{i+j-1}^{j-1}] p'(x) + C_{i+j}^{j-1} q(x) \tag{20}$$

Thus, for all $k = 0, \dots, i+j-2$, we have:

$$b_{ij}^{(k)}(a) = 0 \tag{21}$$

So we have:

$$b_{ij}^{(i+j-1)}(a) = (-1)^{i+j} C_{i+j-2}^{j-1} \tag{22}$$

$$b_{ij}^{(i+j)}(a) = (-1)^{i+j-1} C_{i+j-1}^{j-1} p(a) \tag{23}$$

The explore Taylor's approximations, in a of $b_{ij}^{(i+j+1)}$ at order $n-2$, give that of b_{ij} at order $n+i+j-1$ following :

$$b_{ij}(x) = b_{(n)ij}(x) + o((x-a)^{n+i+j-1}) \tag{24}$$

with

$$b_{(n)ij}(x) = -\frac{(a-x)^{i+j-1}}{(i+j-1)(i-1)!(j-1)!} - p(a) \frac{(a-x)^{i+j}}{(i+j)!i!(j-1)!} \tag{25}$$

For $i \geq 1$ and $j \geq 1$, we put :

$$\tilde{b}_{ij}(x) = \frac{(a-x)^{i+j-1}}{(i-1)!(j-1)!} \tilde{b}_{ij}(x) \tag{26}$$

We have:

$$\tilde{b}_{ij}(x) = \tilde{b}_{(n)ij}(x) + o((x-a)^n) \tag{27}$$

with:

$$\begin{aligned} \tilde{b}_{(n)ij}(x) &= \frac{1}{i+j-1} - \frac{p(a)}{(i+j)i} (x-a) \\ &+ \frac{j-1}{i(i+1)(i+j)} \sum_{l=2}^n \left[\frac{1}{C_{i+j+l-1}^{l-1}} \right] \frac{p^{(l-1)}(a)}{(l-1)!} (x-a)^l \\ &+ \frac{1}{i(i+1)(i+j+1)} \sum_{l=2}^n \left[\frac{1}{C_{i+j+l-1}^{l-2}} \right] \frac{q^{(l-2)}(a)}{(l-1)!} (x-a)^l \end{aligned} \tag{28}$$

So the approximation in a , at order n , of \tilde{b}_{ij} use the approximation at order $n-1$ of p and the approximation at order $n-2$ of q .

We put: $\tilde{B}_n(x) = (\tilde{b}_{ij}(x))_{i,j=1,\dots,n}$ and $\tilde{B}_{(n)}(x) = (\tilde{b}_{(n)ij}(x))_{i,j=1,\dots,n}$

Let $D_n(x)$ the diagonal matrix having $\frac{(a-x)^{i-1}}{(i-1)!}$ as i -th diagonal coefficient. For all : $i=1, \dots, n$.

$$D_n(x) = \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & (a-x) & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & 0 \\ 0 & \dots & 0 & \frac{(a-x)^{n-1}}{(n-1)!} \end{pmatrix} \tag{29}$$

We have :

$$B_n(x) = (x-a)D_n(x)\tilde{B}_n(x)D_n(x) \tag{30}$$

$$B_{(n)}(x) = (x-a)D_n(x)\tilde{B}_{(n)}(x)D_n(x) \tag{31}$$

$$\tilde{B}_n(x) - \tilde{B}_{(n)}(x) = o((x-a)^n) \tag{32}$$

Approximation of F_n

We have:

$$f_i(x) = -y(a) \frac{(a-x)^i}{i!} + [y'(a) + p(a)y(a)] \frac{(a-x)^{i+1}}{(i+1)!} + \int_a^x \frac{(t-x)^{i+1}}{(i+1)!} g(t) dt \tag{33}$$

We are going to use the explore Taylor's approximations, in a of $f_i(x)$ at order $i+n$. For $k \leq i+1$, we have :

$$\frac{d^k}{dx^k} \int_a^x \frac{(t-x)^{i+1}}{(i+1)!} g(t) dt = (-1)^k \int_a^x \frac{(t-x)^{i+1-k}}{(i+1-k)!} g(t) dt \tag{34}$$

and

$$f_i^{(i+2)} = (-1)^{i+1} g(x) \tag{35}$$

Thus we have:

$$f_i(x) = f_{(n)_i}(x) + o((x-a)^{i+n}) \tag{36}$$

with

$$f_{(n)_i}(x) = -y(a) \frac{(a-x)^i}{i!} + [y'(a) + p(a)y(a)] \frac{(a-x)^{i+1}}{(i+1)!} + \frac{(-1)^{i+1}}{(i+2)!} \sum_{k=0}^{n-2} \frac{1}{C_{k+i+2}^k} \frac{g^{(k)}(a)}{k!} (x-a)^{k+i+2} \tag{37}$$

We put:

$$F_{(n)}(x) = \begin{pmatrix} f_{(n)_1}(x) \\ f_{(n)_2}(x) \\ \vdots \\ f_{(n)_n}(x) \end{pmatrix} \tag{38}$$

We have:

$$F_n(x) = (x-a)D_n(x)\tilde{F}_{(n)}(x) \tag{39}$$

with:

$$\tilde{F}_{(n)}(x) = \begin{pmatrix} f_{(n)_1}(x) \\ \tilde{f}_{(n)_1}(x) \\ \tilde{f}_{(n)_2}(x) \\ \vdots \\ \tilde{f}_{(n)_n}(x) \end{pmatrix} \tag{40}$$

For $i=1, \dots, n$, we have:

$$\tilde{f}_{(n)_i}(x) = \frac{1}{i} y(a) + \frac{1}{i(i+1)} [y'(a) + p(a)y(a)] + \frac{1}{i(i+1)(i+2)} \sum_{k=2}^n \frac{1}{C_{k+i}^{k-2}} \frac{g^{(k-2)}(a)}{(k-2)!} (x-a)^k \tag{41}$$

The construction of $F(n)$ uses $p(a)$, the approximate of g in a at order $n-2$. as well as the initial condition of $y(a)$ and $y'(a)$. We have :

$$F_n(x) = (x-a)D_n(x)[\tilde{F}_{(n)}(x) + o((x-a)^n)] \tag{42}$$

Lemma:

1. The matrix $\left(\frac{1}{i+j-1}\right)_{i,j=1, \dots, n}$ has positive determinant.
2. We have :

$$\frac{\det\left(\frac{1}{i+j-1}\right)_{i,j=1, \dots, n}}{\det\left(\frac{1}{i+j}\right)_{i,j=1, \dots, n}} = C_{2n} \tag{43}$$

Proof: We provide $C[X]$ by scalar product: $\langle v, w \rangle = \int_{-1}^1 v(x)\overline{w(x)} dx$

1. Let v_0, v_1, \dots the orthogonal basis given by:

$$v_k = \frac{\sqrt{2k+1}}{\sqrt{2}} L_k$$

with $L_k = \frac{1}{2^k k!} \frac{d^k}{dx^k} (x^2 - 1)^k$ Legendre polynomial of degree k .

We put: $e_k = \frac{(x+1)^k}{2^k \sqrt{2}}$. The Gram Schmidt matrix of linearly independent system (e_0, \dots, e_{n-1}) is $\tilde{B}_n(a) = \left(\frac{1}{i+j-1}\right)_{i,j=1, \dots, n}$.

In particular we have : $\det(\tilde{B}_n(a)) > 0$.

2. We put: $V_n = \left(\frac{1}{i+j}\right)_{i,j=1, \dots, n}$. For: $j=1, \dots, n-1$

$$V_n e_j = \begin{pmatrix} \langle e_0, e_j \rangle \\ \vdots \\ \langle e_{n-1}, e_j \rangle \end{pmatrix}, \quad \tilde{B}_n(a) e_j = \begin{pmatrix} \langle e_0, e_{j-1} \rangle \\ \vdots \\ \langle e_{n-1}, e_{j-1} \rangle \end{pmatrix}$$

where e_j is the j -th vector of the canonical basis.

Let $P = e_n - \langle e_n, v_n \rangle v_n = \sum_{j=0}^{n-1} x_j e_j$ orthogonal projection of e_n on $C_{n-1}[X]$.

We have : $\forall k = 0, \dots, n-1, \langle e_k, e_n \rangle = \langle e_k, P \rangle = \sum_{j=0}^{n-1} x_j \langle e_k, e_j \rangle$.

Thus, n -th column of (V_n) equal:

$$x_0 \begin{pmatrix} \langle e_0, e_0 \rangle \\ \vdots \\ \langle e_{n-1}, e_0 \rangle \end{pmatrix} + \sum_{j=1}^{n-1} x_j \begin{pmatrix} \langle e_0, e_j \rangle \\ \vdots \\ \langle e_{n-1}, e_j \rangle \end{pmatrix}$$

Thus: $\det(V_n) = (-1)^{n-1} x_0 \det(\tilde{B}_n(a))$. We have :

$$P(-1) = -\langle e_n, v_n \rangle v_n(-1) = x_0 \frac{1}{\sqrt{2}}$$

n -th order vector entry of e_n equal n -th order vector entry of $\langle e_n, v_n \rangle v_n$. One arrives at :

$$\begin{aligned} x_0 \frac{1}{\sqrt{2}} &= \frac{n\text{-th order vector entry of } e_n}{n\text{-th order vector entry of } v_n} v_n(-1) \\ &= \frac{n\text{-th order vector entry of } e_n}{n\text{-th order vector entry of } Q} Q(-1) \end{aligned}$$

such as $Q = \frac{1}{n!} \frac{d^n}{dx^n} (x^2 - 1)^n$. Taylor's expansion : $Q(-1) = (-2)^n$. Thus :

$$x_0 \frac{1}{\sqrt{2}} = -\frac{2^n \sqrt{2}}{(2n)!} (-2)^n = (-1)^{n-1} \frac{1}{\sqrt{2}} C_{2n}$$

Thus:

$$\frac{\det\left(\frac{1}{i+j-1}\right)_{i,j=1, \dots, n}}{\det\left(\frac{1}{i+j}\right)_{i,j=1, \dots, n}} = C_{2n}$$

We have : $\tilde{B}_{(n)}(a) = \tilde{B}_n(a) = \left(\frac{1}{i+j-1}\right)_{i,j=1, \dots, n}$.

As, $\det(\tilde{B}_{(n)}(a)) \neq 0$, thus taking x close of a , $\det(\tilde{B}_{(n)}(x)) \neq 0$.

Hence, taking x close of a ; $x \neq a$ $B_{(n)}(x) \neq 0$. Similarly, defines to be :

$$Y_{(n)}(x) = B_{(n)}(x)^{-1} F_{(n)}(x) \tag{44}$$

We put :

$$Y_{(n)}(x) = \begin{pmatrix} y_{(n)0}(x) \\ y_{(n)1}(x) \\ \vdots \\ y_{(n)n-1}(x) \end{pmatrix} \tag{45}$$

From (38) and (41) we have:

$$\tilde{B}_{(n)}(x)D_n(x)Y_{(n)}(x) = F_{(n)}(x) \tag{46}$$

The first element of $D_n(x)Y_{(n)}(x)$ is $y_{(n)0}$ just what we look for.

Theorem:

1. Taking $k=0, \dots, n-1$, we have :

$$y_{(n)k}(x) - y^{(k)}(x) = 0((x-a)^{n-k-1}) \tag{47}$$

2. We have :

$$y(x) = y_{(n)0}(x) + \frac{1}{C_{2n}^n} \frac{y^{(n)}(a)}{n!} (x-a)^n + o((x-a)^n) \tag{48}$$

We have:

$$F_n(x) = F_{(n)}(x) + (x-a)D_n(x) o((x-a)^n)$$

Taking x close of a , $x \neq a$, we have :

$$B_n(x)Y_n(x) = B_{(n)}(x)Y_{(n)}(x)Y_{(n)}(x) + (x-a)D_n(x) o((x-a)^n)$$

one arrives at,

$$\tilde{B}_n(x)D_n(x)Y_n(x) = \tilde{B}_{(n)}(x)D_n(x)Y_{(n)}(x) + \frac{y^{(n)}(a)}{n!} (x-a)^n (-1)^{n-1} \begin{pmatrix} 1 \\ 1+n \\ 1 \\ 2+n \\ \vdots \\ \vdots \\ 1 \\ 2n \end{pmatrix} + o((x-a)^n)$$

Thus :

$$\tilde{B}_n(x)[D_n(x)Y_n(x) - D_n(x)Y_{(n)}(x)] = \frac{y^{(n)}(a)}{n!} (x-a)^n (-1)^{n-1} \begin{pmatrix} 1 \\ 1+n \\ 1 \\ 2+n \\ \vdots \\ \vdots \\ 1 \\ 2n \end{pmatrix} + o((x-a)^n)$$

Furthermore:

$$D_n(x)Y_n(x) - D_n(x)Y_{(n)}(x) = \frac{y^{(n)}(a)}{n!} (x-a)^n (-1)^{n-1} \tilde{B}_{(n)}^{-1}(a) \begin{pmatrix} 1 \\ 1+n \\ 1 \\ 2+n \\ \vdots \\ \vdots \\ 1 \\ 2n \end{pmatrix} + o((x-a)^n)$$

By looking at the first constituent and allowing that $\tilde{B}_{(n)}(a) = \left(\frac{1}{i+j-1} \right)_{i,j=1,\dots,n}$, we obtain:

$$y(x) - y_{(n)0}(x) = \frac{y^{(n)}(a)}{n!} (x-a)^n (-1)^{n-1} \begin{pmatrix} \frac{1}{1+n} & \frac{1}{2} & \dots & \frac{1}{n} \\ \frac{1}{2+n} & \frac{1}{3} & \dots & \frac{1}{1+n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{1}{2n} & \frac{1}{1+n} & \dots & \frac{1}{2n} \end{pmatrix} + o((x-a)^n)$$

Thus:

$$y(x) - y_{(n)0}(x) = \frac{1}{C_{2n}^n} \frac{y^{(n)}(a)}{n!} (x-a)^n + o((x-a)^n)$$

Example: We consider the following second-order differential equation with variable coefficients

$$y''(x) - 2y'(x)y(x) = 0 \tag{49}$$

under the initial condition

$$y(0) = 0, y'(0) = 1 \tag{50}$$

This equation has an exact solution

$$y(x) = \tan(x) \tag{51}$$

Using the present method, we can evaluate its approximate solution. For example, when $n=3$ one can find the corresponding third-order approximations as:

$$y_3(x) = \frac{162x^8 - 8904x^6 + 70560x^4}{-19x^9 + 1734x^7 - 32760x^5 + 70560x^3} \tag{52}$$

Of course, when taking a larger n , it is expected that more accurate approximate results can be given. In order to show the variation of the accuracy of approximations (Table 1).

It indicates that the present method gives a very fast convergence, even for x near some unbounded points.

Conclusions

This paper proposes an effective method for determining the approximate solution of second-order differential equation with constant and varying coefficients. The second-order differential equations are transformed into a Volterra integral equation. By using

x	Exact solution y(x)	Approximate solution y ₃ (x)
0.1	0.1003347	0.1003394
0.5	0.5463025	0.5469448
1	1.5574077	1.5644186
1.57	1.2557656 × 10 ³	1.2536283 × 10 ³
1.57078	6.1249009 × 10 ⁴	6.0755230 × 10 ⁴

Table 1: Indicates that the present method gives a very fast convergence, even for x near some unbounded points.

Taylor's expansion of the unknown function, the resulting Volterra equation can be approximately solved [7]. Based on these results, an approximation of second-order differential equations can be directly constructed. Test example is given, and obtained results turn out that the present method is efficient, and simple. Moreover, it can be implemented by symbolic computation at any personal computer.

References

1. Xie F, Gao X (2004) Exact traveling wave solutions for a class of nonlinear partial differential equations. *Chaos, Solutions & Fractals* 19: 1113-1117.
2. Abbasbandy S (2006) Homotopy perturbation method for quadratic Riccati differential equation and comparison with Adomian's decomposition method. *Appl Math Comput* 172: 485-490.
3. Li XF (2006) Approximate solution of linear ordinary differential equations with variable coefficients. *Math Comput Simul* 75: 113-125.
4. Tang BQ, Li XF (2007) A new method for determining the solution of Riccati differential equations. *Appl Math Comput* 194: 431-440.
5. Adomian G, Rach R (1992) Noise terms in decomposition series solution. *Comput Math Appl* 11: 61-64.
6. El-Tawil MA, Bahnasawi AA, Abdel-Naby A (2004) Solving Riccati differential equation using Adomian's decomposition method. *Appl Math Comput* 157: 503-514.
7. Chen Y, Li Biao (2004) General projective Riccati equation method and exact solutions for generalized KdV-type and KdV-Burgers-type equation with nonlinear terms of any order. *Chaos, Solitons Fractals* 19: 977-984.