

## The Role of Albumin Level and Blood Urea Nitrogen/ Albumin Ratio in Prediction of Prognosis of Community Acquired Pneumonia

Evrım Eylem AKPINAR MD<sup>1</sup>, Derya HOŞGÜN MD<sup>1</sup>, Beyza DOĞANAY PhD<sup>2</sup>, Meral GÜLHAN Prof<sup>1</sup>

<sup>1</sup>Ufuk University, Department of Chest Diseases, Ankara, Turkey

<sup>2</sup>Ankara University, Department of Biostatistics, Ankara, Turkey

### Abstract

**Background and aim:** Community-Acquired Pneumonia (CAP) is an important cause of morbidity and mortality worldwide. The accuracy of scoring scales is lower regarding the evaluation of mortality and need for ICU treatment. This study aimed to evaluate the role of the albumin level and Blood urea nitrogen /Albumin (BUN/Alb) ratio in the prediction of disease severity and one-month mortality.

**Materials and methods:** Patients hospitalized for CAP were included. Venous blood samples were obtained to measure albumin levels and calculate the ratio of BUN/Alb. The correlations of serum albumin levels and BUN/Alb ratio with the requirement of ICU, development of complications and one-month mortality were evaluated.

**Results:** 216 patients were enrolled. Patients who needed ICU treatment had a higher BUN/Alb ratio ( $p=0.029$ ). The cut-off level of BUN/Alb in prediction of ICU need was found 4.15 (sensitivity 76%, specificity 49%). Low albumin level was an independent risk factor for ICU need (OR: 5.263, 95% CI: 1.996 to 13.889,  $p=0.001$ ). The cut-off level of albumin in prediction of ICU need was 3.39 g/dl (sensitivity 71%, specificity 71%). Low albumin level was independent predictive factor for the development of complications (OR: 4.902, 95% CI: 1.595 to 14.925,  $p=0.005$ ). The cut-off level of albumin in prediction of development of complication was 3.44 g/dl (sensitivity 79%, specificity 69%).

**Conclusions:** The CAP patients who have higher BUN/Alb ratio are under higher risk of the development of need for ICU treatment. Low albumin level is a more valuable predictor than BUN/Alb ratio for prognosis of CAP.

### Introduction

Community-Acquired Pneumonia (CAP) is an important cause of morbidity and mortality worldwide [1]. The mortality rate of CAP ranges from less than 5% among outpatients to 12% among hospitalized patients [2]. The need for an intensive care unit (ICU) is also an important problem for clinicians to overcome during the course of CAP [3,4]. CURB-65 (confusion, urea nitrogen, respiratory rate, blood pressure,  $\geq 65$  years) and Pneumonia Severity Index (PSI) are the most frequently used scoring scales to assess the disease severity [5,6]. These scoring tools are useful to decide between hospitalization and outpatient treatment with an oral antibiotic. However, their accuracy is lower regarding the evaluation of mortality and need for ICU treatment [4,7]. In recent years, the role of several biomarkers, such as C-Reactive Protein (CRP) and procalcitonin, have been studied to predict the severity and prognosis of CAP and its correct diagnosis and microbiological etiology [8].

Serum albumin concentration has been used as an indicator of nutritional status for years [9]. Hypoalbuminemia was related with poor outcome in several clinical conditions, including CAP [9-11]. In a recent study by Lee et al., it was reported that albumin was associated with 28-day mortality in patients hospitalized with a CAP diagnosis [12]. The inflammatory reaction was reported as a primary reason for hypoalbuminemia in elderly patients with CAP [13]. The Blood Urea Nitrogen (BUN) level is an important biochemical parameter showing renal hypoperfusion. It is one of the contributing parameters for both the CURB-65 and PSI severity scoring scales. Previous studies have shown that patients with CAP who had higher BUN levels [14,15] and lower albumin levels [6,15] had higher mortality rates. It has recently been reported that the BUN/Albumin (BUN/Alb) ratio was an independent predictor for both mortality and the severity of CAP [16].

This study aimed to evaluate the role of the albumin and BUN/Alb ratio in the prediction of development of complications, need of ICU and one-month mortality. The secondary end-point of the study was to evaluate other factors affecting development of complications, need of ICU and one-month mortality.

### Materials and Methods

The study included 216 consecutive patients hospitalized for CAP between the dates of January 2011 and June 2012. This was a prospective observational study. Patients who were  $\geq 18$  years, admitted from the community, had at least two clinical signs and symptoms related to pneumonia (fever  $> 38^{\circ}\text{C}$ , cough, chest pain, dyspnea, or crackles on auscultation), and presented new infiltration on chest radiography were included in the study. Patients were excluded if they were immunocompromised (i.e., using immunosuppressive therapy, having a human immunodeficiency virus infection, malignancy, or undergoing chemotherapy), had been hospitalized and/or used any antibiotic in last two weeks, or had chronic renal or liver failure. The design of the study is shown in Figure 1. The study was approved by the local ethics committee, and informed consent was obtained from all patients included in the study.

### Blood sampling


Venous blood samples were obtained from patients upon admission before starting antibiotic treatment. Hemogram measurements and biochemical analyses to measure the levels of albumin, CRP, BUN,

\*Corresponding author: Evrim Eylem AKPINAR, Chest Diseases Specialist, Ufuk University Medical Faculty, Dr. Ridvan Ege Hospital, Department of Chest Diseases, Mevlana Bulvarı (Konya Yolu) No: 86-88 Balgat 06540- Ankara, Turkey, Tel: +90 312 204 43 31 ; Fax: +90 312 204 40 55 ; E-mail: [drevrimeylem@gmail.com](mailto:drevrimeylem@gmail.com)

Received August 20, 2013; Accepted October 18, 2013; Published October 21, 2013

Citation: AKPINAR EE, HOŞGÜN D, DOĞANAY B, GÜLHAN M (2013) The Role of Albumin Level and Blood Urea Nitrogen/ Albumin Ratio in Prediction of Prognosis of Community Acquired Pneumonia. J Pulm Respir Med 3: 159. doi:10.4172/2161-105X.1000159

Copyright: © 2013 AKPINAR EE, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



<b>Minor criteria</b>	Respiratory rate $\geq 30$ breaths/min PaO <sub>2</sub> /FIO <sub>2</sub> ratio $\leq 250$ Multilobar infiltrates Confusion/disorientation Uremia (BUN level, $\geq 20$ mg/dl) Leukopenia (WBC count, $<4000$ /mm <sup>3</sup> ) Thrombocytopenia (platelet count, $<100.000$ /mm <sup>3</sup> ) Hypothermia (core temperature $<36^{\circ}\text{C}$ ) Hypotension requiring excessive fluid resuscitation
<b>Major criteria</b>	Invasive mechanic ventilation Septic shock with the need for vasopressors

\*The patients who had at least 1 major or 3 minor criterion were accepted to intensive care unit (ICU).

**Table 1:** Criteria for ICU need\*.

creatinine were made, and the ratio of BUN/Alb was calculated for all participants.

### Evaluation of patients

A detailed medical history of patients was recorded. Comorbidities [i.e., Chronic Obstructive Pulmonary Disease (COPD), Diabetes Mellitus (DM), Congestive Heart Failure (CHF), bronchiectasis, malignancy] were noted. The severity of CAP was assessed using the CURB-65 and PSI for all patients [5,6]. A blood culture was performed for patients who had a fever  $>38^{\circ}\text{C}$ . An arterial blood gas analysis was performed. Empirical antibiotic treatments were started according to the recommendations of the Turkish Thoracic Society [17]. The need for ICU was decided according to ICU criteria of Turkish Thoracic Society (Table 1) [17]. Thoracocentesis was performed to the patients who had pleurisy upon admission or who developed pleural effusion during treatment to evaluate the nature of the accompanying fluid. Patients who developed respiratory failure (paO<sub>2</sub> $<60$  mmHg and/or paCO<sub>2</sub> $>45$  mmHg) during the course of the disease were also noted.

The endpoints of the study were the requirement of ICU, development of complications [e.g., parapneumonic effusion (exudative

pleurisy on admission or during antibiotic treatment) or respiratory failure], or mortality within one month.

### Statistical analysis

Continuous data were described as the mean $\pm$ standard deviation, and median (minimum-maximum) categorical data were described as frequencies and percentages. Differences between groups were evaluated by Student's t test and Mann Whitney U test depending of the normality of continuous data. A Chi-Square test or Fisher's Exact test were used for categorical data when applicable. A Spearman correlation coefficient was used to evaluate the relationship between continuous variables. Logistic regression with backward elimination was used to evaluate independent risk factors or the development of complications and need for ICU treatment. The results were expressed as odds ratios (OR) with 95% Confidence Intervals (CI). Variables significantly associated with the development of complications or need for ICU treatment at the 0.20 levels in the univariate analysis were considered in a backward elimination of variables in a logistic regression. IBM SPSS Statistics for Windows Version 20 was used for all analyses. A p value  $<0.05$  was considered statistically significant.

### Results

In total, 216 patients (117 male, 99 female) were enrolled in the study. Demographic properties and severity of pneumonia (CURB-65, PSI) laboratory findings are shown on Table 2. Complications developed in 19 patients (16 parapneumonic effusion, 3 respiratory failure). Twenty-one (9.7%) of the patients who were hospitalized needed ICU treatment. The one-month mortality rate of CAP in our population was 6.7%. The levels of albumin of patients who were inpatient, non-ICU treated were significantly higher than the albumin levels of patients who required treatment in the ICU ( $p<0.0001$ ). Microbiological investigations were conducted in 53 (24.5%) of patients. Isolation of pathogen bacteria was


<b>Gender</b>	<b>n %</b>
Male	117 (54.2)
Female	99 (45.8)
<b>Age years (n=216)</b>	<b>n %</b>
mean $\pm$ SD	68.79 $\pm$ 13.27
median (min-max)	70 (17-96)
<b>CURB-65</b>	<b>n %</b>
2	177 (75)
3	39 (25)
<b>PSI-stage</b>	<b>n %</b>
3	197 (91.21)
4	15 (6.94)
5	4 (1.85)
<b>WBC (/mm<sup>3</sup>)</b>	<b>n %</b>
$>10000$	115 (53.3)
$<6000$	101 (46.7)
<b>ESR (mm/h) (n=216)</b>	<b>n %</b>
mean $\pm$ SD	37.62 $\pm$ 23.54
median (min-max)	32 (1-97)
<b>CRP (mg/dl) (n=216)</b>	<b>n %</b>
mean $\pm$ SD	61.59 $\pm$ 65.14
median (min-max)	39.5 (0.18-360)
<b>Serum Albumin (g/dl) (n=216)</b>	<b>n %</b>
mean $\pm$ SD	3.63 $\pm$ 0.53
median (min-max)	3.58 (2.25-5 07)
<b>BUN/ Alb (n=216)</b>	<b>n %</b>
mean $\pm$ SD	5.53 $\pm$ 3.58
median (min-max)	4.3 (1.57-25.66)
<b>paO<sub>2</sub> (mmHg) (n=208)</b>	<b>n %</b>
mean $\pm$ SD	61.08 $\pm$ 12.32
median (min-max)	60 (32-107)

**Table 2:** Demographic properties, severity of pneumonia, and laboratory findings of patients.

Microorganism	n (%)	Culture
<i>Streptococcus pneumoniae</i>	9 (4.16)	Sputum
<i>Mycoplasma pneumoniae</i>	2 (0.92)	Sputum
<i>Chlamydia pneumoniae</i>	1 (0.46)	Sputum
<i>Haemophilus influenzae</i>	3 (1.38)	Sputum
<i>Klebsiella pneumoniae</i>	3 (1.38)	TTA
	1 (0.46)	Blood
<i>Pseudomonas aeruginosa</i>	2 (0.92)	TTA
MRSA	2 (0.92)	TTA
<i>Acinetobacter</i>	1 (0.46)	TTA
	1 (0.46)	Blood
<b>Total</b>	<b>25 (11.57)</b>	

MRSA: *meticillin-resistant staphylococcus aureus*, TTA: transtracheal aspirate.

Table 3: Causative agents in positive cultures of patients.



achieved in 25 (11.57%) of these patients. Table 3 shows the pathogen bacteria isolated from cultures.

The factors affecting the development of complications were evaluated. A positive relationship was found between complication rate and severity of pneumonia (CURB-65=2 vs. CURB-65 > 2, PSI=3 vs. PSI >3, respectively; p<0.0001, p<0.0001). The patients who needed the ICU developed more complications compared with those who did not (p<0.0001). The rate of complications was not affected by age (<65 years vs. ≥65 years), gender, paO<sub>2</sub>, BUN, BUN/Alb ratio or co-morbidity (COPD, bronchiectasis, diabetes mellitus, malignancy, CHF) (all p>0.05). Logistic regression analysis showed that low albumin levels and poor PSI were independent predictive factors for the development of complications (OR: 4.902, 95% CI: 1.595 to 14.925, p=0.005; OR: 7.444, 95% CI: 2.565 to 21.605, p<0.0001, respectively). The cut-off level of albumin in prediction of development of complication was found to be 3.44 g/dl (sensitivity 79%, specificity 69%). The area under the ROC curve was found to be 0.751 (95% CI: 0.631-0.871) (p<0.001) (Figure 2).

The factors related to mortality were also investigated. Age (<65 years vs. ≥65 years), gender, BUN, BUN/ Alb ratio, albumin, paO<sub>2</sub>, and the presence of co-morbidities, including COPD, bronchiectasis, DM, malignancy did not affect mortality of the population studied. Patients who had more severe pneumonia (CURB-65 = 2 vs. CURB-65 > 2, PSI=3 vs. PSI>3) needed ICU treatment had higher rate of mortality (respectively; p=0.011, p=0.003, p=0.001). Because the rate of mortality was low, logistic regression analysis could not be done for mortality. The factors influencing mortality and the development of complications are shown in Table 4 and 5.

The need of ICU treatment was not affected by age (<65 years vs. ≥65 years), gender, or the presence of co-morbidities (all p>0.05). The severity of pneumonia (CURB-65=2 vs. CURB-65 > 2, PSI=3 vs. PSI>3) and low levels of albumin affected the need for ICU treatment (for all; p<0.0001). Patients who needed ICU treatment had a higher BUN/Alb ratio than those who did not (p=0.029). The cut-off level of BUN/Alb ratio to predict the need of ICU was found to be 4.15 (sensitivity 76%, specificity 49%). The area under the ROC curve was found to be 0.645 (95% CI: 0.525-0.766) (p=0.029) (Figure 3). The factors influencing the need for ICU treatment are shown in Table 6. Logistic regression analysis revealed that a low albumin level was an independent risk factor for the need of ICU treatment (OR: 5.263, 95% CI: 1.996 to 13.889, p=0.001). The cut-off level of albumin to predict the need of ICU was found to be 3.39 g/dl (sensitivity 71%, specificity 71%). The area under the ROC curve was found to be 0.726 (95% CI: 0.608-0.843) (p=0.001) (Figure 4).

## Discussion

This study showed the CAP patients who had higher BUN/Alb ratio were under higher risk of the development of need for ICU treatment. The cut-off level of BUN/Alb in prediction of ICU need was found 4.15 (sensitivity 76%, specificity 49%). But, the ratio did not predict the development of complications and mortality. Low albumin level was independent predictor for development of complication and need for ICU. The cut-off level of albumin in prediction of ICU need was 3.39 g/dl (sensitivity 71%, specificity 71%). Low albumin level was independent predictive factor for the development of complications the cut-off level of albumin in prediction of development of complication was 3.44 g/dl (sensitivity 79%, specificity 69%).

Biomarkers, including procalcitonin and B-type natriuretic peptide were found to be predictive for the prognosis of CAP [18,19]. But

	Complication n (%)		p	Mortality n (%)		p
	+	-		NS	S	
CURB-65=2	9 (4.1)	168 (78)	<0.0001 <sup>b</sup>	2 (0.93)	175 (81.02)	0.011 <sup>b</sup>
CURB-65>2	10 (4.6)	29 (13.3)		4 (1.85)	35 (16.2)	
PSI=3	6 (2.8)	161 (74.6)	<0.0001 <sup>b</sup>	1 (0.46)	166 (76.85)	0.003 <sup>b</sup>
PSI>3	13 (6)	36 (16.6)		5 (2.31)	44 (20.37)	
<b>Age (years)</b>						
<65	4 (1.85)	66 (30.56)	0.268 <sup>a</sup>	1 (0.46)	69 (31.94)	0.666 <sup>b</sup>
≥65	15 (6.94)	131 (60.65)		5 (2.31)	141 (65.28)	
<b>Gender</b>						
Male	8 (3.7)	109 (50.46)	0.269 <sup>a</sup>	3 (1.39)	114 (52.78)	1.000 <sup>b</sup>
Female	11 (5.09)	88 (40.74)		3 (1.39)	96 (44.44)	
<b>Need for ICU</b>						
Present	9 (4.17)	12 (5.56)	<0.0001 <sup>b</sup>	4 (1.85)	17 (7.87)	0.001 <sup>b</sup>
Absent	10 (4.63)	185 (85.65)		2 (0.93)	193 (89.35)	


CURB-65: Confusion, urea nitrogen, respiratory rate, blood pressure, ≥65 years, PSI: Pneumonia severity index, ICU: Intensive care unit, NS: Non-surviving, S: Surviving, a: p value was based on Chi-Square test, b: p value was based on Fisher's Exact Test.

Table 4: The factors influencing the development of complications and mortality.

	Complication		p	Mortality		p
	Absent n=19 Mean ± SD Median (Min-Max)	Present n=97 Mean ± SD Median (Min-Max)		NS n=6 Mean ± SD Median (Min-Max)	S n=210 Mean ± SD Median (Min-Max)	
<b>BUN</b> (mg/dl)	19.51 ± 11.09 16(4.68-76.48)	19.32 ± 12 15.9(7.63-57.03)	0.939 <sup>d</sup>	24.47 ± 17.2 18.89(9.01-57.03)	19.35 ± 11 16(4.68-76.48)	0.444 <sup>d</sup>
<b>Albumin</b> (g/dl)	3.66 ± 0.51 0.84(0.22-3.55)	3.19 ± 0.52 3.2(2.25-4.45)	<0.001 <sup>c</sup>	3.32 ± 0.61 3.32(2.61-4.25)	3.63 ± 0.53 3.6(2.25-5.07)	0.166 <sup>c</sup>
<b>BUN/Alb</b>	5.48 ± 3.57 4.3(1.57-25.66)	6.01 ± 3.62 4.6(2.5-16.57)	0.375 <sup>d</sup>	7.32 ± 5.09 4.92(3.40-16.57)	5.47 ± 3.53 4.3(1.57-25.66)	0.249 <sup>d</sup>
<b>Creatinin</b> (mg/dl)	0.92 ± 0.44 0.84(0.22-3.55)	0.86 ± 0.38 0.79(0.31-1.93)	0.656 <sup>d</sup>	1.04 ± 0.53 0.95(0.42-1.93)	0.91 ± 0.44 0.835(0.22-3.55)	0.493 <sup>d</sup>

SD: Standard deviation, BUN: Blood urea nitrogen, BUN/Alb: Blood urea nitrogen/ albumin, NS: Non-surviving, S: Surviving, c: p value was based on Student's t-test, d: p value was based on Mann Whitney U test.

**Table 5:** The laboratory parameters influencing the development of complications and mortality.



measurement of these biomarkers is not easily reachable and cheap. However, albumin and BUN levels are routinely measured in all hospitalized CAP patients.


The rate of the albumin synthesis is decreased in the acute phase of inflammation. The endotoxin from Gram-negative bacteria, cytokines like IL-6, chemokines cause capillary leakage of albumin. Hypoalbuminemia is a predictor of worse prognosis in hospitalized and critically ill patients [20]. Lee et al. [12] reported that albumin was associated with mortality in patients hospitalized due to CAP. Similarly, the serum albumin level upon admission was found to be predictive for mortality in CAP patients [6,21]. In our study, albumin was not an effective factor for one-month mortality in patients with CAP. But, the low albumin level was strongly associated with increase in development of complication and ICU need. The addition of albumin level of CAP patients to the parameters of severity scales (CURB-65 and/or PSI) may reinforce their predictive role in prognosis of the disease.

Previous studies have shown that patients with CAP who had higher BUN levels had higher mortality rates [6,14]. A high BUN level is one of the components of both the CURB-65 score and PSI [5,6].

BUN levels show a decrease in renal perfusion and indirectly predict the severity of pneumonia. The patients who have pneumonia are usually dehydrated that results from increase of BUN excretion from the kidneys. We investigated whether the ratio of these two easily measurable parameters (BUN and albumin) predicts the prognosis of CAP or not. In our study, the BUN/albumin ratio only predicted the ICU need in CAP patients.

Recently, Ugajin et al. [16] proposed that the BUN/Alb ratio is an independent predictor for both the severity of CAP and resulting mortality. Our study population had relatively milder disease. The ratio of patients who had greater than 2 points according to CURB-65 scale was 25%. This ratio was 40.6% in the study of Ugajin et al. [16]. The mortality rate also was lower in our study (6.7% vs 10.8%). These differences may be effective in contradictory results of the two studies about the role of the BUN/Alb ratio in the prediction of prognosis of CAP.

Lee et al. [12] found a 28-day mortality rate of 13.7% in patients hospitalized due to CAP. One-month mortality was lower (6.7%) in our study. Two factors of this study may have been more effective against



	ICU need (n %)		P
	Present n=21 n(%) Mean ± SD Median (Min-Max)	Absent n=195 n(%) Mean ± SD Median (Min-Max)	
CURB-65=2	1 (0.46)	176 (81.48)	<0.0001 <sup>b</sup>
CURB-65>2	20 (9.26)	19 (8.80)	
PSI=3	0 (0)	167 (77.31)	<0.0001 <sup>b</sup>
PSI>3	21 (9.72)	28 (12.96)	
<b>Age (years)</b>			
<65	5 (2.31)	65 (30.09)	0.376 <sup>a</sup>
≥65	16 (7.41)	130 (60.19)	
<b>Gender</b>			
Male	15 (6.94)	102 (47.22)	0.095 <sup>a</sup>
Female	6 (2.78)	93 (43.06)	
<b>BUN (mg/dl)</b>	23.08 ± 13.5 19.65(7.63-56.51)	19.11 ± 10.85 15.96(4.68-76.48)	0.198 <sup>d</sup>
<b>Albumin (g/dl)</b>	3.24 ± 0.49 3.23(2.43-4.25)	3.66 ± 0.52 3.61(2.25-5.07)	<0.0001 <sup>c</sup>
<b>BUN/Alb</b>	7.13 ± 4.3 5.26(2.68-18.65)	5.35 ± 3.45 4.19(1.57-25.66)	0.029 <sup>d</sup>

ICU: Intensive care unit, CURB-65: Confusion, urea nitrogen, respiratory rate, blood pressure, ≥65 years, PSI: Pneumonia severity index, SD: Standard deviation, BUN: Blood urea nitrogen, BUN/Alb: Blood urea nitrogen/ albumin, a: p value was based on Chi-Square test, b: p value was based on Fisher's Exact Test, c: p value was based on Student's t-test, d: p value was based on Mann Whitney U test.

**Table 6:** The factors influencing the need for intensive care unit.

mortality. First, empirical antibiotic treatment was applied following the national guidelines for CAP. Second, most of the patients included in our study had relatively milder pneumonia according to their severity scores.

This study had some limitations; it contained the patients with CAP from only one center. The patient population may not be representative for all CAP patients. Most of the patients who were included to the study

had relatively milder disease. This may be effective on the predictive role of BUN/Alb ratio in the course of CAP.

## Conclusions

CAP patients who have higher BUN/Alb ratio are under higher risk of the need for ICU treatment. The level of albumin is a more valuable predictor than BUN/Alb ratio in prediction of prognosis of CAP. Larger studies are necessary to evaluate the exact role and the cut-off value of the BUN/Alb ratio in predicting the severity and mortality of CAP.

## References

- Almirall J, Bolibar I, Vidal J, Sauca G, Coll P, et al. (2000) Epidemiology of community-acquired pneumonia in adults: a population-based study. *Eur Respir J* 15: 757-763.
- Niederman MS (2009) Community-acquired pneumonia: the U.S. perspective. *Semin Respir Crit Care Med* 30: 179-188.
- Lim WS, Baudouin SV, George RC, Hill AT, Jamieson C, et al. (2009) BTS guidelines for the management of community acquired pneumonia in adults: update 2009. *Thorax* 64 Suppl 3: iii1-55.
- Angus DC, Marrie TJ, Obrosky DS, Clermont G, Dremsizov TT, et al. (2002) Severe community-acquired pneumonia: use of intensive care services and evaluation of American and British Thoracic Society Diagnostic criteria. *Am J Respir Crit Care Med* 166: 717-723.
- Fine MJ, Auble TE, Yealy DM, Hanusa BH, Weissfeld LA, et al. (1997) A prediction rule to identify low-risk patients with community-acquired pneumonia. *N Engl J Med* 336: 243-250.
- Lim WS, van der Eerden MM, Laing R, Boersma WG, Karalus N, et al. (2003) Defining community acquired pneumonia severity on presentation to hospital: an international derivation and validation study. *Thorax* 58: 377-382.
- Ewig S, de Roux A, Bauer T, Garcia E, Mensa J, et al. (2004) Validation of predictive rules and indices of severity for community acquired pneumonia. *Thorax* 59: 421-427.
- Lippi G, Meschi T, Cervellini G (2011) Inflammatory biomarkers for the diagnosis, monitoring and follow-up of community-acquired pneumonia: clinical evidence and perspectives. *Eur J Intern Med* 22: 460-465.
- Randle NW, Hubert-Hartmann K, Mulheran LC (1984) Serum albumin levels: relationship to length of hospital stay. *Hosp Pharm* 19: 802-805.
- Ortqvist A, Hedlund J, Grillner L, Jalonen E, Kallings I, et al. (1990) Aetiology, outcome and prognostic factors in community-acquired pneumonia requiring hospitalization. *Eur Respir J* 3: 1105-1113.
- Anderson CF, Wochos DN (1982) The utility of serum albumin values in the nutritional assessment of hospitalized patients. *Mayo Clin Proc* 57: 181-184.
- Lee JH, Kim J, Kim K, Jo YH, Rhee J, et al. (2011) Albumin and C-reactive protein have prognostic significance in patients with community-acquired pneumonia. *J Crit Care* 26: 287-294.
- Hedlund JU, Hansson LO, Ortqvist AB (1995) Hypoalbuminemia in hospitalized patients with community-acquired pneumonia. *Arch Intern Med* 155: 1438-1442.
- Farr BM, Sloman AJ, Fisch MJ (1991) Predicting death in patients hospitalized for community-acquired pneumonia. *Ann Intern Med* 115: 428-436.
- Potgieter PD, Hammond JM (1996) The intensive care management, mortality and prognostic indicators in severe community-acquired pneumococcal pneumonia. *Intensive Care Med* 22: 1301-1306.
- Ugajin M, Yamaki K, Iwamura N, Yagi T, Asano T (2012) Blood urea nitrogen to serum albumin ratio independently predicts mortality and severity of community-acquired pneumonia. *International J General Med* 5: 583-589.
- Ozlu T, Bulbul Y, Alatas F (2009) Turkish Thoracic Society Guideline for diagnosis and treatment of community-acquired pneumonia in adults. *Turkish Thoracic Journal* 10: 3-18.
- Ramirez P, Ferrer M, Martí V, Reyes S, Martínez R, et al. (2011) Inflammatory biomarkers and prediction for intensive care unit admission in severe community-acquired pneumonia. *Crit Care Med* 39: 2211-2217.
- Christ-Crain M, Breidhardt T, Stolz D, Zobrist K, Bingisser R, et al. (2008) Use of B-type natriuretic peptide in the risk stratification of community-acquired pneumonia. *J Intern Med* 264: 166-176.

- 
20. Nicholson JP, Wolmarans MR, Park GR (2000) The role of albumin in critical illness. *Br J Anaesth* 85: 599-610.
21. Fry AM, Shay DK, Holman RC, Curns AT, Anderson LJ (2005) Trends in hospitalizations for pneumonia among persons aged 65 years or older in the United States, 1988-2002. *JAMA* 294: 2712-2719.