

Using of Coffee and Cardamom Mixture to Ameliorate Oxidative Stress Induced in γ -irradiated Rats

Refaat G. Hamzaa* and Nadia N. Osman

Department of food irradiation, Atomic Energy Authority, Cairo, Egypt

Abstract

Human exposure to ionizing radiation induced overproduction of free radicals leading to oxidative stress. This study aims to evaluate the possibility of using of coffee and cardamom mixture, as natural antioxidant compounds, to ameliorate the damage effect of oxidative stress. Phenolic contents in coffee and essential oils in cardamom were identified by using HPLC chromatography and GC/MS analysis. Four groups of adult male rats were used; the control group (A), the second group (B) received orally mixture extract of coffee and cardamom (60 mg/100g body weight) for 8 weeks, the third group (C) γ -irradiated (6 GY) and the fourth group (D) received orally mixture extract for 8 weeks and exposed to γ -irradiation at the 4th week. The results revealed that the administration of mixture extract of coffee and cardamom were significantly reduced the damage effect induced by γ -irradiation exposure. Accordingly, it could be concluded that, via the adjustment of the antioxidant status, decreasing the releasing of lipid peroxides and the subsequent amending of different biochemical parameters and some hormones, mixture of coffee and cardamom could ameliorate the hazardous effects of oxidative stress induced by irradiation exposure.

Keywords: Ionizing radiation; Coffee and cardamom; Phenolic; Essential oil; Antioxidants; Hormones

Introduction

Ionizing radiation being used in large number of therapeutic, industrial and other applications of nuclear power generation and developing new varieties of high yielding of new crops and enhancing storage-period of food materials[1-3]. In addition, Gamma radiation is associated with the generation of reactive oxygen species (ROS), causing oxidative damage particularly to various tissues and induce damages in DNA and cellular membrane [3-5].

A number of medicinal plants evaluated for their radioprotective efficacy have shown protective effects against the damaging effects of ionizing radiation. Plant extracts eliciting radioprotective efficacy contain a plethora of compounds including antioxidants, immunostimulants, cell proliferation stimulators, anti-inflammatory and antimicrobial agents, some of which may act in isolation as well as in combination with other constituents from the same plant [6-10].

Coffee (*Coffea Arabica*) has scientific interests because it is a rich source of a number of phenol compounds with antioxidant effects in vitro and contains several species of xanthines such as caffeine, theobromine and theophylline [11,12]. Main polyphenols in coffee are chlorogenic acids such as caffeic, ferulic, and p-coumaric acid, caffeoylquinic acid, with 5-O-caffeoyl-quinic acid [13,14]. Meta-analyses have concluded that moderate to high coffee consumption (three to six cups/day) is not significantly associated with an increased risk of coronary death or heart attack [15,16]. In addition, habitual coffee drinking has been associated with the prevention of diseases including cancer, cardiovascular disorders, obesity and diabetes as well as neurodegenerative disorders [15-20].

Cardamom (*Elettaria cardamomum* Maton), the Queen of all spices has a history as old as human race [21]. The major use of Cardamom on worldwide is for domestic culinary purpose and in medicine. The aroma and medicinal properties of cardamom are due to the volatile oil present in it and it is obtained from the seeds by steam distillation. The composition of cardamom oil has been studied by various workers and the major compounds found were 1, 8 cineole (20-60%) and α -terpinyl acetate (20-55%) [21]. Studies have implicated cardamom's potential

therapeutic value as an inhibitor of human platelet aggregation [22] and it has antihypertensive, gastroprotective, anticancer and antidiabetic properties [23]. Cardamom has antioxidant properties and can increase levels of glutathione and antioxidant enzymes in the body [24,25]. Reports are available on the antioxidant activity of extract of cardamom from different countries [26-30].

Due to oxidative stress induced by γ -irradiation exposure and the occurrence of different tissue damage, this study aims to evaluate ameliorating effects of coffee and cardamom mixture against biochemical changes induced by γ -irradiation.

Material and Methods

Materials

Coffee and cardamom powder as well as standard commercial rodent diet were purchased from local herbal market (Cairo, Egypt).

Preparation of coffee and cardamom mixture

The mixture of coffee and cardamom were prepared by adding 5 g of cardamom powder for 100 g of mixture (95 g coffee + 5 g cardamom).

Extraction and determination of phenolic compounds of coffee

Grounded dry powder of coffee (10 g) was weighed into a test tube. A total of 100 ml of 80% aqueous methanol was added, and the

*Corresponding author: Refaat G. Hamzaa, Department of Food Irradiation, Atomic Energy Authority, 3 Ahmed El Zomor St., El Zohour District, 8th District, Nasr City, Cairo, Tel: +201224109349; Fax: +20222749298; E-mail: Refaat.galal2009@yahoo.com

Received July 06 2012; Accepted July 19, 2012; Published July 30, 2012

Citation: Hamzaa RG, Osman NN (2012) Using of Coffee and Cardamom Mixture to Ameliorate Oxidative Stress Induced in γ -irradiated Rats. Biochem Anal Biochem 1:113. doi:10.4172/2161-1009.1000113

Copyright: © 2012 Hamzaa RG, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

suspension was stirred slightly. Tubes were sonicated twice for 15 min and one left at room temperature ($\sim 20^{\circ}\text{C}$) for 24 h. The extract was centrifuged for 10 min (10 min, 1500 x g), and supernatants were filtered through a 0.2 μm Millipore membrane filter then 1-3 ml was collected in vial for the HPLC analysis of phenolic compounds. According to the method of Goupy et al. [31], phenolic compounds of coffee were analyzed with high performance liquid chromatography (Hewlett Packard series 1050) equipped with autosampling injector, solvent degasser, ultraviolet (UV) detector set at 280 nm and quarter HP pump (series 1050). The column temperature was maintained at 35°C . Gradient separation was carried out with methanol and acetonitrile as a mobile phase at flow rate of 1 ml/min. The phenolic standards from Sigma Co. were dissolved in a mobile phase and injected into HPLC. Retention time and peak area were used to calculation of Phenolic compounds concentration by the data analysis of HEWLETT Packard Software.

Essential oil determination

Separation and identification of essential oil components were performed by using Gas chromatography instrument, Model Hewlett-Packard- MS (5970) series II at the Agriculture Research Center, Giza, Egypt. Condition analysis as follows: Column: 30 m hp Methyl silicon 0.1 mm; Temperature: Initial 60°C ; Rate: $3^{\circ}\text{C} / \text{min}$ up to 240°C ; Carrier gas: Helium 1.0 ml/min; Injection port; Temperature: 250°C ; Detector temperature: 270°C ; Integration: By using HP software Data; Injection volume: 0.3 ml. The isolated peaks were identified by matching with data from the library of mass spectra and compared to those of authentic compounds and published data [32]. Quantitative determination was carried out based on peak area integration.

Radiation facility

Whole body gamma irradiation at dose level of 6 Gy was performed using a Canadian gamma cell-40, (137 Cs) housed at the National Center for Radiation Research and Technology (NCRRT), Cairo, Egypt. The dose rate was 0.44 Gy/min at the time of the experiment.

Experimental animals

Adult male albino rats, reared in NCRRT animal house, were used in the present experiments. Matched weight animals (150 ± 10 g) were selected and housed in plastic cages under controlled condition and fed on standard commercial rodent diet. Gamma irradiated rats were orally received extract of coffee and cardamom mixture at dose level (60 mg/100 g body weight/day) [20].

Experimental design

Animals (28 rats) were randomly divided into 4 groups each of six animals as follows:

Group A: Rats fed on balanced diet for 8 weeks, served as control.

Group B: Rats fed on balanced diet and received orally the extract of coffee and cardamom mixture at a dose level of 60 mg/100 g body weight for 8 weeks, served as positive control.

Group C: Rats fed daily on balanced diet for 8 weeks and exposed to γ -irradiation (6 GY) at the 4th week, served as negative control.

Group D: Rats fed daily on balanced diet and received orally the extract of coffee and cardamom mixture for 8 weeks and exposed to γ -irradiation at the 4th week.

At the end of the experiment, animals from each group were

sacrificed 24 hrs after the last dose of treatment. Blood sample were collected through heart puncture after light anesthesia and allowed to coagulate and centrifuged to obtain serum for biochemical analysis. Also, liver was removed for biochemical investigation.

Biochemical analysis

The lipid peroxidation was determined colorimetrically as malondialdehyde (MDA) [33]. Hepatic glutathione content (GSH) and activity of superoxides dismutase (SOD) were measured by the method of Gross et al. [34] and Minami and Youshikawa [35], respectively. Serum glucose was evaluated by the method of Trinder [36]. The activity of serum aspartate transaminase (AST) and alanine transaminase (ALT) were assayed by the method of Reitman and Frankel [37], as well as serum gamma glutamyl transferase (GGT) was assessed according to Rosalk, [38]. In addition, total cholesterol (TC), triglycerides (TG) and high-density lipoprotein-cholesterol (HDL-C) were determined according to procedure described by Allain et al. [39], Fossati and Prencipel, [40] and Demacker et al. [41], respectively while low-density lipoprotein cholesterol and very Low-density lipoprotein-cholesterol were evaluated according to Friedwald et al. [42] and Norbert [43] formulas, respectively by the following equations: $\text{LDL-C (mg/dl)} = \text{TC} - (\text{TG}/5 + \text{HDL-C})$, $\text{vLDL (mg/dl)} = \text{TG}/5$. Finally, the serum testosterone was analyzed by the enzyme linked immunosorbent assay (ELISA) according to the method of Engrall and Perlman, [44] and also insulin hormone was determined by radioimmunoassay kit supplied by Diasari, Italy.

Statistical analysis

Statistical analyses were performed using computer program Statistical Packages for Social Science [45] and values compared with each other using suitable tests.

Results

The phenolic compounds of coffee were identified by HPLC against standard compounds and the results were listed in Table 1. The most abundant components of coffee were caffeine, pyrogallol and caffeic "820.59, 65.23 and 17.98 mg/100g", respectively.

The essential oil contents found in cardamom were analyzed and the results showed that the main essential oil components were terpenyl acetate (60.65%) and 1,8 Cineol (13.63%) (Table 2).

Results of irradiated rats revealed a significant increase in the level of MDA and a remarkable decrease in the content of GSH and SOD activity while irradiated rats received oral extract of coffee and cardamom mixture had a lower level of MDA accompanied by

Phenolic compounds	mg/100g
Pyrogallol	65.23
Gallic	4.40
Protocatechuic	8.15
Catechin	8.67
Caffeic	17.98
Caffeine	820.59
Ferulic	2.40
P.coumaric	2.83
Ellagic	15.42
Cholchecien	7.32
Salicylic	16.69
Cinnamic	5.50

Table 1: Phenolic compounds of coffee (mg/100g).

Essential oil	%
α -Pinene	1.36
Sabimene	1.08
Limonene	7.12
1,8 Cineol	13.63
γ -Terpinene	0.25
Terpinolene	0.31
Linaiol	0.92
Terpineol	5.10
Octyl acetate	0.05
Linalyl acetate	2.33
Geraniol	0.57
Terpenyl acetate	60.65
Neryl acetate	2.11
Total unknown	4.52

Table 2: Essential oil contents of cardamom.

Parameters	A	B	C	D
MDA (n mol/ml)	178.71 \pm 0.82 ^c	166.46 \pm 1.03 ^d	396.07 \pm 1.52 ^a	211.26 \pm 0.85 ^b
GSH (mg/g tissue)	24.62 \pm 0.35 ^a	25.58 \pm 0.46 ^a	15.48 \pm 0.22 ^c	22.18 \pm 0.41 ^b
SOD (U/mg protein)	43.60 \pm 0.65 ^a	43.86 \pm 0.71 ^a	36.65 \pm 0.47 ^c	40.96 \pm 0.47 ^b

Values are expressed as means \pm S.E. (n=6).

Values in the same raw with different superscripts are differing significantly at P<0.05.

Table 3: Effect of coffee and cardamom mixture extract administration on MDA, GSH and SOD activity of γ -irradiated rats.

increasing in GSH level and the activity of SOD (Table 3).

On the other hand, the data revealed a significant elevation in AST, ALT and GGT activity accompanied by high glucose level in irradiated group compared to control group. Irradiated rats treated with the mixture extract had a significant reduction in the activity of the above mentioned liver enzymes and glucose concentration (Table 4).

Also, results in Table 5 indicated that exposure of rats to γ -irradiation resulted in a significant increase in TC, TG, LDL-C and vLDL-C concentration and decrease in the level of HDL-C. However, administration of coffee and cardamom mixture extract induced moderate reduction in the bad effect obtained due to whole body γ -irradiation.

The reducing effect of γ -irradiation on the level of testosterone and insulin were obtained in the Figures 1 & 2, while it was noticed that the treatment by using coffee and cardamom mixture to eliminate the adverse effects of γ -irradiation pointed to an obvious enhancement in the level of the both hormones.

Discussion

Exposure to ionizing radiation might happen as a consequence of military or industrial accidents, or after therapeutic treatments [46,47]. Also, it is well known that ionizing radiation generates reactive oxygen species (ROS) that interact with cellular molecules, including DNA, lipids, and proteins [48].

Coffee polyphenolic components (cafestol, kehweol, cholinergic acid and caffeine) [49], as well as essential oil of cardamom [21] function as an antioxidant by scavenging reactive oxygen species and have chemoprotective and anti-inflammatory property [26,50] reverse lipid peroxidation, enzymatic leakage and enhance cellular antioxidant defense mechanism, reported by Dreosti et al. [51].

Parameters	A	B	C	D
AST U/ml	27.57 \pm 0.49 ^c	26.09 \pm 0.43 ^c	26.09 \pm 0.43 ^c	32.35 \pm 0.29 ^b
ALT U/ml	15.92 \pm 0.05 ^c	15.16 \pm 0.16 ^c	47.49 \pm 0.54 ^a	29.31 \pm 0.55 ^b
GGT U/ml	3.24 \pm 0.01 ^c	3.20 \pm 0.02 ^c	6.13 \pm 0.02 ^c	4.46 \pm 0.04 ^b
Glucose (mg/dl)	112.49 \pm 1.34 ^c	109.46 \pm 0.68 ^c	195.95 \pm 1.69 ^a	126.00 \pm 0.96 ^b

Values are expressed as means \pm S.E. (n=6).

Values in the same raw with different superscripts are differing significantly at P<0.05.

Table 4: Effect of coffee and cardamom mixture extract administration on the activity of some liver enzymes and glucose level of γ -irradiated rats.

Parameters	A	B	C	D
TC mg/dl	152.48 \pm 0.83 ^c	149.45 \pm 1.09 ^c	208.13 \pm 1.26 ^a	186.42 \pm 1.17 ^b
TG mg/dl	112.47 \pm 0.91 ^d	108.41 \pm 0.90 ^c	181.45 \pm 1.06 ^a	156.50 \pm 0.86 ^b
HDL-C mg/dl	45.15 \pm 0.72 ^a	47.44 \pm 0.65 ^b	37.53 \pm 0.82 ^d	41.35 \pm 0.51 ^c
LDL-C mg/dl	84.84 \pm 0.22 ^d	80.32 \pm 0.35 ^c	134.31 \pm 0.44 ^a	113.77 \pm 0.76 ^b
vLDL-C mg/dl	22.49 \pm 0.18 ^c	21.68 \pm 0.18 ^c	36.29 \pm 0.21 ^a	31.30 \pm 0.17 ^b

Values are expressed as means \pm S.E. (n=6).

Values in the same raw with different superscripts are differing significantly at P<0.05.

Table 5: Effect of coffee and cardamom mixture extract administration on lipid profile of γ -irradiated rats.




Figure 1: Effect of administration of mixture extract on testosterone level of γ -irradiated rats.




Figure 2: Effect of administration of mixture extract on insulin level of γ -irradiated rats.

Results in the present study demonstrated a significant increase in lipid peroxidation (MDA) and reduction in GSH concentration and SOD activity due to exposure of rat to γ -irradiation (6 Gy). The results are in agreement with those recorded by several authors [52-54]. They recorded a significant depletion in the antioxidant system accompanied by enhancement of lipid peroxides after whole body gamma-irradiation. This could be due to an enhanced utilization of the antioxidant system as an attempt to detoxify the free radicals generated by radiation exposure.

On the other hand, results of gamma irradiated rats received oral extract of coffee and cardamom mixture obtained a significant low level of liver MDA and high GSH concentration and SOD activity in comparison with irradiated group. Therefore, these results reflect the protective effects of both coffee and cardamom against the damage effect of γ -irradiation. Abreu et al., [55] found that the consumption of coffee decreased the lipid peroxidation of the membranes; increased the concentration of glutathione, a potent endogenous antioxidant; and increased the activity of two antioxidant enzymes, glutathione reductase and superoxide dismutase. Undoubtedly, these beneficial effects of coffee on antioxidant properties are partly due to caffeine, kahweol and cafestol [56-58]. In addition to the effect of coffee, cardamom also have a potential antioxidants such as 1, 8-ceineoil and alphaterpineol, protocatechualdehyde and protocatechuic acid which have health benefits by inhibiting lipid peroxidation and activation of antioxidant enzymes [24,25].

The effect of γ -irradiation on liver function in this study were determined by detection of AST, ALT and GGT activity which were highly increased under the effect of γ -irradiation, while the increase in these enzymes significantly reduced by administration of the extract of coffee and cardamom mixture to irradiated rats. Also, the results indicted the increasing effect of γ -irradiation exposure in the level of serum glucose which highly declined after treatment with the mixture extract.

The significant increase of serum transaminases activity may be attributed to a radiation induced state of hypoxia in the parenchymal cells of contracting fibrous tissue as well as to extensive body tissue breakdown includes liver parenchyma and renal tubules and that can cause change in tissue permeability and could enhance the release of the transaminase enzymes from their subcellular sites of production to extracellular and consequently to the blood circulation [59,60]. On the other hand, the increased glucose level might be related to endocrine abnormalities induced by irradiation that promote the secretion of biologically active peptide which has relation carbohydrate metabolism by increasing gluconeogenesis in liver [61].

The effect of both coffee and cardamom resulted in significant enhancement in liver function and serum glucose level by reducing the damage effect induced by γ -irradiation. The negative association between coffee and the development of increased serum AST, ALT and GGT activity might be to some extent to caffeine, which is one of the major ingredients contained in coffee with various biological actions and possibly plays a crucial role in the observed associations of coffee intake [63]. Also, the coffee constituent chlorogenic acid may have the potential to influence glucose metabolism processes to prevent hyperglycemia by inhibition activity of glucose-6-phosphatase, an important regulator of blood glucose levels [63,64].

As well as, Abdel-Wahab and Aly [65] observed that treatment with clove and cardamom effectively decreased liver enzyme levels in the serum. This can be attributed to the presence of antioxidant

in clove and cardamom which contain phenolic compounds that can act by scavenging free radicals. Also, the lowering effect of cardamom on glucose level was observed by El-Yamani [23], who studied the hypoglycemic effect of cardamom as one of spices that possess antioxidant compounds.

The present study showed a significant increase in cholesterol, triglycerides and LDL-C levels associated with reduction in the concentration of HDL after exposure to 6 Gy irradiation compared to control group. In contrast, a moderate decrease in TC, TG and LDL-C contents and rise in HDL level were observed in group D. El-Khafif et al. [66] and Mansour [67] reported that whole body exposure to gamma radiation induced high level of serum cholesterol fractions due to its release from tissues, destruction of cell membranes and increase rate of cholesterol biosynthesis in the liver and other tissues. Also, irradiation could modify low and high density lipoproteins metabolism indirectly through the action of various inflammatory products. Gamma irradiation might decrease the lipoprotein lipase activity in adipose tissue, leading to reduction in the uptake of triacylglycerols.

Choi et al. [20] reported that the cholesterol-raising effect of coffee has long been controversial, and observe significant increase in TC, decrease in HDL-C, and no effect on plasma TG levels after coffee intake (10 cups/day) and the authors attributed this cholesterol-raising effect of coffee to its contents of diterpenes cafestol and kahweol. Therefore, the moderate effect of coffee and cardamom mixture intake on lipid profile in this study was related to the effect of cardamom. The cholesterol and triglyceride lowering effect of cardamom reflecting its protective hepatocellular effects and the ability of its phenolic contents and essential oil to reduce the hyperlipidemia [68-70]. Furthermore, it has been reported that spices may inhibit hepatic 3-hydroxyl-3-methylglutaryl coenzyme A reductase activity, resulting in lowering hepatic and serum cholesterol levels [71].

It was well clear that gamma irradiation induced a significant decrease in testosterone and insulin hormones concentration and these results agreed with those of Michael and Amer [2] and Liu et al. [72] who recorded decreased testosterone level after whole-body irradiation dose of 4 and 5 Gy due to alterations in DNA-single strand break, cell apoptosis and oxidative stress. Also, Popoff and Kapich [73] found a positive correlation between a decline in testosterone affinity and exposure to gamma irradiation. In addition, the lowering effect of γ -irradiation exposure on insulin level was observed due to production of free radicals that induced oxidative stress resulted in reduction in insulin secretion and DNA damage [3].

This study reported that coffee and cardamom ingestion to irradiated rats elevates testosterone secretion. When cardamom is combined with coffee it has especially strong effects on the genital reactions and sexual function and it was traditionally used in Arabic coffee for this purpose. Sanovane et al. [74] reported that cardamom intake stimulates the nervous system which in turn stimulates the pituitary gland to increase the secretion of hormones induced gonads to produce testosterone. In addition, previous animal experiments had found that coffee infusion elevated plasma concentrations of testosterone and that related to the effect of bioactive contents of coffee [75].

Also, coffee and cardamom administration reduced the damage effect due to γ -irradiation exposure and enhance the elevation of insulin concentration in the serum of γ -irradiated rats. Effects of coffee and cardamom were related to their antioxidant contents such as caffeine and terpenyl acetate which reduced oxidative stress by scavenging free radicals and stimulate pancreatic cells to secrete insulin [21,76].

Conclusion

The observed improvement in the antioxidants, liver functions, lipid profile and level of testosterone and insulin in irradiated rats suggested that the radioprotective effect of the mixture of coffee and cardamom was attributed to the phenolic compounds and essential oil contents which possess antioxidant potential can reduce the oxidative stress induced by γ -irradiation exposure.

References

1. Maurya DK, Devasagayam TP, Nair CK (2006) Some novel approaches for radioprotection and the beneficial effect of natural products. *Indian J Exp Biol* 44: 93-114.
2. Michael MI, Amer MM (2010) Is the use of antioxidants mixture ameliorating the double stress effect of radiation and hyperlipidemia? *J Rad Res Appl Sci* 3: 855-874.
3. Lee TK, O'Brien KF, Wang W, Johnke R M, Sheng C, et al. (2010) Radioprotective effect of American ginseng on human lymphocytes at 90 minutes postirradiation: a study of 40 cases. *J Altern Complement Med* 16: 561-567.
4. Batra V, Devasagayam TP (2009) Interaction between cytotoxic effects of gamma-radiation and folate deficiency in relation to choline reserves. *Toxicology* 255: 91-99.
5. Batra V, Sridhar S, Devasagayam TP (2010) Enhanced one-carbon flux towards DNA methylation: Effect of dietary methyl supplements against gamma-radiation-induced epigenetic modifications. *Chem Biol Interact* 183: 425-433.
6. Jagetia GC, Baliga MS (2002) Influence of the leaf extract of *Mentha arvensis* Linn. (mint) on the survival of mice exposed to different doses of gamma radiation. *Strahlenther Onkol* 178: 91-98.
7. Maharwal J, Samarth RM, Saini MR (2003) Radiomodulatory influence of Rajgira (*Amaranthus paniculatus*) leaf extract in Swiss albino mice. *Phytother Res* 17: 1150-1154.
8. Arora R, Gupta D, Prasad J, et al (2003a) Bioprospection of Himalayan highaltitude region flora for radioprotective biomolecules. In Proceedings of the National Seminar on New Millenium Strategies for Quality, Safety and GMPs for Herbal Drugs/Products, NBRI, Lucknow, India.
9. Gupta D, Arora R, Gupta S, Gupta S, Garg AP, Goel HC (2003) Radiationinduced oxidative damage to mitochondria and its modification by endogenous antioxidants in vivo: role of some herbal preparations. *Indian J Nuclear Med* 18: 14-18.
10. Arora R, Gupta D, Sharma AK, et al (2004) Modification of radiationinduced damage in mammalian model systems by natural plant products: implications for radiation protection. In Proceedings of the IUPAC International Conference on Biodiversity and Natural Products, Delhi University, Delhi, India; 33.
11. Hostettmann K, Marston A, Ndjoko K, Wolfender JL (2000) The Potential of African plants as a source of drugs. *Current Organic Chemistry* 4: 973-1010.
12. Nardini M, Cirillo E, Natella F, Scaccini C (2002) Absorption of phenolic acids in humans after coffee consumption. *J Agric Food Chem* 50: 5735-5741.
13. Arts IC, van De Putte B, Hollman PC (2000) Catechin contents of foods commonly consumed in the Netherlands. 2. Tea, wine, fruit juices, and chocolate milk. *J Agric Food Chem* 48: 1752-1757.
14. Clifford MN, Knight S (2004) The cinnamoyl-amino acid conjugates of green robusta coffee beans. *Food Chem* 87: 457-463.
15. Kuriyama S, Shimazu T, Ohmori K, Kikuchi N, Nakaya N, et al. (2006) Green tea consumption and mortality due to cardiovascular disease, cancer, and all causes in Japan: the Ohsaki study. *JAMA* 296: 1255-1265.
16. Thielecke F, Boschmann M (2009) The potential role of green tea catechins in the prevention of the metabolic syndrome - a review. *Phytochemistry* 70: 11-24.
17. Ogunleye AA, Xue F, Michels KB (2010) Green tea consumption and breast cancer risk or recurrence: a meta-analysis. *Breast Cancer Res Treat* 119: 477-484.
18. Hu G, Bidel S, Jousilahti P, Antikainen R, Tuomilehto J (2007) Coffee and tea consumption and the risk of Parkinson's disease. *Mov Disord* 22: 2242-2248.
19. Eskelinen MH, Ngandu T, Tuomilehto J, Soininen H, Kivipelto M (2009) Midlife coffee and tea drinking and the risk of late-life dementia: a population-based CAIDE study. *J Alzheimers Dis* 16: 85-91.
20. Choi EY, Park SY, Cho YO (2011) Freeze-dried instant coffee can promote the activities of antioxidant enzymes and induce weight loss but also aggravate the plasma cholesterol profile in rats. *Nutrition* 27: 1202-1205.
21. Padmakumari Amma KP, Rani MP, Sasidharan I, Premakumari Nisha VN (2010) Chemical Composition, Flavonoid - Phenolic Contents and Radical Scavenging Activity of Four Major Varieties of Cardamom. *Int J Biol Med Res* 1: 20-24.
22. Sunnetha WJ, Krishnakantha TP (2005) Cardamom extract as inhibitor of human platelet aggregation. *Phytother Res* 19: 437-440.
23. El-yamani MA (2011) Cinnamon, cardamom and ginger impact as evaluated on hyperglycemic rats. *J. of specific education* 20: 664-679.
24. Verma SK, Rajeevan V, Bordia A, Jain V (2010) Greater cardamom (*Amomum subulatum* Roxb.) - A cardio-adaptogen against physical stress. *J Herb Med Toxicol* 4: 55-58.
25. Bisht VK, Negi J S, Bhandari A K, Sundriyal RC (2011) *Amomum subulatum* Roxb: Traditional, phytochemical and biological activities-An overview. *Afr J Agric Res* 6: 5386-5390.
26. Hossain MB, Brunton NP, Barry-Ryan C, Martin-Diana AB, Wilkinson M (2008) ANTIOXIDANT ACTIVITY OF SPICE EXTRACTS AND PHENOLICS IN COMPARISON TO SYNTHETIC ANTIOXIDANTS. *Rasayan J Chem* 4: 751-756.
27. Hinneburg I, Damien Dorman HJ, Hiltunen R (2006) Antioxidant activities of extracts from selected culinary herbs and spices. *Food Chemistry* 97: 122-129.
28. Nooman AK, Ashok KS, Atif AO, Zaha EA, Husni F (2008) Antioxidant Activity of Some Common Plants. *Turk J Biol* 32: 51-55.
29. Souri E, Amin G, Farsam H, Barazandeh TM (2008) Screening of antioxidant activity and phenolic content of 24 medicinal plant extracts. *DARU* 16: 83-87.
30. Lilian Regina BM, Gisela Pizarro dE MB, Neura B, Adriana Zerlotti M (2008) Free radical scavenging activity of ethanolic extracts from herbs and spices commercialized in Brazil. *Braz arch biol technol* 51: 1225-1232.
31. Goupy P, Hugues M, Boivin P, Amiot MJ (1999) Antioxidant compounds of barley (*Hordeum vulgare*) and malt extracts. *J Sci of Food and Agri* 79: 1625-1634.
32. Adams RP (1995) Identification of essential oil components by gas chromatography/mass spectrometry. (4th edn) Allured Publ Corp. USA.
33. Yoshioka T, Kawada K, Shimada T, Mori M (1979) Lipid peroxidation in maternal and cord blood and protective mechanism against activated-oxygen toxicity in the blood. *Am J Obstet Gynecol* 135: 372-376.
34. Gross RT, Bracci R, Rudolph N, Schroeder E, Kochen JA (1967) Hydrogen peroxide toxicity and detoxification in the erythrocytes of newborn infants. *Blood* 29: 481-493.
35. Minami M, Yoshikawa H (1979) A simplified assay method of superoxide dismutase activity for clinical use. *Clin Chim Acta* 92: 337-342.
36. Trinder P (1969) Enzymatic colorimetric determination of glucose. *Ann Clin Biochem* 6,24. C/F, Bio Merieux Comp, France.
37. Rosalk SB (1967) In advances clinical chemistry. Academic Press Chap17: 53.
38. REITMAN S, FRANKEL S (1957) A calorimetric method for determination of serum glutamic oxalacetic and glutamic pyruvic transaminases. *Am J Clin Pathol* 28: 56-63.
39. Allain CC, Poon LS, Chan CS, Richmond W, Fu PC (1974) Enzymatic determination of total serum cholesterol. *Clin Chem* 20: 470-475.
40. Fossati P, Prencipe L (1982) Serum triglycerides determined colorimetrically with an enzyme that produces hydrogen peroxide. *Clin Chem* 28: 2077-2080.
41. Demacker P, Vos-janssen H, Hifman A, Vants Lear A, Jansen A (1980) Messurement of high density lipoprotein cholesterol in serum. *Clin Chem* 26: 1780.
42. Friedewald WT, Levy RI, Fredrickson DS (1972) Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem* 18: 499-502.
43. Norbert WT (1995) *Clinical Guide to Laboratory Tests* (Third ed), Saunders W.B. Company, Philadelphia.

44. Engrall E, Pelmann P (1971) Qualitative enzyme linked immunoassay. *Immunochem* 8: 871.
45. SPSS (1998) Statistical Package for Social Science. Computer Software, Ver. 10. SPSS Company, London, UK.
46. Guelman LR, Cabana JI, del Lujan Pagotto RM, Zieher LM (2005) Ionizing radiation-induced damage on developing cerebellar granule cells cultures can be prevented by an early amifostine post-treatment. *Int J Dev Neurosci* 23: 1-7.
47. Lee WH, Sonntag WE, Lee YW (2010) Aging attenuates radiation-induced expression of pro-inflammatory mediators in rat brain. *Neurosci Lett* 476: 89-93.
48. Tominaga H, Kodama S, Matsuda N, Suzuki K, Watanabe M (2004) Involvement of reactive oxygen species (ROS) in the induction of genetic instability by radiation. *J Radiat Res* 45: 181-188.
49. Son S, Lewis BA (2002) Free radical scavenging and antioxidative activity of caffeic acid amide and ester analogues: structure-activity relationship. *J Agric Food Chem* 50: 468-472.
50. Sud'ina GF, Mirzoeva OK, Pushkareva MA, Korshunova GA, Sumbatyan NV, et al. (1993) Caffeic acid phenethyl ester as a lipooxygenase inhibitor with antioxidant properties. *FEBS Lett* 329: 21-24.
51. Dreosti IE, Wargovich MJ, Yang CS (1997) Inhibition of carcinogenesis by tea: the evidence from experimental studies. *Crit Rev Food Sci Nutr* 37: 761-770.
52. Bhatia AL, Jain M (2004) Spinacia oleracea L. protects against gamma radiations: a study on glutathione and lipid peroxidation in mouse liver. *Phytomedicine* 11: 607-615.
53. Koc M, Taysi S, Buyukokuroglu ME, Bakan N (2003) Melatonin protects rat liver against irradiation-induced oxidative injury. *J Radiat Res* 44: 211-215.
54. Smarth RM, Kumar A (2003) Radioprotection of Swiss albino mice by plant extract *Mentha piperita* (Linn.). *J Radiat Res* 44:101-109.
55. Abreu RV, Silva-Oliveira EM, Moraes MF, Pereira GS, Moraes-Santos T (2011) Chronic coffee and caffeine ingestion effects on the cognitive function and antioxidant system of rat brains. *Pharmacol, Biochem Behav* 99: 659-664.
56. Huber WW, Scharf G, Nagel G, Prustomersky S, Schulte-Hermann R, et al. (2003) Coffee and its chemopreventive components Kahweol and Cafestol increase the activity of O6-methylguanine- DNA methyltransferase in rat liver--comparison with phase II xenobiotic metabolism. *Mutat Res* 522: 57-68.
57. Inkielewicz-Stepniak I, Czarnowski W (2010) Oxidative stress parameters in rats exposed to fluoride and caffeine. *Food Chem Toxicol* 48: 1607-1611.
58. Nobre HV Jr, Cunha GM, de Vasconcelos LM, Magalhães HI, Oliveira Neto RN, et al (2010) Caffeine and CSC, adenosine A2A antagonists, offer neuroprotection against 6-OHDA-induced neurotoxicity in rat mesencephalic cells. *Neurochem Int* 56: 51-58.
59. Ashry O M (2003) Modulation of radiation induced toxicity by caffeine pre injection in Rats. *Egypt. J Rad Sce Appl* 16: 1-11.
60. Ramadan LA, Roushdy HM, Abu Senna GM, Amin NE, El-Deshw OA (2002) Radioprotective effect of silymarin against radiation induced hepatotoxicity. *Pharmacol Res* 45: 447-454.
61. Kafafy YA, Roushdy HML, Ashry OM, Salama SF, Abdel-Halim M, et al. (2005) Green tea Antioxidative Potential in Irradiated Pregnant rats. *Egyptian Journal of Radiation Sciences and Applications* 18: 313-333.
62. Thakur S, Du J, Hourani S, Ledent C, Li JM (2010) Inactivation of Adenosine A2A Receptor Attenuates Basal and Angiotensin II-induced ROS Production by Nox2 in endothelial cells. *J Biol Chem* 285: 40104-40113.
63. Chou JY, Matern D, Mansfield BC, Chen YT (2002) Type I glycogen storage diseases: disorders of the glucose-6-phosphatase complex. *Curr Mol Med* 2: 121-143.
64. McCarty MF (2005) A chlorogenic acid-induced increase in GLP-1 production may mediate the impact of heavy coffee consumption on diabetes risk. *Med Hypotheses* 64: 848-853.
65. Abdel-Wahhab MA, Aly SE (2005) Antioxidant property of *Nigella sativa* (black cumin) and *Syzygium aromaticum* (clove) in rats during aflatoxicosis. *J Appl Toxicol* 25: 218-223.
66. El-Khafif M, Ragab M, El-Dawy H, Tawfik S (2003) Effect of taurine treatment on some biochemical parameters in gamma irradiated mice. *Environ Sci* 6: 393-401.
67. Mansour HH (2006) Protective role of carnitine ester against radiation-induced oxidative stress in rats. *Pharmacol Res* 54: 165-171.
68. Kapoor IPS, Singh B, Singh G, Isidorov V, Szczepaniak L (2008) Chemistry antifungal and antioxidant activities of cardamom (*Amomum subulatum*) essential oil and oleoresins. *International Journal of Essential Oil Therapeutics* 2: 29-40.
69. EL-Segay O, Ab-Alla A, Abu- Al-Nman S (2007) Experimental study of antioxidant and Hepatoprotective effects of clove and cardamom in ethanol induced hepatotoxicity. *Tanta Medical Sciences Journal* 2: 27-36.
70. Sadeek EA, Abd El-Razek FH (2010) The Chemo-Protective Effect of Turmeric, Chili, Cloves and Cardamom on Correcting Iron Overload-Induced Liver Injury, Oxidative Stress and Serum Lipid Profile in Rat Models. *J am Sci* 6.
71. Shobana S, Naidu KA (2000) Antioxidant activity of selected Indian spices. *Prostaglandins Leukot Essent Fatty Acids* 62: 107-110.
72. Liu Y, Zhang H, Zhang L, Zhang X, Xie Y, et al. (2009) Melatonin modulates acute testicular damage induced by carbon ions in mice. *Pharmazie* 64: 685-689.
73. Popoff EH, Kapich AN (2010) The effect of ionising radiation on testosterone binding globulin characteristics: correction of the protein' parameters by lipid polyene complexes of fungus *Laetiporus sulfureus*. *Int J Radiat Biol* 86: 238-251.
74. Beaven CM, Hopkins WG, Hansen KT, Wood MR, Cronin JB, et al. (2008) Dose effect of caffeine on testosterone and cortisol responses to resistance exercise. *Int J Sport Nutr Exerc Metab* 18: 131-141.
75. Sonavane G, Sarveiya V, Kasture V, Kasture SB (2001) BEHAVIOURAL ACTIONS OF MYRISTICA FRAGRANS SEEDS. *Indian J Pharmacol* 33: 417-424.
76. Kempf K, Herder C, Erlund I, Kolb H, Martin S, et al. (2010) Effects of coffee consumption on subclinical inflammation and other risk factors for type 2 diabetes: a clinical trial. *Am J Clin Nutr* 91: 950-957.